



**HAL**  
open science

# Designing innovative productive cropping systems with quantified and ambitious environmental goals

Caroline Colnenne-David, Thierry Doré

► **To cite this version:**

Caroline Colnenne-David, Thierry Doré. Designing innovative productive cropping systems with quantified and ambitious environmental goals. *Renewable Agriculture and Food Systems*, 2015, 30 (06), pp.487-502. 10.1017/S1742170514000313 . hal-01354945

**HAL Id: hal-01354945**

**<https://agroparistech.hal.science/hal-01354945>**

Submitted on 7 Sep 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 doi:10.1017/S1742170514000313

2

3 **Designing innovative productive cropping systems with quantified and ambitious**  
4 **environmental goals**

5

6 Colnenne-David Caroline<sup>1</sup>, Doré Thierry<sup>2</sup>

7 <sup>1</sup> INRA, UMR 211 Agronomie, 78850 Thiverval-Grignon, France, e-mail:

8 caroline.colnenne@grignon.inra.fr

9 <sup>2</sup> AgroParisTech, UMR 211 Agronomie, 78850 Thiverval-Grignon, France e-mail:

10 thierry.dore@agroparistech.fr

11

12

13

14

15

16

17

18

19

20

21

22

23 Corresponding author: COLNENNE-DAVID Caroline

24 INRA, UMR 211 Agronomie, 78850 Thiverval-Grignon, France

25 Phone: (+33) 1.30.81.53.28

26 Fax: (+33) 1.30.81.54.25

27 e-mail: caroline.colnenne@grignon.inra.fr

28

## 29 **Abstract**

30 Agriculture must face a number of very pressing environmental issues. We used the prototyping method to design three  
31 innovative cropping systems, each satisfying three ambitious goals simultaneously: (i) overcoming a major  
32 environmental constraint, which represents a major break regarding objectives to be reached in current cropping  
33 systems (differing between systems: a ban on all pesticides but with chemical N fertilizer permitted; reducing fossil  
34 energy consumption by 50%; or decreasing greenhouse gas emissions by 50%), (ii) meeting a wide range of  
35 environmental criteria, and (iii) maximizing yields, given the major constraint and environmental targets. A fourth  
36 cropping system was designed, in which the environmental and yield targets were achieved with no major constraint  
37 (PHEP system). The performances of these innovative cropping systems were compared to a conventional system in the  
38 Ile-de-France region. We used a three-step prototyping method: (1) new cropping systems were designed on the basis of  
39 scientific and expert knowledge, (2) these system prototypes were assessed with tools and a model (*ex ante* assessment)  
40 adjusted to the set of constraints and targets, with optimization by an iterative process until the criteria were satisfied  
41 and (3) evaluation in a long-term field experiment (*ex post* assessment), which is currently underway. We describe only  
42 the first two steps here, together with the results of the prototypes assessment with tools and a model. The pesticide,  
43 energy and greenhouse gas constraints were fulfilled. All these innovative systems satisfied environmental criteria in  
44 terms of nitrogen and phosphorus management, pesticide use, energy consumption and crop diversity. For the pesticide-  
45 free system, the soil organic matter indicator was lower than expected due to frequent plowing (every two years) and  
46 yields were 20 to 50% lower than for the PHEP system, depending on the crop considered. We focus our discussions on  
47 the design methodology and the availability of scientific knowledge and tools for projects of this type.

48

## 49 **Keywords**

50 Innovative cropping systems, environmental goals, *ex ante* assessment.

51

## 52 **Introduction**

53 Agriculturalists are faced with challenges relating to very pressing environmental and health issues, including the need  
54 to decrease pesticide use. In many countries, high levels of pesticides have frequently been found in rivers, lakes and  
55 groundwater<sup>1,2</sup>. A second pressing environmental issue is the consumption of fossil fuel. Energy use has markedly  
56 increased over the last decade<sup>3</sup>, and some scientists agree that oil availability will decline in the near future<sup>4</sup>, leading to  
57 a sharp increase in oil prices<sup>5</sup>. In this context, new ways of optimizing or reducing energy use have been proposed<sup>6</sup>.  
58 Global warming is a third challenging environmental issue facing agriculture. About 12% of global greenhouse gas  
59 (GHG) emissions emanate from agricultural lands<sup>7</sup>, and this proportion is expected to rise in the future, due to increases  
60 in the amount of land used for agricultural purposes and the intensification of agricultural practices<sup>8</sup>. Carbon (C)  
61 sequestration in the soil, through the return of crop residues, root deposition and organic amendments, may help to  
62 decrease GHG emissions<sup>9</sup>. Sustainable development is another pressing social issue. Sustainable agriculture must  
63 satisfy environmental criteria<sup>10</sup>. The harmful impact of agriculture on the environment can be lessened by optimizing  
64 fertilization (N, P) and increasing crop diversity. Currently pesticide use and energy consumption should also be  
65 reduced<sup>1,2</sup>, and soil fertility should be maintained. Since the 1950s, alternative crop management systems have been  
66 proposed<sup>11</sup> and legislation and inspection services have controlled the use of inputs (pesticides, N fertilization). Finally,  
67 agricultural production must satisfy the food needs of a soaring world population<sup>10</sup>. Global agricultural production  
68 currently feeds a population of approximately seven billion. Current projections suggest that the world population will  
69 have reached nine billion by 2050<sup>12</sup>. The resulting increase in the need for land for housing will reduce the amount of  
70 land available for agricultural purposes<sup>13</sup>. The availability of arable land per capita differs greatly between regions (*e.g.*,  
71 between China and South America), and major cropping systems must take this scarcity into account. Foley<sup>14</sup> have  
72 suggested that feeding a population of this size will be possible only if agricultural systems change, along with human  
73 eating habits.

74 Many studies in recent years have focused on the design and assessment of new cropping systems. New crop  
75 management strategies have been proposed to decrease pesticide use<sup>15</sup>, to decrease energy consumption<sup>16</sup>, or to enhance  
76 C sequestration through crop management practices<sup>17</sup>. Energy use and greenhouse gas emissions have been calculated  
77 and assessed for different systems<sup>18</sup>. At the cropping system level, long-term trials also have been set up to investigate  
78 the effects of different technical operations, such as N fertilization, on soil physical and chemical properties<sup>19</sup> and on  
79 soil biology<sup>20</sup>. Several studies have assessed the differences between organic, integrated and conventional cropping  
80 systems in terms of C sequestration<sup>21</sup>, energy efficiency and use<sup>22</sup>, profitability<sup>23</sup> or productivity<sup>24</sup>. Some authors have  
81 analyzed the impact of different degrees of tillage on productivity<sup>25</sup> or biological activity<sup>26</sup>. Others have focused on the  
82 effect of cropping systems on biodiversity<sup>27</sup>, or have used life cycle assessment methods to analyze the sustainability of

83 various farming systems<sup>28</sup>. In most of these cases, new cropping systems were designed by modifying a few agricultural  
84 practices targeting a single goal (*e.g.*, no chemicals to be used in organic cropping systems; no ploughing to increase C  
85 sequestration; more legumes in the rotation to reduce energy consumption; and more inputs to enhance profitability),  
86 without considering the other dimensions of sustainability. Despite the evidence that the future of agriculture must  
87 address a wide range of issues, no study has designed innovative cropping systems with specific and quantitative  
88 objectives covering a broad range of issues.

89 The objective of our project was to design, by prototyping<sup>29</sup>, innovative cropping systems meeting three quantitative  
90 objectives: (i) to satisfy a major environmental constraint, which represents a major break regarding objectives to be  
91 reached in current cropping systems (the banning of pesticide use, reducing fossil energy consumption by 50%, or  
92 reducing gas emissions by 50%); (ii) to satisfy a wide range of environmental criteria with specific quantitative targets,  
93 and (iii) to produce the maximum yield possible given the constraint and the environmental targets. The ultimate aim of  
94 this work is to improve arable cropping systems throughout northern Europe. Prototyping was developed to enable  
95 agronomists to design, test and improve more sustainable cropping systems<sup>29</sup>. With this approach, newly designed  
96 cropping systems could satisfy several of the issues mentioned above and contributed to identify the weaknesses of  
97 cropping systems. System prototypes were assessed with tools and a model (*ex ante* assessment), with discussion of  
98 their potential performances, before their assessment in a long-term field experiment (*ex post* assessment). We set up  
99 this long-term field experiment in 2008, and its results will be published in due course. We focus here on the  
100 prototyping and assessment of the cropping system prototypes. We discuss our results in terms of innovative design  
101 methodology, the innovation of agricultural practices, the availability of suitable tools, models and crop management,  
102 and the yields achievable.

103

## 104 **1. Materials and methods**

### 105 **1.1. Design method**

106 The method used for cropping system design was based on the prototyping approach<sup>29,30,31,32</sup>, which involves four major  
107 steps:

- 108 (i) Defining and ranking the constraints and targets;
- 109 (ii) Designing innovative cropping system prototypes on the basis of current knowledge;
- 110 (iii) Assessments of cropping system prototypes with tools and models adapted for the constraints and targets used,  
111 with improvement of the cropping systems (in terms of rotation or crop management aspects) by an iterative  
112 process, until satisfaction of the constraints or achievement of results considered the best possible; and

113 (iv) Assessment of the most promising cropping system candidates in a long-term field trial. This practical  
114 assessment is currently being carried out in a long-term field experiment, initiated in 2008.  
115

## 116 **1.2. Constraints and targets for innovative cropping systems**

117 Four different cropping systems with quantified constraints (*i.e.*, conditions that had to be fulfilled), environmental and  
118 yield targets, were designed for the agricultural conditions and principal crops of northern France. These constraints and  
119 targets were prioritized as follows: an environmental constraint had to be satisfied first; a set of environmental targets  
120 then had to be attained, and finally, yield had to be maximized. The quantitative levels of the constraints did not  
121 correspond to any regulations and reflected a major break to be reached in current cropping systems. Inclusion of the  
122 use of organic fertilizers (manure, compost, etc.), which are currently not readily available to many farmers in large  
123 areas of Western Europe, was not permitted in the design of the cropping systems.  
124

### 125 **1.2.1. The productive high environmental performance cropping system (PHEP).**

126 No major environmental constraint was placed on this cropping system, which was designed to reach environmental  
127 targets. Eleven environmental indicators, according to the INDIGO® tool<sup>33</sup>, were used to assess the effects of the  
128 cropping system on groundwater pollution (nitrate and pesticides), crop diversity, energy use and soil quality (organic  
129 matter content and phosphorus concentration). To reach environmental goals, all these environmental indicators,  
130 calculated over an entire crop rotation sequence, had to have values higher than 7 (graduated scale from 1 to 10)<sup>33</sup>. This  
131 system was used as the reference system for comparisons with the other three systems.  
132

### 133 **1.2.2. No-pesticide cropping system (No-Pest).**

134 This cropping system was subject to a specific pesticide constraint: no pesticide use was tolerated, even using  
135 substances (e.g. acetic acid) at levels usually considered acceptable in organic cropping systems. However, inorganic  
136 chemical fertilizers were allowed (these fertilizers are not permitted in organic farming systems). This system had to  
137 achieve the same environmental targets as the PHEP cropping system.  
138

### 139 **1.2.3. Low energy cropping system (L-EN).**

140 This cropping system was subjected to a specific energy constraint: it had to have fossil fuel consumption levels no  
141 greater than half those of the PHEP cropping system. It had to reach the same environmental targets as the PHEP  
142 cropping system.  
143

### 144 **1.2.4. Low greenhouse gas emission cropping system (L-GHG).**

145 This cropping system was subject to a specific constraint concerning greenhouse gas emissions: its greenhouse gas  
146 emissions had to be no more than half those of the PHEP cropping system by increasing carbon sequestration in the soil  
147 and decreasing N<sub>2</sub>O emissions. It had to meet the same environmental targets as the PHEP cropping system.

148

149 For each cropping system, once the constraint had been satisfied and environmental targets had been reached, the  
150 combination of agricultural practices giving the highest yields was retained.

151

### 152 **1.3. Design of the four innovative cropping systems**

153 The innovative cropping systems were designed from published knowledge, quantitative data from field experiments  
154 and individual or group expertise provided by scientists, extension service staff and farmers. For each cropping system,  
155 one prototype, consisting of the species in the rotation and the combination of agricultural practices used, was designed.

156 If the constraints were not satisfied, the candidate was modified iteratively (changes to the crops in the rotation or  
157 agronomic practices) until they were. At the beginning of the process, a modification of a crop led to a multitude of  
158 changes; at the end, changes were only one at a time. When the constraints were satisfied, environmental targets were  
159 optimized by an iterative procedure until improvement was observed. Maximum achievable yields were then  
160 determined by experts knowledge or from trial results, for the various cropping systems. The candidate cropping  
161 systems selected for further assessment in a field experiment were, those with the best performances in terms of  
162 constraints, environmental targets and achievable yields.

163 We carefully selected agronomic strategies from previous publications, to satisfy the given constraints. Examples of  
164 such strategies are presented in Table 1. Current knowledge, based on conventional cropping systems, had to be adapted  
165 for innovative cropping systems, and it was necessary to combine strategies. Agronomic strategies were translated into  
166 decision rules (as described by Debaeke<sup>34</sup>) to meet the requirements of future cropping systems and to cope with the  
167 variability of weather and agronomic conditions.

168

### 169 **1.4. Assessment with tools and a model and fine-tuning of innovative cropping system prototypes**

170 During the design process, the cropping system prototypes were assessed with various tools and a model, to determine  
171 the best ways to satisfy the set of constraints and targets imposed. Direct and indirect non-renewable energy  
172 consumption was assessed with the INDIGO® tool (v. 1.9). Direct energy consumption concerned the fuel, lubricants  
173 and electricity used to power farm machinery and tractors. Indirect energy consumption concerned the energy used in  
174 the manufacture, formulation, packaging and maintenance of inputs, such as machinery, fertilizer or pesticides. The  
175 energy outputs of the cropping systems were calculated as the gross energy content of the harvested produce. Energy

176 consumption was calculated on a per hectare basis, per tonne of crop product and per calorie produced, over a complete  
177 crop sequence.

178 C sequestration in the soil was assessed with (i) the Roth C 26.3 model<sup>35</sup> and (ii) the SIMEOS® tool (v.2010) based on  
179 the AMG model<sup>36</sup>. We used climatic data (*i.e.*, monthly mean air temperature, monthly precipitation, and monthly open  
180 pan evaporation) from a meteorological station located in Grignon (Ile-de-France region, 30 km west of Paris). The soil  
181 characteristics (plow layer, 0-30 cm) used to drive simulations were as follows: clay content 20.6%, bulk density 1.4,  
182 initial C content 8 g/kg dry matter. The expected annual yields were estimated from experimental data obtained under  
183 the same conditions (*i.e.*, Ile de France region) and adjusted by expert knowledge. These values were used to estimate  
184 the expected annual dry matter production of roots and stubble, as described by Van Groenigen et al.<sup>37</sup>. Direct and  
185 indirect GHG emissions were estimated with the GES'TIM database<sup>38</sup>. We focused on two main greenhouse gases:  
186 nitrous oxide (N<sub>2</sub>O) and carbon dioxide (CO<sub>2</sub>). Direct emissions included N<sub>2</sub>O emissions from N fertilizers, calculated  
187 with Intergovernmental Panel on Climate Change coefficients<sup>39</sup>, and the CO<sub>2</sub> produced by the combustion of fossil fuels  
188 by farm machinery; CO<sub>2</sub> respired by soil organisms was not taken into account in calculations. Indirect emissions  
189 corresponded to the use of fossil energy in the manufacture and maintenance of farm inputs. GHG balances (C  
190 sequestration plus GHG emissions) were determined over periods of 25 and 50 years, in accordance with  
191 Intergovernmental Panel on Climate Change proposals<sup>39</sup> and current knowledge of C sequestration kinetics in the soil.  
192 In this investigation, any GHG entering the system is counted negatively whereas GHG leaving the system is counted  
193 positively. Therefore, the overall balance is a positive value if more greenhouse gases are emitted than sequestered in  
194 the system.

195  
196 Environmental indicators, such as in the INDIGO® tool (v. 1.9), were used to assess the environmental effects of  
197 cropping system prototypes. Three indicators of nitrogen effects provided information about ammonia volatilization,  
198 nitrous oxide emissions into the air, and nitrate leaching into the groundwater. Four pesticide indicators were studied:  
199 three providing information about pesticide volatilization, pesticide runoff and pesticide leaching into groundwater and  
200 one taking the global effect of pesticides into account. The last four indicators used provided information about crop  
201 diversity, energy consumption, organic matter in the soil, and phosphorus management. Each indicator takes a value  
202 between 1 (worst) and 10 (best). For rotations of more than five crops, the crop diversity indicator was calculated from  
203 the coefficients of Leteinturier<sup>40</sup>. For example, values 0.5, 4.1 and 7.6 respectively correspond to a wheat monoculture,  
204 a wheat-maize rotation and a wheat-sunflower-spring barley-maize rotation.

205 All these tools and the model were chosen on the basis of their relevance for assessing compliance with constraints and  
206 environmental targets. In a comparison of the performance of nine soil organic C models, using different datasets from  
207 long-term experiments from different parts of the world, Smith<sup>41</sup> found that the RothC model was among those that



208 performed best. Bockstaller<sup>42</sup> analyzed four methods for assessing the sustainability of agricultural systems. They found  
209 that the INDIGO® tool was the most relevant for conditions corresponding to those used here. These tools and the  
210 model have been regularly used in different countries. For example, Roth C has been used by De Li Liu<sup>43</sup> and Cerri<sup>44</sup>,  
211 and INDIGO® has been used by Bockstaller<sup>42</sup>.

212

### 213 **1.5 Current cropping system in the Ile-de-France region**

214 The current system in the Ile-de-France region was defined on the basis of data collected in 2006 (Agreste<sup>45</sup>), the most  
215 recent data available at the initiation of this program. We defined the current cropping system in terms of agronomic  
216 practices and crop descriptions. This system was validated by various experts (farmers, extension service staff) with few  
217 adjustments in terms of types and numbers of crops in the rotation. This system was used as a reference for further  
218 comparisons.

219

## 220 **2. Results**

### 221 **2.1. Design methodology step**

222 The systems were designed over a six-month period, by a panel of about 15 experts. For each innovative cropping  
223 system, the first candidate characteristics were based on the current cropping system in the Ile-de-France region. The  
224 system was then optimized through an iterative process, which produced approximately 70 prototypes, to find the four  
225 most promising candidates. These candidates corresponded to the prototypes satisfying the constraint and  
226 environmental targets imposed and yielding the best results in the assessment. For example, for the PHEP system, the  
227 value of the crop diversity indicator was gradually increased from 5 to 7 during the fine-tuning of the system, with  
228 simultaneous improvement of the values of the other indicators. The first prototype was based on a 3-year rotation  
229 (winter oilseed rape, winter barley and winter wheat), currently used in the Ile-de-France region. In the best prototype, a  
230 winter legume and spring barley with a mustard catch crop were gradually introduced, leading to the following rotation:  
231 winter field beans (*Vicia faba*), winter wheat, winter oilseed rape, winter wheat and spring barley with a mustard catch  
232 crop. In the design process, we began by determining the crop rotation and then defined the crop management practices.

233

### 234 **2.2. Description of the innovative cropping system prototypes**

235 For each cropping system, we present only the most promising prototype. The systems are first described in terms of the  
236 crop rotation, crop management practices and yield targets. We then present the results of the final assessment with  
237 respect to constraints (*i.e.*, pesticide use, energy consumption and greenhouse gas emissions) and, finally, we evaluate  
238 the systems in terms of environmental targets. The crop rotations and targeted yields are presented in Tables 2 and 3.

239

240 **2.2.1. The productive high environmental performance cropping system (PHEP)**

241 The PHEP cropping system was designed with multiple environmental targets in mind and was based on the following  
242 agronomic strategies: (i) to reduce pesticide use, we increased crop diversity (four different crops instead of the three  
243 currently sown), (ii) to reduce the amount of N used and indirect energy consumption, we included at least one legume  
244 in the rotation, (iii) to decrease nitrogen leaching, a catch crop was always sown before the spring crop and N  
245 fertilization was forbidden during autumn and winter, (iv) to reduce direct energy consumption, plowing was allowed  
246 only once in the rotation, before the spring crop, (v) to reduce pesticide use and crop loss due to insects and diseases,  
247 highly resistant varieties were used, together with optimal sowing dates and densities and (vi) to stabilize or/and to  
248 enrich the soil organic matter (SOM) content of the soil, crop residues were not removed. As the system had to satisfy  
249 environmental targets requiring the use of fewer inputs, the target yields set were similar to those currently achieved  
250 with low-input cropping systems in the Ile-de-France region.

251

252 **2.2.2. The no-pesticide cropping system (No-Pest)**

253 Pesticide use was prohibited in the “No-Pest” cropping system. Therefore, this cropping system was designed as  
254 follows: (i) to break the cycles of some common soil-borne pathogens, we used a long rotation including a range of  
255 species (five different crops), with the alternate sowing of host and non-host plants, (ii) to reduce weed emergence from  
256 year to year, we sowed species with different sowing dates in spring and in winter successively, (iii) to decrease pest  
257 and disease pressure and damage, we used highly resistant varieties and species mixtures, and excluded crops highly  
258 susceptible to some enemies but with few non-chemical solutions, such as oilseed rape or potatoes, from the rotation  
259 (iv) to increase the competitiveness of the crop with respect to weeds, we sowed species with rapid shoot growth, such  
260 as hemp and triticale, (v) to maximize weed emergence before sowing, we used the stale seed-bed technique, (vi) to  
261 reduce weed emergence after sowing, plowing was carried out before each spring crop and (vii) we adapted sowing  
262 densities to make it possible to use mechanical weeding techniques and to decrease pathogen propagation. We used the  
263 following approaches to reach environmental targets: (i) to reduce nitrate leaching, catch crops were always sown  
264 before spring crops and the spreading of nitrogen fertilizer was allowed only in the spring, (ii) to decrease direct and  
265 indirect energy consumption, we decreased the number of plowing events and N fertilization was calculated according  
266 to yield objectives, and (iii) to stabilize SOM, crop residues were not removed. Yield targets were lower than those for  
267 the PHEP cropping system, because no pesticides were used. However, they were higher than those achieved in organic  
268 systems because chemical fertilizers were allowed, increasing flexibility in the management of crop nitrogen nutrition.  
269 For the integration of these features, in accordance with current knowledge of pest and disease pressures in the Ile-de-

270 France region, experts suggested yield potentials 30% lower than those for the PHEP system for cereals and 25% lower  
271 for field beans.

272

### 273 **2.2.3. Low energy cropping system (L-EN)**

274 The L-EN cropping system was designed, to have a much lower energy consumption than the PHEP cropping system,  
275 as follows: (i) to reduce indirect fuel consumption due to N fertilization, we included as many legumes as possible in  
276 the rotation (field beans as a main crop, clover as a catch crop, and a white clover-winter wheat mixture), and we used  
277 species or varieties with high N use efficiency (*e.g.*, oats <sup>46</sup>) and forms of mineral N fertilizers requiring less energy for  
278 their manufacture, (ii) to decrease direct fuel consumption, we omitted plowing, which is a very resource-intensive  
279 operation, and used a direct drilling system, and (iii) we decreased the amounts of mineral fertilizer (N, P, K) applied,  
280 implying a decrease in target yields. We also designed the L-EN cropping system along the same lines as the PHEP  
281 system, to achieve environmental targets for crop diversity, length of rotation, date of nitrogen spreading, and catch  
282 crop sowing. Target yields were 20% lower than for the PHEP cropping system, except for field beans.

283

### 284 **2.2.4. Low greenhouse gas emission cropping system (L-GHG)**

285 The L-GHG cropping system was designed to decrease greenhouse gas emissions by increasing C sequestration in the  
286 soil and decreasing N<sub>2</sub>O emissions.

287 C sequestration in the soil was increased by (i) including as many cereals as possible in the rotation, to ensure the  
288 production of large amounts of residues (*i.e.*, maize, winter wheat, winter barley or triticale), (ii) maintaining  
289 continuous soil cover to increase the amounts of organic residues (*i.e.*, cover or catch crops were always sown between  
290 main crops, and volunteers were left to grow after harvest), (iii) targeting high yields for the main and catch crops, to  
291 ensure the production of large amounts of residues, and (iv) excluding moldboard plowing, which increases C  
292 mineralization.

293 N<sub>2</sub>O emissions were reduced by (i) decreasing the amount of N fertilizer required at rotation scale<sup>47</sup>, and consequently  
294 direct emissions of N<sub>2</sub>O, by sowing legumes in the rotation (main and catch crops), (ii) improving and optimizing N  
295 fertilization practices according to climatic conditions, through the use of appropriate decision rules to prevent  
296 applications in conditions favoring N<sub>2</sub>O emissions, and (iii) sowing species with taproots to reduce soil compaction and  
297 N<sub>2</sub>O emissions.

298 The L-GHG cropping system was also designed according to the same principles as the PHEP system, to reach  
299 environmental targets for crop diversity, length of rotation, pesticide use, date of N spreading, and catch crop sowing.

300 Target yields were considered to be a compromise between the production of large amounts of C residues (*i.e.*, high

301 yields) and the decrease in N<sub>2</sub>O emissions (*i.e.*, low N fertilization). Experts thought that potential yields would be  
302 similar to those achieved by the PHEP cropping system.

303

### 304 **2.2.5. The current cropping system in the Ile-de-France region**

305 This system is based on a cereal crop rotation, with five cereal crops over a six-year rotation (Table 2). In order to  
306 secure high yields, the agronomic practices were as follows: regular plowings, four times over a 6-year rotation. The  
307 amounts of N fertilizer exceed crop requirements, to prevent yield shortfalls in the event of unfavorable climatic  
308 conditions or unexpected nitrogen losses. Pesticides and growth regulators were used liberally to prevent diseases,  
309 weeds, pests and lodging (3 to 5 pesticides every year).

310

## 311 **3. Cropping system prototypes assessments with tools and a model**

### 312 **3.1. Constraint assessment**

#### 313 **3.1.1. Pesticide constraint in the No-Pest cropping system**

314 This constraint was achieved by not applying pesticides in the No-Pest cropping system.

315

#### 316 **3.1.2. Energy constraint in the L-EN cropping system**

317 Mean total fossil energy consumption (direct and indirect energy), calculated over a single rotation, was 4517 MJ ha<sup>-1</sup>  
318 year<sup>-1</sup> for the L-EN system and 8826 MJ ha<sup>-1</sup> year<sup>-1</sup> for the PHEP system (Figure 1). Chemicals, including N fertilizers,  
319 the largest component, accounted for 1271 MJ ha<sup>-1</sup> year<sup>-1</sup> (43% of total indirect energy consumption) in the L-EN  
320 system and 4345 MJ ha<sup>-1</sup> year<sup>-1</sup> (95% of total indirect energy consumption) in the PHEP system. The use of machinery  
321 for tillage, fertilization, harvesting, sowing and crop protection was the only component of direct energy consumption  
322 that was nearly halved in the L-EN system (2976 MJ ha<sup>-1</sup> year<sup>-1</sup> and 4228 MJ ha<sup>-1</sup> year<sup>-1</sup> for the L-EN and the PHEP  
323 systems, respectively). The difference between these two systems can be accounted for by the absence of tillage and the  
324 use of less N fertilizer in the L-EN cropping system.

325 When expressed in MJ ha<sup>-1</sup>, the total fossil energy in the L-EN system is 49% lower than that in the PHEP system  
326 (Table 4). However, if expressed in MJ t<sup>-1</sup>, the energy performance of the L-EN system is lower (*i.e.* difference between  
327 the PHEP and the L-EN systems of only 24% in term of total fossil energy per tonne of produce), because the target  
328 yield is about 20% lower. A similar reduction in energy use (about 29%) was observed for the calculation in kJ kcal<sup>-1</sup>.

329

#### 330 **3.1.3. The greenhouse gas constraint in the L-GHG cropping system**

331 C sequestration was assessed for the optimized prototypes of the PHEP and the L-GHG systems, for the mean soil  
332 organic matter content in the Ile de France region (1.6%). Both the Roth C model and the SIMEOS® tool predicted that  
333 C would be sequestered throughout the study period, from the start, in both cropping systems. The highest values were  
334 obtained with the L-GHG system over a 50-year period, for both assessment tools (Table 5). For both systems, total C  
335 sequestration was higher during the first 25-year period than during the second 25-year period. When expressed in t  
336 CO<sub>2</sub>-eq ha<sup>-1</sup>, C sequestration values were systematically higher with the Roth C model than with the SIMEOS® tool.  
337 Nevertheless, after 25 and 50 years, the differences between the L-GHG and the PHEP systems calculated with the Roth  
338 C model and the SIMEOS® tool were similar if the results were expressed in relative values.

339 Direct and indirect greenhouse gas emissions were calculated with the GES'TIM database, over one rotation period, for  
340 the L-GHG and the PHEP cropping systems (Figure 2). Mean total greenhouse gas emissions were 1104 kg CO<sub>2</sub>-eq ha<sup>-1</sup>  
341 year<sup>-1</sup> and 1273 kg CO<sub>2</sub>-eq ha<sup>-1</sup> year<sup>-1</sup> for the L-GHG and the PHEP cropping systems, respectively. Direct and indirect  
342 greenhouse gas emissions accounted for similar proportions of total emissions: 48% and 52% for direct greenhouse gas  
343 emissions for the L-GHG and PHEP systems, respectively (Table 6). Chemical fertilizers caused both direct and  
344 indirect greenhouse gas emissions. They represented 76% and 73% of total greenhouse gas emissions for the L-GHG  
345 and the PHEP systems, respectively. Soil cultivation, accounting for 19% and 23% of total greenhouse gas emissions  
346 for the L-GHG and the PHEP systems, respectively, was the second most important component of these emissions.  
347 When results were expressed per ha, total greenhouse gas emissions were 13% lower in the L-GHG system than in the  
348 PHEP system. When expressed per tonne of produce, the larger decrease (22%) may be accounted for by the higher  
349 yields, calculated at rotation scale, of the L-GHG system than of the PHEP system.

350 In terms of the overall balance of GHG emissions (Tables 7-8), GHG values were negative for the L-GHG system,  
351 except for the 50-year period with the SIMEOS® tool. All GHG balance values were lower for the L-GHG system than  
352 for the PHEP system. The difference in GHG balance between the two systems increased over time, and was greater  
353 with the Roth C model, which gave decreases in GHG emission of 51% for the 25-year period and 76% for the 50-year  
354 period.

355

### 356 **3.2. Assessment of environmental targets**

357 The results of assessments of environmental targets with the INDIGO® tool are shown in Figure 3. For all optimized  
358 cropping systems, all 11 indicators had values of at least 7 (*i.e.*, environmental criteria were satisfied), except for the  
359 organic matter indicator for the No-Pest cropping system (OMI = 5.7). The large number of species (more than three in  
360 each rotation) and the small quantities of pesticides sprayed on crops (0 to 2 pesticides used per crop), the systematic  
361 restitution of residues and the optimization of tillage and fertilization management (*i.e.*, the small number of ploughing  
362 operations, optimizing P and N fertilization in terms of both the amounts applied and the timing of applications),

363 resulted in high values for the indicators for crop diversity, pesticide use, soil organic matter, phosphorus, nitrogen and  
364 fossil fuel, respectively.  
365 The low value of the organic matter indicator for the No-Pest cropping system (5.7) could be accounted for by both the  
366 large number of plowing operations (alternate years), encouraging mineralization, and the lower yields, resulting in  
367 smaller amounts of C residues.

368

## 369 **4. Discussion**

### 370 **4.1. Design and assessment of innovative cropping systems**

371 The main challenge of this study was to design innovative cropping systems. Our approach is original in the multiplicity  
372 of purposes assigned to these systems (*i.e.*, association of one major constraint with environmental and yield targets). In  
373 most previous studies, these issues have been analyzed separately. For example, Zentner<sup>48</sup> and Gefland<sup>49</sup> studied energy  
374 efficiency, whereas Nemecek<sup>28</sup> used life cycle assessment methods to evaluate environmental criteria, Nowacki<sup>50</sup>  
375 studied profitability, and Chikowo<sup>51</sup> studied new cropping systems with a lower reliance on pesticides. However,  
376 several other studies are currently investigating system sustainability including assessments of several different  
377 criteria<sup>52</sup>, or numerous environmental parameters<sup>22</sup>. In our project, we combined one major constraint with  
378 environmental and yield targets, reflecting the multifunctionality of agriculture. Furthermore, there was a clear,  
379 particular hierarchy throughout the design process. In most published experiments, environmental consequences are  
380 assessed only during the assessment of technical innovations in the trials, or environmental goals exist but are not  
381 quantified at the start of the study. In our work, satisfying the major constraints and the precise environmental targets  
382 were major aims, which became the conditions determining yield, with target yields set as high as possible under the  
383 conditions concerned. In addition, the clear definition of the constraint (*i.e.*, reducing energy consumption or GHG  
384 emissions by 50%) and the environmental targets (*i.e.*, having a value of at least 7 for all INDIGO environmental  
385 indicators) was also original. The quantitative levels of the constraints did not correspond to any regulations (*i.e.*, these  
386 constraints reflected a major break with the regulations). However these innovative cropping systems were considered  
387 as research tools which enabled to identify the most relevant agronomic practices combinations which could be used in  
388 more restrictive legislative contexts. The level quantifications of the constraints and environmental targets were very  
389 useful during the design process which required calculations.

390 Before the assessment of the prototypes in a long-term field trial, candidate systems were assessed and improved in an  
391 iterative process until the constraints were satisfied and environmental performance with respect to targets was  
392 optimized. This theoretical process of improvement has rarely been reported in previous studies. Cropping systems are  
393 usually assessed or compared in systems defined on the basis of the main standardized characteristics, essentially

394 relating to one major aspect: *e.g.*, organic *versus* conventional systems<sup>53,54,24</sup>, no-tillage *versus* conventional tillage<sup>54,55</sup>,  
395 or integrated *versus* conventional systems<sup>50,56</sup>. Quantitative data for environmental criteria<sup>26</sup>, yield<sup>57</sup> or economic  
396 performance<sup>58,59</sup> were therefore recorded in experiments. The results of these comparisons can be used to compare the  
397 impact of different systems, but not to identify all solutions for their improvement. Even though inductive reasoning can  
398 bring about some conclusions in regards to general principles, another round of conception and evaluation is required,  
399 to strengthen cropping systems. Our prototyping approach is totally different. Innovative systems were assessed by  
400 modeling until they satisfied specific constraints and were optimized in terms of specific environmental targets. A field  
401 trial was then set up to determine whether each of the selected prototypes could satisfy its multiple constraints and  
402 targets. In this case, the various environmental targets were included in the agricultural strategies from the start of the  
403 design process, facilitating identification of the weaknesses of the system and making it possible to propose solutions  
404 for improvement before undertaking field trials. After the assessment of these innovative cropping systems in a field  
405 trial, their costs and economic performances will be calculated in different economic scenarios, to determine the  
406 likelihood of their being adopted by farmers.

407 Our approach required a large panel of experts (scientists, farmers, and extension service staff) to design and to support  
408 prototypes throughout the design process (*i.e.*, from the first to the last candidates). This was necessary because (i) the  
409 best crop management system may not correspond simply to the sum of individual agricultural practices, but may  
410 instead involve a set of agronomic strategies and their interactions, and (ii) a breadth of agro-ecological knowledge is  
411 required to identify sets of agricultural rules likely to be responsive to such strict constraints and environmental targets.  
412 Moreover, this approach provided a more realistic view of cropping systems, making the adoption of the proposed  
413 innovations more likely<sup>60</sup>. However, the field trial assessment step is still absolutely necessary because some innovative  
414 agronomic practices, not currently used in cropping systems, have never been evaluated by experts.

415 During the design process, about 15 experts attended individual sessions or group meetings, to provide knowledge  
416 unavailable from published work. The definition of crop rotations and agronomic practices took about six months, and a  
417 further 18 months were required for the writing of the decision rules. Published studies involving design processes have  
418 differed considerably in the number of experts involved and the time spent by individual experts, depending on the  
419 availability of the experts and the difficulties encountered in achieving the goals assigned to systems<sup>52,61</sup>.

420

## 421 **4.2. Achievement of multiple constraint and targets**

422 For all innovative systems, the constraints were satisfied with no consequences for other environmental components,  
423 except for the organic matter indicator of the No-Pest system. In this case, regular tillage combined with the restitution  
424 of only small amounts of organic matter had an adverse effect on soil environmental characteristics (indicator value for  
425 soil organic matter of 5.7, according to the INDIGO® tool). Within this system, it did not appear to be possible to

426 satisfy both the constraint and this environmental target with the available non-chemical techniques for pest control.  
427 Moreover, this was only possible with the available techniques by reducing yield targets with respect to those of current  
428 regional systems (Agreste<sup>45</sup>). Nevertheless, progress in integrated pest management is being made, and new techniques  
429 may make it possible to improve environmental and yield performances. In the design of the L-EN system, we managed  
430 various agricultural processes, decreasing both direct energy consumption (due to tillage, for example) and indirect  
431 energy consumption (due to the use of mineral fertilizers). We halved fossil energy consumption by greatly decreasing  
432 N fertilizer inputs, which was associated with a 20% yield loss. However, the energy performance of the L-EN system  
433 was expressed relative to that of the PHEP system, which also had a relatively low level of fossil fuel consumption with  
434 respect to current practices in Ile-de-France. The total energy consumption of the L-EN system was about 35% that of  
435 the current system in Ile-de-France (Agreste<sup>45</sup>), when energy was expressed in MJ ha<sup>-1</sup> (Table 4). For both the L-GHG  
436 and the PHEP systems, decreases in pesticide were taken into account by considering the maximum achievable yields to  
437 be similar to those of current low-input systems in Ile-de-France (Agreste<sup>45</sup>), and much lower than those of  
438 conventional systems in the region (Agreste<sup>45</sup>). Achievable yields for the L-GHG and the PHEP systems were  
439 considered to be 13% and 21% lower, respectively, than those of the current system in Ile-de-France. However,  
440 considering all the innovative systems together, it would appear to be possible to satisfy such ambitious constraints and  
441 environmental targets at the expense of only relatively small yield losses.

442 Available knowledge and current techniques suggested that it would not be possible to overcome all constraints in a  
443 single cropping system, because the agronomic practices used in one innovative system were incompatible with the  
444 constraints imposed on others. Plowing, one of the most effective practices against weeds used in the No-Pest system, is  
445 incompatible with large decrease in fossil energy consumption and the increase in C sequestration achieved with the L-  
446 EN and L-GHG systems, respectively. The large decrease in N fertilizer levels of the L-EN system is not compatible  
447 with the achievable yields defined for the PHEP and the L-GHG systems. Winter wheat sowing was delayed to avoid  
448 pest pressure in the No-Pest system, whereas it was brought forward in the direct drilling conditions of the L-EN and  
449 the L-GHG systems. This pattern was already evident during the design of the L-GHG system (*i.e.*, GHG emission  
450 processes were managed in hierarchical order). Consequently, the development of a system without pesticides, with  
451 ambitious constraints in terms of GHG emissions and fossil fuel use, and with other environmental and yield targets,  
452 will require further progress in agronomic knowledge. For example, a better understanding of the interactions between  
453 cash and cover crops in terms of cooperative and competitive effects might allow the introduction of a living cover crop  
454 during cash crop growth in the L-GHG system<sup>62</sup>. The field trial assessment again proves essential to gain a better  
455 understanding of these interactions.

456

### 457 **4.3. Improving the design process**



458 In the design of the L-GHG system, we had to rank the secondary objectives (C sequestration had to be enhanced first,  
459 and then N<sub>2</sub>O emissions had to be reduced) to satisfy the GHG constraint. In this case, several practices had effects on  
460 both processes involved: no-tillage increased C sequestration in the soil but increased N<sub>2</sub>O emissions; ample N fertilizer  
461 applications were required to obtain high yields and, thus, abundant C residues, but this also generated more N<sub>2</sub>O  
462 emissions. We decided to promote C sequestration, because N<sub>2</sub>O emission assessments were highly uncertain due to the  
463 lack of published data about N<sub>2</sub>O emissions, for field bean residues for example (IPPC<sup>39</sup>), and the variability of results  
464 due to differences in soil and climatic conditions<sup>63</sup>. Nevertheless, knowledge about the effects of cropping systems on  
465 N<sub>2</sub>O emissions is increasing, and it should be possible to improve the adjustment of cropping systems in the future.

466 The cropping systems assessment required tools and models adapted to the set of objectives and convenient to use  
467 during the iterative optimization process. Some approximations were used, due to the lack of data. In the L-GHG  
468 system, the coefficient defined by the Intergovernmental Panel on Climate Change (*Tier 1*) was used to calculate N<sub>2</sub>O  
469 release from the amount of N applied, rather than using different values for different soil and climatic conditions<sup>63</sup>.

470 There were also uncertainties in the assessment of C sequestration. The two simulations provided similar ranks for the  
471 PHEP and the L-GHG systems, but the SIMEOS® tool and the Roth C model gave different quantitative results. One  
472 difference between these tools relates to tillage. In the SIMEOS® tool, mineralization mechanisms in the tilled layer are  
473 modified by tillage<sup>36</sup>. In the Roth C model, tillage is taken into account only indirectly, through the C inputs to the soil  
474 from crop biomass, which are affected by tillage practices<sup>43</sup>. Further uncertainties arise from the lack of root biomass  
475 data. Furthermore, these tools partially take into account changes in soil organic matter content, which greatly influence  
476 N transformation mechanisms in the soil. Anyway, the difference in the values generated by these models suggested  
477 that neither may accurately predict actual performances of these contrasting cropping systems. If cropping systems  
478 assessment is to be considered relevant, it must make use of tools that are regularly updated, including: (i) “new”  
479 machinery, such as harrows, hoes or direct-sowing drills with adjustable parameters in terms of energy consumption  
480 and GHG emissions, and (ii) new more forms of pesticides or fertilizers. It may also be useful to improve systems  
481 assessments with crop models, but many parameters are unavailable for marginal crops (such as hemp or flax), for crop  
482 mixtures (*e.g.*, cereal and legume combinations), or species mixtures used as catch crops (*e.g.*, the association of spring  
483 oats, mustard and clover).

484 Several economic and social aspects were not taken into account during this design process, which focused on the  
485 cropping system rather than the farm scale. Nevertheless, we excluded some crops that are not grown by farmers in Ile-  
486 de-France due to the lack of a market (*i.e.*, with low economic performances), but we included others for which the  
487 market is poorly developed in this area (*e.g.*, lucerne and hemp). We did not take the organization of farm work into  
488 account either, although the identification and quantification of pest pressures in the field, to decrease pesticide use, is  
489 known to be time-consuming. From a technical point of view, we assumed that all farms owned the specific machinery

490 used in the innovative systems, such as direct-sowing drills or mechanical weeding tools. For all these reasons, further  
491 investigations of the most relevant innovative cropping systems identified in a field trial assessment are needed, to  
492 forecast their possible application in an area.

#### 493 494 **4.4. Toward field assessment of the innovative cropping system prototypes**

495 The innovative systems prototypes designed here were based on different hypotheses about soil and climate effects,  
496 because the true impact on crops and soil are unknown. For example we assumed that agricultural practices and climatic  
497 conditions would not affect crop emergence. In our conditions, crops are not affected by direct drilling because this  
498 technique is widely used in France (<http://agriculture-de-conservation.com/>). However, conservation tillage and mulch  
499 tillage practices remain largely empiric<sup>64</sup>, and additional knowledge is required for the optimization of these techniques,  
500 to make it possible to achieve good results in terms of sowing management and target yields. We also assumed that  
501 rainfall would provide enough water throughout the year in northern France to allow catch crop and main crop  
502 emergence, but this may no longer be the case if the climate changes radically. We assumed that new equilibria would  
503 appear with certain practices, allowing inputs to be reduced. In the L-GHG system, the practice of leaving crop residues  
504 on the soil should decrease weed emergence, making it possible to decrease the amount of herbicide used, and the  
505 absence of plowing should lead to the maintenance or increase in size of ground beetle populations, making  
506 molluscicide use unnecessary. The achievable yield was determined on the basis of several assumptions. In the L-GHG  
507 system, we promoted C sequestration, which is highly dependent on cereal yields and difficult to estimate in the case of  
508 no-tillage systems, because the presence of excessive amounts of crop residues may decrease emergence. The chemical  
509 properties of the soil associated with different tillage practices are also poorly characterized<sup>64</sup>. In this system, the effects  
510 on water availability of sowing cover crops every year also should be analyzed. In the L-EN system, the mixture of  
511 winter wheat and white clover might also decrease cereal yields. In the No-Pest system, yields were defined by  
512 approximation to organic systems, in which no mineral fertilizers are permitted. In this system, late winter cereal  
513 sowing might lead to higher levels of damping-off and plant death during winter, decreasing yields to a greater extent  
514 than anticipated. These are just a few of the uncertainties remaining about the real performance of the systems we have  
515 devised. For these reasons, the field assessment of these system prototypes, which is currently being carried out (the  
516 experiment started in 2008), is absolutely necessary.

517 Finally, it should be stressed that approaches of this type could be used in many agronomic situations, with a diversity  
518 of challenges, provided that sufficient knowledge is available for the development of innovative strategies. In our  
519 opinion, this approach meets the need expressed by Foley<sup>14</sup> to “search for practical solutions” for a cultivated planet.

520

521 **Acknowledgments:**

522 We thank the experts who took part in the design process, Marie-Hélène Jeuffroy, Jane Jonhson, Laurence Guichard,  
523 Marie Gosme and David Makowski for their useful comments on a previous version of the paper, Suzette Tanis-Plant  
524 for linguistic help, and Alain Bone for assistance with bibliographic research.

525

526 **References**

527 1. IFEN, 2007. Les pesticides dans les eaux – données 2005. Les dossiers – n°9, ISSN: 1776-8411, 35 p.  
528 <[http://www.ifen.fr/publications/le-catalogue-despublications/les-dossiers/2007/les-pesticides-dans-les-eaux-donnees-](http://www.ifen.fr/publications/le-catalogue-despublications/les-dossiers/2007/les-pesticides-dans-les-eaux-donnees-2005.html)  
529 [2005.html](http://www.ifen.fr/publications/le-catalogue-despublications/les-dossiers/2007/les-pesticides-dans-les-eaux-donnees-2005.html)>.

530 2. Schrack D., Coquil X., Ortar A., and Benoît M., 2009. Rémanence des pesticides dans les eaux issues de parcelles  
531 agricoles récemment converties à l'Agriculture Biologique. *Innovations Agronomiques* 4, 259-268

532 3. Höök M., Sivertsson A., and Aleklett K., 2010. Validity of the fossil fuel production outlooks in the IPCC Emission  
533 Scenarios. *Natural Resources Research: Volume 19, Issue 2*, 63-81

534 4. Jakobsson K., Söderbergh B., Höök M. and Aleklett K., 2009. How reasonable are oil production scenarios from  
535 public agencies? *Energy Policy*, Vol. 37, 4809-4818.

536 5. Shafiee S and Topal E., 2009. When will fossil fuel reserves be diminished? *Energy Policy*, vol. 37, issue 1, pages  
537 181-189

538 6. Mohr S.H., Evans G.M., 2010. Long term prediction of unconventional oil production. *Energy Policy*, Volume 38,  
539 Issue 1, 265-276

540 7. Smith P., Martino D., Cai Z., Gwary D., Janzen H., Kumar P., McCarl B., Ogle S., O'Mara F., Rice, C., Scholes B.,  
541 and Sirotenko O., 2007. Agriculture. In *Climate Change 2007: Mitigation. Contribution of Working Group III to the*  
542 *Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Metz, B. and Davidson, O. R. and  
543 Bosch, P. R. and Dave, R. and Meyer, L. A.[eds.]. Cambridge University Press, Cambridge, United Kingdom and New  
544 York, NY, USA.

545 8. Galloway J. N., Townsend A. R., Erisman J. W., Bekunda M., Cai Z. C., Freney J. R., Martinelli L. A., Seitzinger  
546 S.P., Sutton M.A., 2008. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science*,  
547 16;320(5878):889-92. doi: 10.1126/science.1136674.

548 9. Arrouays D., Balesdent J., Germon J.C., Jayet P.A., Soussana J.F., and Stengel P., 2002. Expertise scientifique  
549 collective "Contribution à la lutte contre l'effet de serre" Stocker du carbone dans les sols agricoles de France ?  
550 Collective Scientific Expert Report. INRA edition, pp 332

- 551 10. Spiertz J.H.J., 2010. Nitrogen, sustainable agriculture and food security. A review *Agronomic Sustainable*  
552 *Development*. 30 (2010) 43–55
- 553 11. Matson P.A., Parton W.J, Power A.G., and Swift M.J., 1997. Agriculture intensification and ecosystems properties,  
554 *Science* 277, 504-509
- 555 12. UNEP 2007 [http://www.unep.org/PDF/AnnualReport/2007/AnnualReport2007\\_en\\_web.pdf](http://www.unep.org/PDF/AnnualReport/2007/AnnualReport2007_en_web.pdf)
- 556 13. Bouman B.A.M., Humphreys E., Tuong T.P., and Barker R., 2007. Rice and Water, *Advances in Agronomy*, 92,  
557 187-237
- 558 14. Foley J.A., Ramankutty N., Brauman K.A., Cassidy E.S., Gerber J.S., Johnston M., Mueller N.D., O’Connell C.,  
559 Ray D.K., West P.C., Balzer C., Bennett L.M., Carpenter S.R., Hill J., Monfreda C., Polasky S., Rockström J., Sheehan  
560 J., and Siebert S., 2011. Solutions for a cultivated planet. *Nature*, 10452
- 561 15. Aubertot J.N., Barbier J.M., Carpentier A., Gril J.J., Guichard L., Lucas P., Savary S., Savini I., and Voltz M., 2005.  
562 Pesticides, agriculture and the environment: reducing the use of pesticides and limiting their environmental impact.  
563 Collective Scientific Expert Report. INRA – CEMAGREF. INRA edition, pp 63
- 564 16. Köpke U. and Nemecek T., 2010. Ecological services of faba bean. *Field Crops Research* 115 (2010) 217–233
- 565 17. Beheydt D., Boeckx P., Ahmed H. P., and Van Cleemput O., 2008. N<sub>2</sub>O emissions from conventional and  
566 minimum-tilled soils. *Biology and Fertility of Soils* 44, 863–873
- 567 18. Van Kessel C., Ventera R., Six J, Arlene M., Borde A., Linqvist B., and Van Groenigen K., 2012. Climate, duration,  
568 and N placement determine N<sub>2</sub>O emissions in reduced tillage systems: a meta-analysis. *Global Change Biology*, doi:  
569 10.1111/j.1365-2486.2012.02779.x
- 570 19. Fu WeiJun, Tunney H., and Zhang Chao Sheng, 2010. Spatial variation of soil test phosphorus in a long-term  
571 grazed, experimental grassland field. *Journal of Plant Nutrition and Soil Science*, 173, 3, 323-331.
- 572 20. Heinze S., Raupp J., and Joergensen R. G., 2010. Effects of fertilizer and spatial heterogeneity in soil pH on  
573 microbial, biomass indices in a long-term field trial of organic agriculture. *Plant and Soil*, 328, 1/2, 203-215
- 574 21. Chirinda N., Olesen J.E., Porter J.R., and Schjønning P., 2010. Soil properties, crop production and greenhouse gas  
575 emissions from organic and inorganic fertilizer-based arable cropping systems. *Agriculture, Ecosystems &*  
576 *Environment*, Volume 139, No. 4, 584-594
- 577 22. Tuomisto H.L., Hodge I.D., Riordan P., and Macdonald D.W., 2012. Comparing energy balances, greenhouse gas  
578 balances and biodiversity impacts of contrasting farming systems with alternative land uses. *Agricultural Systems* 108 :  
579 42–49
- 580 23. Shadbolt N., Kelly T., Horne D., Harrington K., Kemp P., Palmer A., and Thatcher A., 2009. Comparisons between  
581 organic & conventional pastoral dairy farming, systems: cost of production and profitability. *Journal of Farm*  
582 *Management*, 13, 10, 671-685

- 583 24. De Ponti T., Rijk B., and Van Ittersum M., 2012. The crop yield gap between organic and conventional agriculture.  
584 *Agricultural Systems* 108: 1–9
- 585 25. Simonsen J., Posner J., Rosemeyer M., and Baldock J., 2010. Endogeic and anecic earthworm abundance in six  
586 Midwestern cropping systems. *Applied Soil Ecology* 44. 147–155
- 587 26. Riley H., Pommeresche R., Eltun R., Hansen S., and Korsæth A., 2008. Soil structure, organic matter and  
588 earthworm activity in a comparison, of cropping systems with contrasting tillage, rotations, fertilizer, levels and manure  
589 use., *Agriculture, Ecosystems Environment*, 124, 3/4, 275-284
- 590 27. Snapp S.S., Gentry L.W., and Harwood R., 2010. Management intensity – not biodiversity – the driver of ecosystem  
591 services in a long-term row crop experiment. *Agriculture, Ecosystems & Environment* 138 : 242–248
- 592 28. Nemecek T., Dubois D., Huguenin-Elie O., and Gaillard G., 2011. Life cycle assessment of Swiss farming systems:  
593 I. Integrated and organic farming. *Agricultural Systems* 104 (2011) 217–232
- 594 29. Vereijken P., 1997. A methodological way of prototyping integrated and ecological arable farming systems  
595 (I/EAFS) in interaction with pilot farms. *European Journal of Agronomy*, 7: 235-250
- 596 30. Dogliotti S., Van Ittersum M.K., and Rossing W.A.H., 2005. A method for exploring sustainable development  
597 options at farm scale: a case study for vegetable farms in South Uruguay. *Agricultural Systems* 86: 29–51
- 598 31. Loyce C. and Wery J., 2006. Les outils des agronomes pour l'évaluation et la conception de systèmes de culture.  
599 *L'agronomie aujourd'hui*, QUAE éditions, 77-95.
- 600 32. Lançon J., Wery J., Rapidel B., Angokaye M., Gérardaux E., Gaborelc C., Ballo D., and Fadegnon B., 2007. An  
601 improved methodology for integrated crop management systems. *Agronomy of Sustainable Development*, 27: 101–110
- 602 33. Bockstaller C., Guichard L., Makowski D., Aveline A., Girardin P., and Plantureux S., 2008. Agri-environmental  
603 indicators to assess cropping and farming systems. A review. *Agronomy of Sustainable Development*, 28: 139-149
- 604 34. Debaeke P., Munier-Jolain N., Bertrand M., Guichard L., Nolot J.M., Faloya V., and Saulas P., 2009. Iterative  
605 design and evaluation of rule-based cropping systems: methodology and case studies. *Agronomy of Sustainable*  
606 *Development*, 29: 73–86
- 607 35. Coleman K. and Jenkinson D.S., 1999. A model for the turnover of carbon in soil; Model description and windows  
608 users guide, IACR – Rothamsted Harpenden Herts AL5 2JQ, p47
- 609 36. Andriulo A., Mary B., and Guerif J. 1999. Modelling soil carbon dynamic with various cropping sequences on the  
610 rolling pampas. *Agronomie*, 19: 365-379
- 611 37. Van Groenigen K.J. , Hastings A., Forristal D., Jones M., and Smith P., 2011. Soil C storage as affected by tillage  
612 and straw management: An assessment using field measurements and model predictions. *Agriculture, Ecosystems &*  
613 *Environment*, 140: 218-225

- 614 38. GES'TIM (2010): Ministère de l'agriculture, de l'alimentation, de la pêche, de la ruralité et de l'aménagement du  
615 territoire. Réf. 0933103, 156p
- 616 39. Intergovernmental Panel on Climate Change. Climate Change 2007. The physical science basis. UK: Cambridge  
617 University Press, 1009p
- 618 40. Leteinturier B., Herman J.L., de Longueville F., Quintin L., and Oger R., 2006. Adaptation of a crop sequence  
619 indicator based on a land parcel management system. *Agriculture, Ecosystems and Environment* 112 (2006) 324–334
- 620 41. Smith P., Smith J., Powlson D., McGill W., Arah J., Chertov O., Coleman K., Frnko U., Froking S., Jenkinson D.,  
621 Jensen L., Kelly R., Klein-Gunnewiek H., Komarov A., Li C., Molina J., Mueller T., Parton W., Thornley J., and  
622 Whitmore A., 1997. A comparison of the performance of nine soil organic matter models using datasets from seven  
623 long-term experiments. *Geoderma* 81, 153–225.
- 624 42. Bockstaller C. Guichard L., Keichinger O., Girardin P., Galan MB., Gaillard G., 2009. Comparison of methods to  
625 assess the sustainability of agricultural systems. A review. *Agronomy of Sustainable Development*. 29: 223–235
- 626 43. De Li Liu, Yin Chan K., Conyers M.K., Li G., and Poile G.J., 2011. Simulation of soil organic carbon dynamics  
627 under different pasture managements using the RothC carbon model. *Geoderma*, 165: 69–77
- 628 44. Cerri C., Easter M., Paustian K., Killian K., Coleman K., Bernoux M., Falloon P., Powlson D., Batjes N., Milne E.,  
629 and Cerri C.C., 2007. Simulating SOC changes in 11 land use change chronosequences from the Brazilian Amazon with  
630 RothC and Century models. *Agriculture, Ecosystems & Environment*, 122 (1), 46–57
- 631 45. Agreste, an agricultural statistical service of the French government <http://agreste.agriculture.gouv.fr/>
- 632 46. Riley H., Hoel B.O., Kristoffersen A.Ø., 2012. Economic and environmental optimization of nitrogen fertilizer  
633 recommendations for cereals in Norway. *Acta Agriculturae Scandinavica, Section B, Soil & Plant Science*, 62, 5, 387-  
634 400
- 635 47. Jeuffroy M.H., Baranger E., Carrouée B., de Chezelles E., Gosme M., Henault C., Schneider A., Cellier P., 2013.  
636 Nitrous oxide emissions from crop rotations including wheat, oilseed rape and dry peas. *Biogeosciences*, 10, 1787–1797
- 637 48. Zentner R.P., Lafond G.P., Derksen D.A., Nagyd C.N., Wall D.D., and May W.E., 2004. Effects of tillage method  
638 and crop rotation on non-renewable energy use efficiency for a thin Black Chernozem in the Canadian Prairies. *Soil &*  
639 *Tillage Research* 77: 125–136
- 640 49. Gelfand I., Snapp S., and Robertson G., 2010. Energy efficiency of conventional, organic, and alternative cropping  
641 systems for food and fuel at a site in the US Midwest. *Environment Sciences Technology*, 44, 4006–4011
- 642 50. Nowacki W., 2008. Comparison of profitability of potato cultivation in organic and, integrated farming systems.  
643 *Progress in Plant Protection*, 48, 4, 1526-1534

- 644 51. Chikowo R., Faloya V., Petit S., and Munier-Jolain N., 2009. Integrated weed management systems allow reduced  
645 reliance on herbicides and long-term weed control. *Agriculture, Ecosystems & Environment*, 132: 237–242
- 646 52. Blazy J.M., Ozier-Lafontaine A., Doré T., Thomas A., and Wery J., 2009. A methodological framework that  
647 accounts for farm diversity in the prototyping of crop management systems. Application to banana-based systems in  
648 Guadeloupe. *Agricultural Systems* 101 . 30–41
- 649 53. Coulter J.A., Sheaffer C.C., Wyse D.L., Haar M.J., Porter P.M., Quiring S.R., and Klossner L.D., 2011. Agronomic  
650 performance of cropping systems with contrasting crop rotations and external inputs. *Agronomy Journal*, 103: 1, 182
- 651 54. Moreno M., Lacasta C., Meco R., and Moreno C., 2011. Rainfed crop energy balance of different farming systems  
652 and crop rotations in a semi-arid environment: Results of a long-term trial. *Soil & Tillage Research*, 114, 1: 18-27
- 653 55. Feiza V., Feiziene D., Auskalnis A., and Kadziene G., 2010. Sustainable tillage: results from long-term field  
654 experiments on Cambisol., *Zemdirbyste (Agriculture)*, 97, 2: 3-14
- 655 56. Baturo A., 2007. Effect of organic system on spring barley stem base health in, comparison with integrated and  
656 conventional farming. *Journal of Plant Protection Research*, 47, 2: 167-178
- 657 57. Klimekova M. and Lehocka Z., 2009. Comparison of yields and qualitative characters of spring barley grown, after  
658 three preceding crops in an organic farming system in the years, 2003-2008. *Agronomy Research*, 7, Special Issue, 335-  
659 340
- 660 58. Liebman M., Gibson L. Sundberg D., Heggenstaller A., Westerman P., Chase C., Hartzler R., Menalled H., Davis  
661 M., and Dixon P., 2008. Agronomic and economic performance characteristics of conventional and low-external-input  
662 cropping systems in the Central Corn Belt. *Agronomy Journal*, 100, 3: 600-610
- 663 59. Cavigelli M. A., Hima B.L., Hanson J.C., Teasdale J.R., Conklin A.E., and Lu Y., 2009. Long-term economic  
664 performance of organic and conventional field crops in the mid-Atlantic region. *Renewable Agriculture and Food  
665 Systems*. 24, 2: 102-119
- 666 60. Le Gal P.Y., Dugué P., Faure G., and Novak S., 2011. How does research address the design of innovative  
667 agricultural production systems at the farm level? A review *Agricultural Systems* 104 : 714–728
- 668 61. Stoorvogel J.J., Bouma J., and Orlich R.A., 2004. Participatory research for systems analysis: Prototyping for a  
669 Costa Rican banana plantation. *Agronomy Journal*, 96, 2: 323-336
- 670 62. Shili-Touzi I., de Tourdonnet S., Nciri R., Le Floch D., Saulas P., and Doré T. (submitted). Is it possible to manage  
671 competition between wheat and alfalfa grown as a service plant in an intercropping system in temperate conditions?  
672 *Agriculture, Ecosystems & Environment*.
- 673 63. Rochette P. and Janzen H., 2005. Towards a revised coefficient for estimating N<sub>2</sub>O emissions from legumes.  
674 *Nutrient Cycling in Agroecosystems* 73, 171–179.

675 64. Alletto L., Coquet Y., and Roger-Estrade J., 2010. Two-dimensional spatial variation of soil physical properties in  
676 two tillage systems. *Soil Use and Management*, 26, 432–444

677 65. Munier-Jolain N.M., Faloya V., Davaine J.B., Biju-Duval L., Meunier D., Martin C., and Charles R., 2004. A  
678 cropping system experiment for testing the principles of integrated weed management: first results, in *Annales AFPP*,  
679 XIIe colloque international sur la biologie des mauvaises herbes, Dijon, 147–156.

680 66. Tonitto C., David M. B., Li C. S., and Drinkwater L. E., 2007. Application of the DNDC model to tile-drained  
681 Illinois agroecosystems: model comparison of conventional and diversified rotations. *Nutrient Cycling in*  
682 *Agroecosystems* 78, 65–81.

683 67. Deike S., Pallutt B., Melander B., Strassemeier J., and Christen O., 2008. Long-term productivity and  
684 environmental effects of arable farming as affected by crop rotation, soil tillage intensity and strategy of pesticide use:  
685 A case-study of two long-term field experiments in Germany and Denmark. *European Journal of Agronomy*, 29: 191–  
686 199

687 68. Nemecek T., von Richthofen J.F., Dubois G., Casta P., Charles R., and Pahl H., 2008. Environmental impacts of  
688 introducing grain legumes into European crop rotations, *European Journal of Agronomy* 28: 380–393

689 70. XueLi Fu; Zhang Hui, and Jia JiZeng., 2009. Yield performance and resources use efficiency of winter wheat and  
690 summer maize in double late-cropping system. *Acta Agronomica Sinica*: 35, 9: 1708-1714

691 71. Freibauer A., Rounsevell M. D. A., Smith P., and Verhagen J., 2004. Carbon sequestration in the agricultural soils  
692 of Europe. *Geoderma* 122, 1–23.

693 72. Mosier A. R., Halvorson A. D., Peterson G. A., Robertson G. P., and Sherrod L., 2005. Measurement of net global  
694 warming potential in three agroecosystems. *Nutrient Cycling in Agroecosystems* 72, 67–76.

695 73. Lehuger S., 2009. Modélisation des bilans de gaz à effet de serre des agro-systèmes en Europe. Thèse de doctorat de  
696 l'Institut des Sciences et Industries du Vivant et de l'Environnement AgroParisTech. p184

697 74. Pelster D.E., Larouche F., Rochette P., Chantigny M.H., Allaire S., and Angers D.A., 2011. N fertilization but not  
698 soil tillage affects nitrous oxide emissions from a clay loam soil under a maize–soybean rotation. *Soil and Tillage*  
699 *Research*. 115-116: 16-26

700



701 **Tables**

702 Table 1: Examples of published agronomic strategies and associated practices for achieving specific goals

Specific goal	Agronomic management strategies	Example of agronomic practices
<b>To reduce pesticide use:</b> (Munier-Jolain <sup>65</sup> ; Aubertot <sup>15</sup> ; Chikowo <sup>51</sup> )	To avoid the coincidence of the pest, disease and weed contamination periods and sensitive stages of crop development	Modification of sowing date, earlier for oilseed rape, later for winter wheat
	To reduce the density of pests	Use of mechanical weeding Use of <i>Trichogramma</i> parasitoid wasps against <i>Ostrinia nubilalis</i> on maize
	To reduce the impacts of pests, diseases and weeds on crops	Decrease in sowing density and N fertilization to decrease shoot biomass Choice of varieties with the highest resistance Maximization of competition against weeds, by sowing winter oilseed rape very early (mid-August in the Ile-de-France region)
	To decrease the pool of pathogenic fungi in the soil, diseases and weeds	Lengthening of the crop rotation Sowing of a wide range of crops, to decrease the pool of fungi in the soil Maximization of the use of stale seed beds to increase weed emergence before sowing
	To maintain beneficial insects	Shallow plowing to maintain populations of carabid beetles (slug predators)
<b>To reduce fossil fuel energy use:</b> (Tonitto <sup>66</sup> ; Deick <sup>67</sup> ; Nemecked <sup>68</sup> ; XueLi <sup>70</sup> ; Gelfand <sup>49</sup> )	<b>Direct energy:</b> To decrease the use of the most power- consuming farm machinery <b>Indirect energy:</b> To decrease chemical N fertilization	Reducing or eliminating the use of deep plowing  Introduction of large numbers of legumes and species with the highest N use efficiency into the crop rotation
<b>To enhance C sequestration in the soil:</b> (Arrouays <sup>9</sup> ; Freibauer <sup>71</sup> ; Mosier <sup>72</sup> ; Beheydt <sup>17</sup> ; Smith <sup>7</sup> ; Lehuger <sup>73</sup> )	To decrease organic matter mineralization  To allow high levels of C accumulation, depending on the nature of the crop residues	Avoidance of plowing, replaced by direct drilling (conversion to no-tillage practices)  Sowing of maize and cereals in rotation
<b>To reduce N<sub>2</sub>O emissions:</b> (Rochette <sup>63</sup> ; Pelster <sup>74</sup> )	To decrease anaerobic conditions, to decrease denitrification mechanisms To decrease the amount of available mineral N in the soil	Decreasing compaction and soil moisture content without deep tillage Optimization of N fertilization, according to soil nitrogen content Increasing the N use efficiency of crops
<b>To decrease nitrogen leaching</b>	To decrease soil nitrate content during autumn and winter	Sowing of catch crops Sowing of oilseed rape after legumes Banning of N fertilization during autumn and winter

703

704

705 Table 2: Crop rotations of the innovative cropping systems designed, and a current cropping system, defined by experts  
 706 on the basis of data from the Ile-de-France region in 2006 (Agreste <sup>45</sup>)

Cropping system	Crop rotation
Productive high environmental performance (PHEP)	Winter field bean - Winter wheat - Winter oilseed rape - Winter wheat – (with mustard as a catch crop) - Spring barley
Without pesticide (No-Pest)	Triticale – (species mixture as a catch crop) Maize – Winter wheat – (species mixture as a catch crop) Spring field bean - Winter wheat - (species mixture as a catch crop) - Hemp
Low energy use: (L-EN)	Winter field bean - Winter wheat - Winter oil flax - Winter wheat-white clover mixture – (white clover as a catch crop) Spring oat
Low greenhouse gas emissions (L-GHG).	Triticale – (frost-sensitive species mixture as a catch crop) - Spring field bean - Winter oilseed rape (volunteers) - Winter wheat – (legumes as a cover crop) - Winter barley – (legume-oat mixture as catch crop) Maize
Ile-de-France (IdF)	Winter oilseed rape – Winter wheat – Winter barley – Maize – Winter wheat - Winter wheat

707

708

709 Table 3: Targeted yields (t ha<sup>-1</sup>) for different species sown in the innovative cropping systems and comparison with  
 710 current yields. Cropping systems: PHEP (productive high environmental performance), No-Pest (no pesticide use), L-  
 711 EN (low energy use), L-GHG (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).

Cropping systems / Crops (t ha <sup>-1</sup> )	PHEP	No-Pest	L-EN	L-GHG	IdF Mean from 1998 to 2007 (Agreste)
Spring barley	6.2	-	-	-	
Winter barley	-	4.6	-	6.6	6.6
Spring field bean	-	4.7	-	4.7	
Winter field bean	3.4	-	3.4		
Hemp	-	8	-	-	
Maize	-	7.3	-	9.1	9.1
Winter oil flax	-	-	1.8	-	
Winter oilseed rape	3.1	-	-	3.1	3.4
Triticale	-	4.2	-	6.1	
Winter wheat	7.9	5.5	6.3	7.9	8.0

712

713

714 Table 4: Comparisons of total fossil energy consumption, expressed in MJ ha<sup>-1</sup>, MJ t<sup>-1</sup> and kJ jCal<sup>-1</sup> (INDIGO® tool,  
 715 v.1.9), between cropping systems. Cropping systems: PHEP (productive high environmental performance), L-EN (low  
 716 energy use) and IdF (current system in Ile-de-France region).

	Total fossil energy consumption (MJ ha <sup>-1</sup> )	Total fossil energy consumption (MJ t <sup>-1</sup> )	Total fossil energy consumption (kJ kCal <sup>-1</sup> )
<u>L-EN - PHEP</u> PHEP	-49%	-24%	-29%
<u>PHEP - IdF</u> IdF	-31%	-21%	-30%
<u>L-EN - IdF</u> IdF	-65%	-40%	-50%

717

718 Table 5: C sequestration (t CO<sub>2</sub>.eq ha<sup>-1</sup>) simulated over 25-year and 50-year periods for the different cropping systems  
 719 with the ROTH C model (V.26-3) and the SIMEOS® tool (2010), for mean soil organic matter content in the Ile-de-  
 720 France region (1.6%. SOM = 123.3 t CO<sub>2</sub>.eq ha<sup>-1</sup>). Cropping systems: PHEP (productive high environmental  
 721 performance), L-GHG (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).

C sequestration (t CO <sub>2</sub> .eq ha <sup>-1</sup> )		Assessment period	Assessment period
		25 years	50 years
Cropping systems	Tool or model		
PHEP	SIMEOS® tool	15.1	19.7
PHEP	ROTH C model	85.2	116.0
L-GHG	SIMEOS® tool	32.0	40.5
L-GHG	ROTH C model	102.9	147.8
IdF	SIMEOS® tool	17.4	22.1
IdF	ROTH C model	87.1	129.2
<u>(L-GHG – PHEP)</u>			
PHEP SOC	SIMEOS® tool	12%	15%
<u>(L-GHG – PHEP)</u>			
PHEP SOC	ROTH C model	9%	13%

722

723 Table 6: Comparison of total simulated greenhouse gas emissions, expressed in t CO<sub>2</sub>-eq ha<sup>-1</sup> and t CO<sub>2</sub>-eq t<sup>-1</sup>, between  
 724 cropping systems. Cropping systems: PHEP (productive high environmental performance), L-GHG (low greenhouse  
 725 gas emissions) and IdF (current system in Ile-de-France region).

	Total greenhouse gas emissions (t CO <sub>2</sub> -eq ha <sup>-1</sup> )	Total greenhouse gas emissions (t CO <sub>2</sub> -eq t <sup>-1</sup> )
<u>L-GHG - PHEP</u> PHEP	-13%	-22%
<u>PHEP - IdF</u> IdF	-46%	-32%
<u>L-GHG - IdF</u> IdF	-53%	-47%

726

727 Table 7: Greenhouse gas balances (t CO<sub>2</sub>-eq ha<sup>-1</sup>), for different cropping systems, determined with the SIMEOS® tool  
 728 (2010) and the ROTH C model (V.26-3), for a current soil in the Ile-de-France region (SOM = 1.6%) and the  
 729 GES'TIM<sup>37</sup> database, for 25-year and 50-year periods. The GHG balance is positive when the amount of carbon emitted  
 730 in greenhouse gases exceeds that sequestered. Cropping systems: PHEP (productive high environmental performance),  
 731 L-GHG (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).

Greenhouse gas balances (t CO <sub>2</sub> -eq ha <sup>-1</sup> ),		Assessment period	Assessment period
		25 years	50 years
Cropping system	Tool or model		
PHEP	SIMEOS® tool	16.7	44.0
PHEP	ROTH C model	-53.4	-52.3
L-GHG	SIMEOS® tool	-4.4	14.4
L-GHG	ROTH C model	-75.3	-92.6
IdF	SIMEOS® tool	41.9	96.4
IdF	ROTH C model	-39.7	-10.8
L-GHG – PHEP	SIMEOS® tool	-21.1	-29.3
L-GHG – PHEP	ROTH C model	-21.9	-40.3

732

733 Table 8: Greenhouse gas balance ratio of the various cropping systems, expressed in t CO<sub>2</sub>-eq ha<sup>-1</sup> , simulated with the  
734 ROTH C (V.26-3) model over two different periods (25 and 50 years). Cropping systems: PHEP (productive high  
735 environmental performance), L-GHG (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).

	GHG balance: 25 years	GHG balance: 50 years
L-GHG / PHEP	1.51	1.76
PHEP /IdF	1.39	4.95
L-GHG / IdF	2.10	8.72


736

737






745 Figure 2: Cropping system prototypes assessment with GESTIM database: Mean total greenhouse gas emissions  
 746 (expressed in kg CO<sub>2</sub>-eq ha<sup>-1</sup> year<sup>-1</sup>) calculated over one rotation period for the different cropping systems, and total  
 747 greenhouse gas emissions for each crop of the various cropping systems (i.ghg: indirect greenhouse gas – d.ghg: direct  
 748 greenhouse gas- w: winter - s: spring). Cropping systems: PHEP (productive high environmental performance), L-GHG  
 749 (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).



750

751 Figure 3: Environmental criteria assessment of the different cropping system prototypes with the INDIGO® tool  
 752 (v.1.9). Cropping systems: PHEP (productive high environmental performance), L-GHG (low greenhouse gas  
 753 emissions), L-EN (low energy), No-Pest (No pesticides) and IdF (current system in Ile-de-France region). Minimum  
 754 values to satisfy environmental targets.



755 — PHEP — L-EN - - - L-GHG — No Pest ..... IdF x Min values to satisfy environmental targets