



Designing innovative productive cropping systems with quantified and ambitious environmental goals

Caroline Colnenne-David, Thierry Doré

► To cite this version:

Caroline Colnenne-David, Thierry Doré. Designing innovative productive cropping systems with quantified and ambitious environmental goals. Renewable Agriculture and Food Systems, 2015, 30 (06), pp.487-502. 10.1017/S1742170514000313 . hal-01354945

HAL Id: hal-01354945

<https://agroparistech.hal.science/hal-01354945>

Submitted on 7 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 doi:10.1017/S1742170514000313

2

3 **Designing innovative productive cropping systems with quantified and ambitious**
4 **environmental goals**

5

6 Colnenne-David Caroline¹, Doré Thierry²

7 ¹ INRA, UMR 211 Agronomie, 78850 Thiverval-Grignon, France, e-mail:

8 caroline.colnenne@grignon.inra.fr

9 ² AgroParisTech, UMR 211 Agronomie, 78850 Thiverval-Grignon, France e-mail:

10 thierry.dore@agroparistech.fr

11

12

13

14

15

16

17

18

19

20

21

22

23 Corresponding author: COLNENNE-DAVID Caroline

24 INRA, UMR 211 Agronomie, 78850 Thiverval-Grignon, France

25 Phone: (+33) 1.30.81.53.28

26 Fax: (+33) 1.30.81.54.25

27 e-mail: caroline.colnenne@grignon.inra.fr

28

29 **Abstract**

30 Agriculture must face a number of very pressing environmental issues. We used the prototyping method to design three
31 innovative cropping systems, each satisfying three ambitious goals simultaneously: (i) overcoming a major
32 environmental constraint, which represents a major break regarding objectives to be reached in current cropping
33 systems (differing between systems: a ban on all pesticides but with chemical N fertilizer permitted; reducing fossil
34 energy consumption by 50%; or decreasing greenhouse gas emissions by 50%), (ii) meeting a wide range of
35 environmental criteria, and (iii) maximizing yields, given the major constraint and environmental targets. A fourth
36 cropping system was designed, in which the environmental and yield targets were achieved with no major constraint
37 (PHEP system). The performances of these innovative cropping systems were compared to a conventional system in the
38 Ile-de-France region. We used a three-step prototyping method: (1) new cropping systems were designed on the basis of
39 scientific and expert knowledge, (2) these system prototypes were assessed with tools and a model (*ex ante* assessment)
40 adjusted to the set of constraints and targets, with optimization by an iterative process until the criteria were satisfied
41 and (3) evaluation in a long-term field experiment (*ex post* assessment), which is currently underway. We describe only
42 the first two steps here, together with the results of the prototypes assessment with tools and a model. The pesticide,
43 energy and greenhouse gas constraints were fulfilled. All these innovative systems satisfied environmental criteria in
44 terms of nitrogen and phosphorus management, pesticide use, energy consumption and crop diversity. For the pesticide-
45 free system, the soil organic matter indicator was lower than expected due to frequent plowing (every two years) and
46 yields were 20 to 50% lower than for the PHEP system, depending on the crop considered. We focus our discussions on
47 the design methodology and the availability of scientific knowledge and tools for projects of this type.

48

49 **Keywords**

50 Innovative cropping systems, environmental goals, *ex ante* assessment.

51

52 **Introduction**

53 Agriculturalists are faced with challenges relating to very pressing environmental and health issues, including the need
54 to decrease pesticide use. In many countries, high levels of pesticides have frequently been found in rivers, lakes and
55 groundwater^{1,2}. A second pressing environmental issue is the consumption of fossil fuel. Energy use has markedly
56 increased over the last decade³, and some scientists agree that oil availability will decline in the near future⁴, leading to
57 a sharp increase in oil prices⁵. In this context, new ways of optimizing or reducing energy use have been proposed⁶.
58 Global warming is a third challenging environmental issue facing agriculture. About 12% of global greenhouse gas
59 (GHG) emissions emanate from agricultural lands⁷, and this proportion is expected to rise in the future, due to increases
60 in the amount of land used for agricultural purposes and the intensification of agricultural practices⁸. Carbon (C)
61 sequestration in the soil, through the return of crop residues, root deposition and organic amendments, may help to
62 decrease GHG emissions⁹. Sustainable development is another pressing social issue. Sustainable agriculture must
63 satisfy environmental criteria¹⁰. The harmful impact of agriculture on the environment can be lessened by optimizing
64 fertilization (N, P) and increasing crop diversity. Currently pesticide use and energy consumption should also be
65 reduced^{1,2}, and soil fertility should be maintained. Since the 1950s, alternative crop management systems have been
66 proposed¹¹ and legislation and inspection services have controlled the use of inputs (pesticides, N fertilization). Finally,
67 agricultural production must satisfy the food needs of a soaring world population¹⁰. Global agricultural production
68 currently feeds a population of approximately seven billion. Current projections suggest that the world population will
69 have reached nine billion by 2050¹². The resulting increase in the need for land for housing will reduce the amount of
70 land available for agricultural purposes¹³. The availability of arable land per capita differs greatly between regions (*e.g.*,
71 between China and South America), and major cropping systems must take this scarcity into account. Foley¹⁴ have
72 suggested that feeding a population of this size will be possible only if agricultural systems change, along with human
73 eating habits.

74 Many studies in recent years have focused on the design and assessment of new cropping systems. New crop
75 management strategies have been proposed to decrease pesticide use¹⁵, to decrease energy consumption¹⁶, or to enhance
76 C sequestration through crop management practices¹⁷. Energy use and greenhouse gas emissions have been calculated
77 and assessed for different systems¹⁸. At the cropping system level, long-term trials also have been set up to investigate
78 the effects of different technical operations, such as N fertilization, on soil physical and chemical properties¹⁹ and on
79 soil biology²⁰. Several studies have assessed the differences between organic, integrated and conventional cropping
80 systems in terms of C sequestration²¹, energy efficiency and use²², profitability²³ or productivity²⁴. Some authors have
81 analyzed the impact of different degrees of tillage on productivity²⁵ or biological activity²⁶. Others have focused on the
82 effect of cropping systems on biodiversity²⁷, or have used life cycle assessment methods to analyze the sustainability of

various farming systems²⁸. In most of these cases, new cropping systems were designed by modifying a few agricultural practices targeting a single goal (*e.g.*, no chemicals to be used in organic cropping systems; no ploughing to increase C sequestration; more legumes in the rotation to reduce energy consumption; and more inputs to enhance profitability), without considering the other dimensions of sustainability. Despite the evidence that the future of agriculture must address a wide range of issues, no study has designed innovative cropping systems with specific and quantitative objectives covering a broad range of issues.

The objective of our project was to design, by prototyping²⁹, innovative cropping systems meeting three quantitative objectives: (i) to satisfy a major environmental constraint, which represents a major break regarding objectives to be reached in current cropping systems (the banning of pesticide use, reducing fossil energy consumption by 50%, or reducing gas emissions by 50%); (ii) to satisfy a wide range of environmental criteria with specific quantitative targets, and (iii) to produce the maximum yield possible given the constraint and the environmental targets. The ultimate aim of this work is to improve arable cropping systems throughout northern Europe. Prototyping was developed to enable agronomists to design, test and improve more sustainable cropping systems²⁹. With this approach, newly designed cropping systems could satisfy several of the issues mentioned above and contributed to identify the weaknesses of cropping systems. System prototypes were assessed with tools and a model (*ex ante* assessment), with discussion of their potential performances, before their assessment in a long-term field experiment (*ex post* assessment). We set up this long-term field experiment in 2008, and its results will be published in due course. We focus here on the prototyping and assessment of the cropping system prototypes. We discuss our results in terms of innovative design methodology, the innovation of agricultural practices, the availability of suitable tools, models and crop management, and the yields achievable.

1. Materials and methods

1.1. Design method

The method used for cropping system design was based on the prototyping approach^{29,30,31,32}, which involves four major steps:

- (i) Defining and ranking the constraints and targets;
- (ii) Designing innovative cropping system prototypes on the basis of current knowledge;
- (iii) Assessments of cropping system prototypes with tools and models adapted for the constraints and targets used, with improvement of the cropping systems (in terms of rotation or crop management aspects) by an iterative process, until satisfaction of the constraints or achievement of results considered the best possible; and

(iv) Assessment of the most promising cropping system candidates in a long-term field trial. This practical assessment is currently being carried out in a long-term field experiment, initiated in 2008.

1.2. Constraints and targets for innovative cropping systems

Four different cropping systems with quantified constraints (*i.e.*, conditions that had to be fulfilled), environmental and yield targets, were designed for the agricultural conditions and principal crops of northern France. These constraints and targets were prioritized as follows: an environmental constraint had to be satisfied first; a set of environmental targets then had to be attained, and finally, yield had to be maximized. The quantitative levels of the constraints did not correspond to any regulations and reflected a major break to be reached in current cropping systems. Inclusion of the use of organic fertilizers (manure, compost, etc.), which are currently not readily available to many farmers in large areas of Western Europe, was not permitted in the design of the cropping systems.

1.2.1. The productive high environmental performance cropping system (PHEP).

No major environmental constraint was placed on this cropping system, which was designed to reach environmental targets. Eleven environmental indicators, according to the INDIGO® tool³³, were used to assess the effects of the cropping system on groundwater pollution (nitrate and pesticides), crop diversity, energy use and soil quality (organic matter content and phosphorus concentration). To reach environmental goals, all these environmental indicators, calculated over an entire crop rotation sequence, had to have values higher than 7 (graduated scale from 1 to 10)³³. This system was used as the reference system for comparisons with the other three systems.

1.2.2. No-pesticide cropping system (No-Pest).

This cropping system was subject to a specific pesticide constraint: no pesticide use was tolerated, even using substances (e.g. acetic acid) at levels usually considered acceptable in organic cropping systems. However, inorganic chemical fertilizers were allowed (these fertilizers are not permitted in organic farming systems). This system had to achieve the same environmental targets as the PHEP cropping system.

1.2.3. Low energy cropping system (L-EN).

This cropping system was subjected to a specific energy constraint: it had to have fossil fuel consumption levels no greater than half those of the PHEP cropping system. It had to reach the same environmental targets as the PHEP cropping system.

1.2.4. Low greenhouse gas emission cropping system (L-GHG).

145 This cropping system was subject to a specific constraint concerning greenhouse gas emissions: its greenhouse gas
146 emissions had to be no more than half those of the PHEP cropping system by increasing carbon sequestration in the soil
147 and decreasing N₂O emissions. It had to meet the same environmental targets as the PHEP cropping system.

148

149 For each cropping system, once the constraint had been satisfied and environmental targets had been reached, the
150 combination of agricultural practices giving the highest yields was retained.

151

152 **1.3. Design of the four innovative cropping systems**

153 The innovative cropping systems were designed from published knowledge, quantitative data from field experiments
154 and individual or group expertise provided by scientists, extension service staff and farmers. For each cropping system,
155 one prototype, consisting of the species in the rotation and the combination of agricultural practices used, was designed.
156 If the constraints were not satisfied, the candidate was modified iteratively (changes to the crops in the rotation or
157 agronomic practices) until they were. At the beginning of the process, a modification of a crop led to a multitude of
158 changes; at the end, changes were only one at a time. When the constraints were satisfied, environmental targets were
159 optimized by an iterative procedure until improvement was observed. Maximum achievable yields were then
160 determined by experts knowledge or from trial results, for the various cropping systems. The candidate cropping
161 systems selected for further assessment in a field experiment were, those with the best performances in terms of
162 constraints, environmental targets and achievable yields.

163 We carefully selected agronomic strategies from previous publications, to satisfy the given constraints. Examples of
164 such strategies are presented in Table 1. Current knowledge, based on conventional cropping systems, had to be adapted
165 for innovative cropping systems, and it was necessary to combine strategies. Agronomic strategies were translated into
166 decision rules (as described by Debaeke³⁴) to meet the requirements of future cropping systems and to cope with the
167 variability of weather and agronomic conditions.

168

169 **1.4. Assessment with tools and a model and fine-tuning of innovative cropping system prototypes**

170 During the design process, the cropping system prototypes were assessed with various tools and a model, to determine
171 the best ways to satisfy the set of constraints and targets imposed. Direct and indirect non-renewable energy
172 consumption was assessed with the INDIGO® tool (v. 1.9). Direct energy consumption concerned the fuel, lubricants
173 and electricity used to power farm machinery and tractors. Indirect energy consumption concerned the energy used in
174 the manufacture, formulation, packaging and maintenance of inputs, such as machinery, fertilizer or pesticides. The
175 energy outputs of the cropping systems were calculated as the gross energy content of the harvested produce. Energy

176 consumption was calculated on a per hectare basis, per tonne of crop product and per calorie produced, over a complete
177 crop sequence.

178 C sequestration in the soil was assessed with (i) the Roth C 26.3 model³⁵ and (ii) the SIMEOS® tool (v.2010) based on
179 the AMG model³⁶. We used climatic data (*i.e.*, monthly mean air temperature, monthly precipitation, and monthly open
180 pan evaporation) from a meteorological station located in Grignon (Ile-de-France region, 30 km west of Paris). The soil
181 characteristics (plow layer, 0-30 cm) used to drive simulations were as follows: clay content 20.6%, bulk density 1.4,
182 initial C content 8 g/kg dry matter. The expected annual yields were estimated from experimental data obtained under
183 the same conditions (*i.e.*, Ile de France region) and adjusted by expert knowledge. These values were used to estimate
184 the expected annual dry matter production of roots and stubble, as described by Van Groenigen et al.³⁷. Direct and
185 indirect GHG emissions were estimated with the GES'TIM database³⁸. We focused on two main greenhouse gases:
186 nitrous oxide (N₂O) and carbon dioxide (CO₂). Direct emissions included N₂O emissions from N fertilizers, calculated
187 with Intergovernmental Panel on Climate Change coefficients³⁹, and the CO₂ produced by the combustion of fossil fuels
188 by farm machinery; CO₂ respired by soil organisms was not taken into account in calculations. Indirect emissions
189 corresponded to the use of fossil energy in the manufacture and maintenance of farm inputs. GHG balances (C
190 sequestration plus GHG emissions) were determined over periods of 25 and 50 years, in accordance with
191 Intergovernmental Panel on Climate Change proposals³⁹ and current knowledge of C sequestration kinetics in the soil.
192 In this investigation, any GHG entering the system is counted negatively whereas GHG leaving the system is counted
193 positively. Therefore, the overall balance is a positive value if more greenhouse gases are emitted than sequestered in
194 the system.

195

196 Environmental indicators, such as in the INDIGO® tool (v. 1.9), were used to assess the environmental effects of
197 cropping system prototypes. Three indicators of nitrogen effects provided information about ammonia volatilization,
198 nitrous oxide emissions into the air, and nitrate leaching into the groundwater. Four pesticide indicators were studied:
199 three providing information about pesticide volatilization, pesticide runoff and pesticide leaching into groundwater and
200 one taking the global effect of pesticides into account. The last four indicators used provided information about crop
201 diversity, energy consumption, organic matter in the soil, and phosphorus management. Each indicator takes a value
202 between 1 (worst) and 10 (best). For rotations of more than five crops, the crop diversity indicator was calculated from
203 the coefficients of Leteinturier⁴⁰. For example, values 0.5, 4.1 and 7.6 respectively correspond to a wheat monoculture,
204 a wheat-maize rotation and a wheat-sunflower-spring barley-maize rotation.

205 All these tools and the model were chosen on the basis of their relevance for assessing compliance with constraints and
206 environmental targets. In a comparison of the performance of nine soil organic C models, using different datasets from
207 long-term experiments from different parts of the world, Smith⁴¹ found that the RothC model was among those that

performed best. Bockstaller⁴² analyzed four methods for assessing the sustainability of agricultural systems. They found that the INDIGO® tool was the most relevant for conditions corresponding to those used here. These tools and the model have been regularly used in different countries. For example, Roth C has been used by De Li Liu⁴³ and Cerri⁴⁴, and INDIGO® has been used by Bockstaller⁴².

1.5 Current cropping system in the Ile-de-France region

The current system in the Ile-de-France region was defined on the basis of data collected in 2006 (Agreste⁴⁵), the most recent data available at the initiation of this program. We defined the current cropping system in terms of agronomic practices and crop descriptions. This system was validated by various experts (farmers, extension service staff) with few adjustments in terms of types and numbers of crops in the rotation. This system was used as a reference for further comparisons.

2. Results

2.1. Design methodology step

The systems were designed over a six-month period, by a panel of about 15 experts. For each innovative cropping system, the first candidate characteristics were based on the current cropping system in the Ile-de-France region. The system was then optimized through an iterative process, which produced approximately 70 prototypes, to find the four most promising candidates. These candidates corresponded to the prototypes satisfying the constraint and environmental targets imposed and yielding the best results in the assessment. For example, for the PHEP system, the value of the crop diversity indicator was gradually increased from 5 to 7 during the fine-tuning of the system, with simultaneous improvement of the values of the other indicators. The first prototype was based on a 3-year rotation (winter oilseed rape, winter barley and winter wheat), currently used in the Ile-de-France region. In the best prototype, a winter legume and spring barley with a mustard catch crop were gradually introduced, leading to the following rotation: winter field beans (*Vicia faba*), winter wheat, winter oilseed rape, winter wheat and spring barley with a mustard catch crop. In the design process, we began by determining the crop rotation and then defined the crop management practices.

2.2. Description of the innovative cropping system prototypes

For each cropping system, we present only the most promising prototype. The systems are first described in terms of the crop rotation, crop management practices and yield targets. We then present the results of the final assessment with respect to constraints (*i.e.*, pesticide use, energy consumption and greenhouse gas emissions) and, finally, we evaluate the systems in terms of environmental targets. The crop rotations and targeted yields are presented in Tables 2 and 3.

239

240 **2.2.1. The productive high environmental performance cropping system (PHEP)**

241 The PHEP cropping system was designed with multiple environmental targets in mind and was based on the following
242 agronomic strategies: (i) to reduce pesticide use, we increased crop diversity (four different crops instead of the three
243 currently sown), (ii) to reduce the amount of N used and indirect energy consumption, we included at least one legume
244 in the rotation, (iii) to decrease nitrogen leaching, a catch crop was always sown before the spring crop and N
245 fertilization was forbidden during autumn and winter, (iv) to reduce direct energy consumption, plowing was allowed
246 only once in the rotation, before the spring crop, (v) to reduce pesticide use and crop loss due to insects and diseases,
247 highly resistant varieties were used, together with optimal sowing dates and densities and (vi) to stabilize or/and to
248 enrich the soil organic matter (SOM) content of the soil, crop residues were not removed. As the system had to satisfy
249 environmental targets requiring the use of fewer inputs, the target yields set were similar to those currently achieved
250 with low-input cropping systems in the Ile-de-France region.

251

252 **2.2.2. The no-pesticide cropping system (No-Pest)**

253 Pesticide use was prohibited in the “No-Pest” cropping system. Therefore, this cropping system was designed as
254 follows: (i) to break the cycles of some common soil-borne pathogens, we used a long rotation including a range of
255 species (five different crops), with the alternate sowing of host and non-host plants, (ii) to reduce weed emergence from
256 year to year, we sowed species with different sowing dates in spring and in winter successively, (iii) to decrease pest
257 and disease pressure and damage, we used highly resistant varieties and species mixtures, and excluded crops highly
258 susceptible to some enemies but with few non-chemical solutions, such as oilseed rape or potatoes, from the rotation
259 (iv) to increase the competitiveness of the crop with respect to weeds, we sowed species with rapid shoot growth, such
260 as hemp and triticale, (v) to maximize weed emergence before sowing, we used the stale seed-bed technique, (vi) to
261 reduce weed emergence after sowing, plowing was carried out before each spring crop and (vii) we adapted sowing
262 densities to make it possible to use mechanical weeding techniques and to decrease pathogen propagation. We used the
263 following approaches to reach environmental targets: (i) to reduce nitrate leaching, catch crops were always sown
264 before spring crops and the spreading of nitrogen fertilizer was allowed only in the spring, (ii) to decrease direct and
265 indirect energy consumption, we decreased the number of plowing events and N fertilization was calculated according
266 to yield objectives, and (iii) to stabilize SOM, crop residues were not removed. Yield targets were lower than those for
267 the PHEP cropping system, because no pesticides were used. However, they were higher than those achieved in organic
268 systems because chemical fertilizers were allowed, increasing flexibility in the management of crop nitrogen nutrition.
269 For the integration of these features, in accordance with current knowledge of pest and disease pressures in the Ile-de-

270 France region, experts suggested yield potentials 30% lower than those for the PHEP system for cereals and 25% lower
271 for field beans.

272

273 **2.2.3. Low energy cropping system (L-EN)**

274 The L-EN cropping system was designed, to have a much lower energy consumption than the PHEP cropping system,
275 as follows: (i) to reduce indirect fuel consumption due to N fertilization, we included as many legumes as possible in
276 the rotation (field beans as a main crop, clover as a catch crop, and a white clover-winter wheat mixture), and we used
277 species or varieties with high N use efficiency (*e.g.*, oats ⁴⁶) and forms of mineral N fertilizers requiring less energy for
278 their manufacture, (ii) to decrease direct fuel consumption, we omitted plowing, which is a very resource-intensive
279 operation, and used a direct drilling system, and (iii) we decreased the amounts of mineral fertilizer (N, P, K) applied,
280 implying a decrease in target yields. We also designed the L-EN cropping system along the same lines as the PHEP
281 system, to achieve environmental targets for crop diversity, length of rotation, date of nitrogen spreading, and catch
282 crop sowing. Target yields were 20% lower than for the PHEP cropping system, except for field beans.

283

284 **2.2.4. Low greenhouse gas emission cropping system (L-GHG)**

285 The L-GHG cropping system was designed to decrease greenhouse gas emissions by increasing C sequestration in the
286 soil and decreasing N₂O emissions.

287 C sequestration in the soil was increased by (i) including as many cereals as possible in the rotation, to ensure the
288 production of large amounts of residues (*i.e.*, maize, winter wheat, winter barley or triticale), (ii) maintaining
289 continuous soil cover to increase the amounts of organic residues (*i.e.*, cover or catch crops were always sown between
290 main crops, and volunteers were left to grow after harvest), (iii) targeting high yields for the main and catch crops, to
291 ensure the production of large amounts of residues, and (iv) excluding moldboard plowing, which increases C
292 mineralization.

293 N₂O emissions were reduced by (i) decreasing the amount of N fertilizer required at rotation scale⁴⁷, and consequently
294 direct emissions of N₂O, by sowing legumes in the rotation (main and catch crops), (ii) improving and optimizing N
295 fertilization practices according to climatic conditions, through the use of appropriate decision rules to prevent
296 applications in conditions favoring N₂O emissions, and (iii) sowing species with taproots to reduce soil compaction and
297 N₂O emissions.

298 The L-GHG cropping system was also designed according to the same principles as the PHEP system, to reach
299 environmental targets for crop diversity, length of rotation, pesticide use, date of N spreading, and catch crop sowing.

300 Target yields were considered to be a compromise between the production of large amounts of C residues (*i.e.*, high

301 yields) and the decrease in N₂O emissions (*i.e.*, low N fertilization). Experts thought that potential yields would be
302 similar to those achieved by the PHEP cropping system.

303

304 **2.2.5. The current cropping system in the Ile-de-France region**

305 This system is based on a cereal crop rotation, with five cereal crops over a six-year rotation (Table 2). In order to
306 secure high yields, the agronomic practices were as follows: regular plowings, four times over a 6-year rotation. The
307 amounts of N fertilizer exceed crop requirements, to prevent yield shortfalls in the event of unfavorable climatic
308 conditions or unexpected nitrogen losses. Pesticides and growth regulators were used liberally to prevent diseases,
309 weeds, pests and lodging (3 to 5 pesticides every year).

310

311 **3. Cropping system prototypes assessments with tools and a model**

312 **3.1. Constraint assessment**

313 **3.1.1. Pesticide constraint in the No-Pest cropping system**

314 This constraint was achieved by not applying pesticides in the No-Pest cropping system.

315

316 **3.1.2. Energy constraint in the L-EN cropping system**

317 Mean total fossil energy consumption (direct and indirect energy), calculated over a single rotation, was 4517 MJ ha⁻¹
318 year⁻¹ for the L-EN system and 8826 MJ ha⁻¹ year⁻¹ for the PHEP system (Figure 1). Chemicals, including N fertilizers,
319 the largest component, accounted for 1271 MJ ha⁻¹ year⁻¹ (43% of total indirect energy consumption) in the L-EN
320 system and 4345 MJ ha⁻¹ year⁻¹ (95% of total indirect energy consumption) in the PHEP system. The use of machinery
321 for tillage, fertilization, harvesting, sowing and crop protection was the only component of direct energy consumption
322 that was nearly halved in the L-EN system (2976 MJ ha⁻¹ year⁻¹ and 4228 MJ ha⁻¹ year⁻¹ for the L-EN and the PHEP
323 systems, respectively). The difference between these two systems can be accounted for by the absence of tillage and the
324 use of less N fertilizer in the L-EN cropping system.

325 When expressed in MJ ha⁻¹, the total fossil energy in the L-EN system is 49% lower than that in the PHEP system
326 (Table 4). However, if expressed in MJ t⁻¹, the energy performance of the L-EN system is lower (*i.e.* difference between
327 the PHEP and the L-EN systems of only 24% in term of total fossil energy per tonne of produce), because the target
328 yield is about 20% lower. A similar reduction in energy use (about 29%) was observed for the calculation in kJ kcal⁻¹.

329

330 **3.1.3. The greenhouse gas constraint in the L-GHG cropping system**

331 C sequestration was assessed for the optimized prototypes of the PHEP and the L-GHG systems, for the mean soil
 332 organic matter content in the Ile de France region (1.6%). Both the Roth C model and the SIMEOS® tool predicted that
 333 C would be sequestered throughout the study period, from the start, in both cropping systems. The highest values were
 334 obtained with the L-GHG system over a 50-year period, for both assessment tools (Table 5). For both systems, total C
 335 sequestration was higher during the first 25-year period than during the second 25-year period. When expressed in t
 336 CO₂-eq ha⁻¹, C sequestration values were systematically higher with the Roth C model than with the SIMEOS® tool.
 337 Nevertheless, after 25 and 50 years, the differences between the L-GHG and the PHEP systems calculated with the Roth
 338 C model and the SIMEOS® tool were similar if the results were expressed in relative values.
 339 Direct and indirect greenhouse gas emissions were calculated with the GES'TIM database, over one rotation period, for
 340 the L-GHG and the PHEP cropping systems (Figure 2). Mean total greenhouse gas emissions were 1104 kg CO₂-eq ha⁻¹
 341 year⁻¹ and 1273 kg CO₂-eq ha⁻¹ year⁻¹ for the L-GHG and the PHEP cropping systems, respectively. Direct and indirect
 342 greenhouse gas emissions accounted for similar proportions of total emissions: 48% and 52% for direct greenhouse gas
 343 emissions for the L-GHG and PHEP systems, respectively (Table 6). Chemical fertilizers caused both direct and
 344 indirect greenhouse gas emissions. They represented 76% and 73% of total greenhouse gas emissions for the L-GHG
 345 and the PHEP systems, respectively. Soil cultivation, accounting for 19% and 23% of total greenhouse gas emissions
 346 for the L-GHG and the PHEP systems, respectively, was the second most important component of these emissions.
 347 When results were expressed per ha, total greenhouse gas emissions were 13% lower in the L-GHG system than in the
 348 PHEP system. When expressed per tonne of produce, the larger decrease (22%) may be accounted for by the higher
 349 yields, calculated at rotation scale, of the L-GHG system than of the PHEP system.
 350 In terms of the overall balance of GHG emissions (Tables 7-8), GHG values were negative for the L-GHG system,
 351 except for the 50-year period with the SIMEOS® tool. All GHG balance values were lower for the L-GHG system than
 352 for the PHEP system. The difference in GHG balance between the two systems increased over time, and was greater
 353 with the Roth C model, which gave decreases in GHG emission of 51% for the 25-year period and 76% for the 50-year
 354 period.

355

356 **3.2. Assessment of environmental targets**

357 The results of assessments of environmental targets with the INDIGO® tool are shown in Figure 3. For all optimized
 358 cropping systems, all 11 indicators had values of at least 7 (*i.e.*, environmental criteria were satisfied), except for the
 359 organic matter indicator for the No-Pest cropping system (OMI = 5.7). The large number of species (more than three in
 360 each rotation) and the small quantities of pesticides sprayed on crops (0 to 2 pesticides used per crop), the systematic
 361 restitution of residues and the optimization of tillage and fertilization management (*i.e.*, the small number of ploughing
 362 operations, optimizing P and N fertilization in terms of both the amounts applied and the timing of applications),

363 resulted in high values for the indicators for crop diversity, pesticide use, soil organic matter, phosphorus, nitrogen and
364 fossil fuel, respectively.
365 The low value of the organic matter indicator for the No-Pest cropping system (5.7) could be accounted for by both the
366 large number of plowing operations (alternate years), encouraging mineralization, and the lower yields, resulting in
367 smaller amounts of C residues.

368

369 **4. Discussion**

370 **4.1. Design and assessment of innovative cropping systems**

371 The main challenge of this study was to design innovative cropping systems. Our approach is original in the multiplicity
372 of purposes assigned to these systems (*i.e.*, association of one major constraint with environmental and yield targets). In
373 most previous studies, these issues have been analyzed separately. For example, Zentner⁴⁸ and Gefland⁴⁹ studied energy
374 efficiency, whereas Nemecek²⁸ used life cycle assessment methods to evaluate environmental criteria, Nowacki⁵⁰
375 studied profitability, and Chikowo⁵¹ studied new cropping systems with a lower reliance on pesticides. However,
376 several other studies are currently investigating system sustainability including assessments of several different
377 criteria⁵², or numerous environmental parameters²². In our project, we combined one major constraint with
378 environmental and yield targets, reflecting the multifunctionality of agriculture. Furthermore, there was a clear,
379 particular hierarchy throughout the design process. In most published experiments, environmental consequences are
380 assessed only during the assessment of technical innovations in the trials, or environmental goals exist but are not
381 quantified at the start of the study. In our work, satisfying the major constraints and the precise environmental targets
382 were major aims, which became the conditions determining yield, with target yields set as high as possible under the
383 conditions concerned. In addition, the clear definition of the constraint (*i.e.*, reducing energy consumption or GHG
384 emissions by 50%) and the environmental targets (*i.e.*, having a value of at least 7 for all INDIGO environmental
385 indicators) was also original. The quantitative levels of the constraints did not correspond to any regulations (*i.e.*, these
386 constraints reflected a major break with the regulations). However these innovative cropping systems were considered
387 as research tools which enabled to identify the most relevant agronomic practices combinations which could be used in
388 more restrictive legislative contexts. The level quantifications of the constraints and environmental targets were very
389 useful during the design process which required calculations.

390 Before the assessment of the prototypes in a long-term field trial, candidate systems were assessed and improved in an
391 iterative process until the constraints were satisfied and environmental performance with respect to targets was
392 optimized. This theoretical process of improvement has rarely been reported in previous studies. Cropping systems are
393 usually assessed or compared in systems defined on the basis of the main standardized characteristics, essentially

relating to one major aspect: *e.g.*, organic *versus* conventional systems^{53,54,24}, no-tillage *versus* conventional tillage^{54,55}, or integrated *versus* conventional systems^{50,56}. Quantitative data for environmental criteria²⁶, yield⁵⁷ or economic performance^{58,59} were therefore recorded in experiments. The results of these comparisons can be used to compare the impact of different systems, but not to identify all solutions for their improvement. Even though inductive reasoning can bring about some conclusions in regards to general principles, another round of conception and evaluation is required, to strengthen cropping systems. Our prototyping approach is totally different. Innovative systems were assessed by modeling until they satisfied specific constraints and were optimized in terms of specific environmental targets. A field trial was then set up to determine whether each of the selected prototypes could satisfy its multiple constraints and targets. In this case, the various environmental targets were included in the agricultural strategies from the start of the design process, facilitating identification of the weaknesses of the system and making it possible to propose solutions for improvement before undertaking field trials. After the assessment of these innovative cropping systems in a field trial, their costs and economic performances will be calculated in different economic scenarios, to determine the likelihood of their being adopted by farmers.

Our approach required a large panel of experts (scientists, farmers, and extension service staff) to design and to support prototypes throughout the design process (*i.e.*, from the first to the last candidates). This was necessary because (i) the best crop management system may not correspond simply to the sum of individual agricultural practices, but may instead involve a set of agronomic strategies and their interactions, and (ii) a breadth of agro-ecological knowledge is required to identify sets of agricultural rules likely to be responsive to such strict constraints and environmental targets. Moreover, this approach provided a more realistic view of cropping systems, making the adoption of the proposed innovations more likely⁶⁰. However, the field trial assessment step is still absolutely necessary because some innovative agronomic practices, not currently used in cropping systems, have never been evaluated by experts.

During the design process, about 15 experts attended individual sessions or group meetings, to provide knowledge unavailable from published work. The definition of crop rotations and agronomic practices took about six months, and a further 18 months were required for the writing of the decision rules. Published studies involving design processes have differed considerably in the number of experts involved and the time spent by individual experts, depending on the availability of the experts and the difficulties encountered in achieving the goals assigned to systems^{52,61}.

420

421 **4.2. Achievement of multiple constraint and targets**

For all innovative systems, the constraints were satisfied with no consequences for other environmental components, except for the organic matter indicator of the No-Pest system. In this case, regular tillage combined with the restitution of only small amounts of organic matter had an adverse effect on soil environmental characteristics (indicator value for soil organic matter of 5.7, according to the INDIGO® tool). Within this system, it did not appear to be possible to

425

satisfy both the constraint and this environmental target with the available non-chemical techniques for pest control. Moreover, this was only possible with the available techniques by reducing yield targets with respect to those of current regional systems (Agreste⁴⁵). Nevertheless, progress in integrated pest management is being made, and new techniques may make it possible to improve environmental and yield performances. In the design of the L-EN system, we managed various agricultural processes, decreasing both direct energy consumption (due to tillage, for example) and indirect energy consumption (due to the use of mineral fertilizers). We halved fossil energy consumption by greatly decreasing N fertilizer inputs, which was associated with a 20% yield loss. However, the energy performance of the L-EN system was expressed relative to that of the PHEP system, which also had a relatively low level of fossil fuel consumption with respect to current practices in Ile-de-France. The total energy consumption of the L-EN system was about 35% that of the current system in Ile-de-France (Agreste⁴⁵), when energy was expressed in MJ ha⁻¹ (Table 4). For both the L-GHG and the PHEP systems, decreases in pesticide were taken into account by considering the maximum achievable yields to be similar to those of current low-input systems in Ile-de-France (Agreste⁴⁵), and much lower than those of conventional systems in the region (Agreste⁴⁵). Achievable yields for the L-GHG and the PHEP systems were considered to be 13% and 21% lower, respectively, than those of the current system in Ile-de-France. However, considering all the innovative systems together, it would appear to be possible to satisfy such ambitious constraints and environmental targets at the expense of only relatively small yield losses.

Available knowledge and current techniques suggested that it would not be possible to overcome all constraints in a single cropping system, because the agronomic practices used in one innovative system were incompatible with the constraints imposed on others. Plowing, one of the most effective practices against weeds used in the No-Pest system, is incompatible with large decrease in fossil energy consumption and the increase in C sequestration achieved with the L-EN and L-GHG systems, respectively. The large decrease in N fertilizer levels of the L-EN system is not compatible with the achievable yields defined for the PHEP and the L-GHG systems. Winter wheat sowing was delayed to avoid pest pressure in the No-Pest system, whereas it was brought forward in the direct drilling conditions of the L-EN and the L-GHG systems. This pattern was already evident during the design of the L-GHG system (*i.e.*, GHG emission processes were managed in hierarchical order). Consequently, the development of a system without pesticides, with ambitious constraints in terms of GHG emissions and fossil fuel use, and with other environmental and yield targets, will require further progress in agronomic knowledge. For example, a better understanding of the interactions between cash and cover crops in terms of cooperative and competitive effects might allow the introduction of a living cover crop during cash crop growth in the L-GHG system⁶². The field trial assessment again proves essential to gain a better understanding of these interactions.

4.3. Improving the design process

458 In the design of the L-GHG system, we had to rank the secondary objectives (C sequestration had to be enhanced first,
459 and then N₂O emissions had to be reduced) to satisfy the GHG constraint. In this case, several practices had effects on
460 both processes involved: no-tillage increased C sequestration in the soil but increased N₂O emissions; ample N fertilizer
461 applications were required to obtain high yields and, thus, abundant C residues, but this also generated more N₂O
462 emissions. We decided to promote C sequestration, because N₂O emission assessments were highly uncertain due to the
463 lack of published data about N₂O emissions, for field bean residues for example (IPPC³⁹), and the variability of results
464 due to differences in soil and climatic conditions⁶³. Nevertheless, knowledge about the effects of cropping systems on
465 N₂O emissions is increasing, and it should be possible to improve the adjustment of cropping systems in the future.

466 The cropping systems assessment required tools and models adapted to the set of objectives and convenient to use
467 during the iterative optimization process. Some approximations were used, due to the lack of data. In the L-GHG
468 system, the coefficient defined by the Intergovernmental Panel on Climate Change (*Tier 1*) was used to calculate N₂O
469 release from the amount of N applied, rather than using different values for different soil and climatic conditions⁶³.

470 There were also uncertainties in the assessment of C sequestration. The two simulations provided similar ranks for the
471 PHEP and the L-GHG systems, but the SIMEOS® tool and the Roth C model gave different quantitative results. One
472 difference between these tools relates to tillage. In the SIMEOS® tool, mineralization mechanisms in the tilled layer are
473 modified by tillage³⁶. In the Roth C model, tillage is taken into account only indirectly, through the C inputs to the soil
474 from crop biomass, which are affected by tillage practices⁴³. Further uncertainties arise from the lack of root biomass
475 data. Furthermore, these tools partially take into account changes in soil organic matter content, which greatly influence
476 N transformation mechanisms in the soil. Anyway, the difference in the values generated by these models suggested
477 that neither may accurately predict actual performances of these contrasting cropping systems. If cropping systems
478 assessment is to be considered relevant, it must make use of tools that are regularly updated, including: (i) “new”
479 machinery, such as harrows, hoes or direct-sowing drills with adjustable parameters in terms of energy consumption
480 and GHG emissions, and (ii) new more forms of pesticides or fertilizers. It may also be useful to improve systems
481 assessments with crop models, but many parameters are unavailable for marginal crops (such as hemp or flax), for crop
482 mixtures (*e.g.*, cereal and legume combinations), or species mixtures used as catch crops (*e.g.*, the association of spring
483 oats, mustard and clover).

484 Several economic and social aspects were not taken into account during this design process, which focused on the
485 cropping system rather than the farm scale. Nevertheless, we excluded some crops that are not grown by farmers in Ile-
486 de-France due to the lack of a market (*i.e.*, with low economic performances), but we included others for which the
487 market is poorly developed in this area (*e.g.*, lucerne and hemp). We did not take the organization of farm work into
488 account either, although the identification and quantification of pest pressures in the field, to decrease pesticide use, is
489 known to be time-consuming. From a technical point of view, we assumed that all farms owned the specific machinery

490 used in the innovative systems, such as direct-sowing drills or mechanical weeding tools. For all these reasons, further
491 investigations of the most relevant innovative cropping systems identified in a field trial assessment are needed, to
492 forecast their possible application in an area.

493 494 **4.4. Toward field assessment of the innovative cropping system prototypes**

495 The innovative systems prototypes designed here were based on different hypotheses about soil and climate effects,
496 because the true impact on crops and soil are unknown. For example we assumed that agricultural practices and climatic
497 conditions would not affect crop emergence. In our conditions, crops are not affected by direct drilling because this
498 technique is widely used in France (<http://agriculture-de-conservation.com/>). However, conservation tillage and mulch
499 tillage practices remain largely empiric⁶⁴, and additional knowledge is required for the optimization of these techniques,
500 to make it possible to achieve good results in terms of sowing management and target yields. We also assumed that
501 rainfall would provide enough water throughout the year in northern France to allow catch crop and main crop
502 emergence, but this may no longer be the case if the climate changes radically. We assumed that new equilibria would
503 appear with certain practices, allowing inputs to be reduced. In the L-GHG system, the practice of leaving crop residues
504 on the soil should decrease weed emergence, making it possible to decrease the amount of herbicide used, and the
505 absence of plowing should lead to the maintenance or increase in size of ground beetle populations, making
506 molluscicide use unnecessary. The achievable yield was determined on the basis of several assumptions. In the L-GHG
507 system, we promoted C sequestration, which is highly dependent on cereal yields and difficult to estimate in the case of
508 no-tillage systems, because the presence of excessive amounts of crop residues may decrease emergence. The chemical
509 properties of the soil associated with different tillage practices are also poorly characterized⁶⁴. In this system, the effects
510 on water availability of sowing cover crops every year also should be analyzed. In the L-EN system, the mixture of
511 winter wheat and white clover might also decrease cereal yields. In the No-Pest system, yields were defined by
512 approximation to organic systems, in which no mineral fertilizers are permitted. In this system, late winter cereal
513 sowing might lead to higher levels of damping-off and plant death during winter, decreasing yields to a greater extent
514 than anticipated. These are just a few of the uncertainties remaining about the real performance of the systems we have
515 devised. For these reasons, the field assessment of these system prototypes, which is currently being carried out (the
516 experiment started in 2008), is absolutely necessary.

517 Finally, it should be stressed that approaches of this type could be used in many agronomic situations, with a diversity
518 of challenges, provided that sufficient knowledge is available for the development of innovative strategies. In our
519 opinion, this approach meets the need expressed by Foley¹⁴ to “search for practical solutions” for a cultivated planet.

520

521 **Acknowledgments:**

522 We thank the experts who took part in the design process, Marie-Hélène Jeuffroy, Jane Jonhson, Laurence Guichard,
523 Marie Gosme and David Makowski for their useful comments on a previous version of the paper, Suzette Tanis-Plant
524 for linguistic help, and Alain Bone for assistance with bibliographic research.

525

526 **References**

- 527 1. IFEN, 2007. Les pesticides dans les eaux – données 2005. Les dossiers – n°9, ISSN: 1776-8411, 35 p.
528 <[http://www.ifen.fr/publications/le-catalogue-despublications/les-dossiers/2007/les-pesticides-dans-les-eaux-donnees-](http://www.ifen.fr/publications/le-catalogue-despublications/les-dossiers/2007/les-pesticides-dans-les-eaux-donnees-2005.html)
529 [2005.html](http://www.ifen.fr/publications/le-catalogue-despublications/les-dossiers/2007/les-pesticides-dans-les-eaux-donnees-2005.html)>.
- 530 2. Schrack D., Coquil X., Ortat A., and Benoît M., 2009. Rémanence des pesticides dans les eaux issues de parcelles
531 agricoles récemment converties à l'Agriculture Biologique. Innovations Agronomiques 4, 259-268
- 532 3. Höök M., Sivertsson A., and Aleklett K., 2010. Validity of the fossil fuel production outlooks in the IPCC Emission
533 Scenarios. Natural Resources Research: Volume 19, Issue 2, 63-81
- 534 4. Jakobsson K., Söderbergh B., Höök M. and Aleklett K., 2009. How reasonable are oil production scenarios from
535 public agencies? Energy Policy, Vol. 37, 4809-4818.
- 536 5. Shafiee S and Topal E., 2009. When will fossil fuel reserves be diminished? Energy Policy, vol. 37, issue 1, pages
537 181-189
- 538 6. Mohr S.H., Evans G.M., 2010. Long term prediction of unconventional oil production. Energy Policy, Volume 38,
539 Issue 1, 265-276
- 540 7. Smith P., Martino D., Cai Z., Gwary D., Janzen H., Kumar P., McCarl B., Ogle S., O'Mara F., Rice, C., Scholes B.,
541 and Sirotenko O., 2007. Agriculture. In Climate Change 2007: Mitigation. Contribution of Working Group III to the
542 Fourth Assessment Report of the Intergovernmental Panel on Climate Change, Metz, B. and Davidson, O. R. and
543 Bosch, P. R. and Dave, R. and Meyer, L. A.[eds.]. Cambridge University Press, Cambridge, United Kingdom and New
544 York, NY, USA.
- 545 8. Galloway J. N., Townsend A. R., Erisman J. W., Bekunda M., Cai Z. C., Freney J. R., Martinelli L. A., Seitzinger
546 S.P., Sutton M.A., 2008. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. Science,
547 16;320(5878):889-92. doi: 10.1126/science.1136674.
- 548 9. Arrouays D., Balesdent J., Germon J.C., Jayet P.A., Soussana J.F., and Stengel P., 2002. Expertise scientifique
549 collective "Contribution à la lutte contre l'effet de serre" Stocker du carbone dans les sols agricoles de France ?
550 Collective Scientific Expert Report. INRA edition, pp 332

551 10. Spiertz J.H.J., 2010. Nitrogen, sustainable agriculture and food security. A review *Agronomic Sustainable*
552 *Development*. 30 (2010) 43–55

553 11. Matson P.A., Parton W.J, Power A.G., and Swift M.J., 1997. Agriculture intensification and ecosystems properties,
554 *Science* 277, 504-509

555 12. UNEP 2007 http://www.unep.org/PDF/AnnualReport/2007/AnnualReport2007_en_web.pdf

556 13. Bouman B.A.M., Humphreys E., Tuong T.P., and Barker R., 2007. Rice and Water, *Advances in Agronomy*, 92,
557 187-237

558 14. Foley J.A., Ramankutty N., Brauman K.A., Cassidy E.S., Gerber J.S., Johnston M., Mueller N.D., O’Connell C.,
559 Ray D.K., West P.C., Balzer C., Bennett L.M., Carpenter S.R., Hill J., Monfreda C., Polasky S., Rockström J., Sheehan
560 J., and Siebert S., 2011. Solutions for a cultivated planet. *Nature*, 10452

561 15. Aubertot J.N., Barbier J.M., Carpentier A., Gril J.J., Guichard L., Lucas P., Savary S., Savini I., and Voltz M., 2005.
562 Pesticides, agriculture and the environment: reducing the use of pesticides and limiting their environmental impact.
563 Collective Scientific Expert Report. INRA – CEMAGREF. INRA edition, pp 63

564 16. Köpke U. and Nemecek T., 2010. Ecological services of faba bean. *Field Crops Research* 115 (2010) 217–233

565 17. Beheydt D., Boeckx P., Ahmed H. P., and Van Cleemput O., 2008. N₂O emissions from conventional and
566 minimum-tilled soils. *Biology and Fertility of Soils* 44, 863–873

567 18. Van Kessel C., Ventera R., Six J, Arlene M., Borde A., Linqvist B., and Van Groenigen K., 2012. Climate, duration,
568 and N placement determine N₂O emissions in reduced tillage systems: a meta-analysis. *Global Change Biology*, doi:
569 10.1111/j.1365-2486.2012.02779.x

570 19. Fu WeiJun, Tunney H., and Zhang Chao Sheng, 2010. Spatial variation of soil test phosphorus in a long-term
571 grazed, experimental grassland field. *Journal of Plant Nutrition and Soil Science*, 173, 3, 323-331.

572 20. Heinze S., Raupp J., and Joergensen R. G., 2010. Effects of fertilizer and spatial heterogeneity in soil pH on
573 microbial, biomass indices in a long-term field trial of organic agriculture. *Plant and Soil*, 328, 1/2, 203-215

574 21. Chirinda N., Olesen J.E., Porter J.R., and Schjønnig P., 2010. Soil properties, crop production and greenhouse gas
575 emissions from organic and inorganic fertilizer-based arable cropping systems. *Agriculture, Ecosystems &*
576 *Environment*, Volume 139, No. 4, 584-594

577 22. Tuomisto H.L., Hodge I.D., Riordan P., and Macdonald D.W., 2012. Comparing energy balances, greenhouse gas
578 balances and biodiversity impacts of contrasting farming systems with alternative land uses. *Agricultural Systems* 108 :
579 42–49

580 23. Shadbolt N., Kelly T., Horne D., Harrington K., Kemp P., Palmer A., and Thatcher A., 2009. Comparisons between
581 organic & conventional pastoral dairy farming, systems: cost of production and profitability. *Journal of Farm*
582 *Management*, 13, 10, 671-685

583 24. De Ponti T., Rijk B., and Van Ittersum M., 2012. The crop yield gap between organic and conventional agriculture.
584 Agricultural Systems 108: 1–9

585 25. Simonsen J., Posner J., Rosemeyer M., and Baldock J., 2010. Endogeic and anecic earthworm abundance in six
586 Midwestern cropping systems. Applied Soil Ecology 44. 147–155

587 26. Riley H., Pommeresche R., Eltun R., Hansen S., and Korsæth A., 2008. Soil structure, organic matter and
588 earthworm activity in a comparison, of cropping systems with contrasting tillage, rotations, fertilizer, levels and manure
589 use., Agriculture, Ecosystems Environment, 124, 3/4, 275-284

590 27. Snapp S.S., Gentry L.W., and Harwood R., 2010. Management intensity – not biodiversity – the driver of ecosystem
591 services in a long-term row crop experiment. Agriculture, Ecosystems & Environment 138 : 242–248

592 28. Nemecek T., Dubois D., Huguenin-Elie O., and Gaillard G., 2011. Life cycle assessment of Swiss farming systems:
593 I. Integrated and organic farming. Agricultural Systems 104 (2011) 217–232

594 29. Vereijken P., 1997. A methodological way of prototyping integrated and ecological arable farming systems
595 (I/EAFS) in interaction with pilot farms. European Journal of Agronomy, 7: 235-250

596 30. Dogliotti S., Van Ittersum M.K., and Rossing W.A.H., 2005. A method for exploring sustainable development
597 options at farm scale: a case study for vegetable farms in South Uruguay. Agricultural Systems 86: 29–51

598 31. Loyce C. and Wery J., 2006. Les outils des agronomes pour l'évaluation et la conception de systèmes de culture.
599 L'agronomie aujourd'hui, QUAE éditions, 77-95.

600 32. Lançon J., Wery J., Rapidel B., Angokaye M., Géraudeau E., Gaborelc C., Ballo D., and Fadegnon B., 2007. An
601 improved methodology for integrated crop management systems. Agronomy of Sustainable Development, 27: 101–110

602 33. Bockstaller C., Guichard L., Makowski D., Aveline A., Girardin P., and Plantureux S., 2008. Agri-environmental
603 indicators to assess cropping and farming systems. A review. Agronomy of Sustainable Development, 28: 139-149

604 34. Debaeke P., Munier-Jolain N., Bertrand M., Guichard L., Nolot J.M., Faloya V., and Saulas P., 2009. Iterative
605 design and evaluation of rule-based cropping systems: methodology and case studies. Agronomy of Sustainable
606 Development, 29: 73–86

607 35. Coleman K. and Jenkinson D.S., 1999. A model for the turnover of carbon in soil; Model description and windows
608 users guide, IACR – Rothamsted Harpenden Herts AL5 2JQ, p47

609 36. Andriulo A., Mary B., and Guerif J. 1999. Modelling soil carbon dynamic with various cropping sequences on the
610 rolling pampas. Agronomie, 19: 365-379

611 37. Van Groenigen K.J. , Hastings A., Forristal D., Jones M., and Smith P., 2011. Soil C storage as affected by tillage
612 and straw management: An assessment using field measurements and model predictions. Agriculture, Ecosystems &
613 Environment, 140: 218-225

614 38. GES'TIM (2010): Ministère de l'agriculture, de l'alimentation, de la pêche, de la ruralité et de l'aménagement du
615 territoire. Réf. 0933103, 156p

616 39. Intergovernmental Panel on Climate Change. Climate Change 2007. The physical science basis. UK: Cambridge
617 University Press, 1009p

618 40. Leteinturier B., Herman J.L., de Longueville F., Quintin L., and Oger R., 2006. Adaptation of a crop sequence
619 indicator based on a land parcel management system. *Agriculture, Ecosystems and Environment* 112 (2006) 324–334

620 41. Smith P., Smith J., Powlson D., McGill W., Arah J., Chertov O., Coleman K., Frnko U., Frolking S., Jenkinson D.,
621 Jensen L., Kelly R., Klein-Gunnewiek H., Komarov A., Li C., Molina J., Mueller T., Parton W., Thornley J., and
622 Whitmore A., 1997. A comparison of the performance of nine soil organic matter models using datasets from seven
623 long-term experiments. *Geoderma* 81, 153–225.

624 42. Bockstaller C. Guichard L., Keichinger O., Girardin P., Galan MB., Gaillard G., 2009. Comparison of methods to
625 assess the sustainability of agricultural systems. A review. *Agronomy of Sustainable Development*. 29: 223–235

626 43. De Li Liu, Yin Chan K., Conyers M.K., Li G., and Poile G.J., 2011. Simulation of soil organic carbon dynamics
627 under different pasture managements using the RothC carbon model. *Geoderma*, 165: 69–77

628 44. Cerri C., Easter M., Paustian K., Killian K., Coleman K., Bernoux M., Falloon P., Powlson D., Batjes N., Milne E.,
629 and Cerri C.C., 2007. Simulating SOC changes in 11 land use change chronosequences from the Brazilian Amazon with
630 RothC and Century models. *Agriculture, Ecosystems & Environment*, 122 (1), 46–57

631 45. Agreste, an agricultural statistical service of the French government <http://agreste.agriculture.gouv.fr/>

632 46. Riley H., Hoel B.O., Kristoffersen A.Ø., 2012. Economic and environmental optimization of nitrogen fertilizer
633 recommendations for cereals in Norway. *Acta Agriculturae Scandinavica, Section B, Soil & Plant Science*, 62, 5, 387-
634 400

635 47. Jeuffroy M.H., Baranger E., Carrouée B., de Chezelles E., Gosme M., Henault C., Schneider A., Cellier P., 2013.
636 Nitrous oxide emissions from crop rotations including wheat, oilseed rape and dry peas. *Biogeosciences*, 10, 1787–1797

637 48. Zentner R.P., Lafond G.P., Derksen D.A., Nagy C.N., Wall D.D., and May W.E., 2004. Effects of tillage method
638 and crop rotation on non-renewable energy use efficiency for a thin Black Chernozem in the Canadian Prairies. *Soil &*
639 *Tillage Research* 77: 125–136

640 49. Gelfand I., Snapp S., and Robertson G., 2010. Energy efficiency of conventional, organic, and alternative cropping
641 systems for food and fuel at a site in the US Midwest. *Environment Sciences Technology*, 44, 4006–4011

642 50. Nowacki W., 2008. Comparison of profitability of potato cultivation in organic and, integrated farming systems.
643 *Progress in Plant Protection*, 48, 4, 1526-1534

644 51. Chikowo R., Faloya V., Petit S., and Munier-Jolain N., 2009. Integrated weed management systems allow reduced
645 reliance on herbicides and long-term weed control. *Agriculture, Ecosystems & Environment*, 132: 237–242

646 52. Blazy J.M., Ozier-Lafontaine A., Doré T., Thomas A., and Wery J., 2009. A methodological framework that
647 accounts for farm diversity in the prototyping of crop management systems. Application to banana-based systems in
648 Guadeloupe. *Agricultural Systems* 101 . 30–41

649 53. Coulter J.A., Sheaffer C.C., Wyse D.L., Haar M.J., Porter P.M., Quiring S.R., and Klossner L.D., 2011. Agronomic
650 performance of cropping systems with contrasting crop rotations and external inputs. *Agronomy Journal*, 103: 1, 182

651 54. Moreno M., Lacasta C., Meco R., and Moreno C., 2011. Rainfed crop energy balance of different farming systems
652 and crop rotations in a semi-arid environment: Results of a long-term trial. *Soil & Tillage Research*, 114, 1: 18-27

653 55. Feiza V., Feiziene D., Auskalnis A., and Kadziene G., 2010. Sustainable tillage: results from long-term field
654 experiments on Cambisol., Zemdirbyste (*Agriculture*), 97, 2: 3-14

655 56. Baturu A., 2007. Effect of organic system on spring barley stem base health in, comparison with integrated and
656 conventional farming. *Journal of Plant Protection Research*, 47, 2: 167-178

657 57. Klimekova M. and Lehocka Z., 2009. Comparison of yields and qualitative characters of spring barley grown, after
658 three preceding crops in an organic farming system in the years, 2003-2008. *Agronomy Research*, 7, Special Issue, 335-
659 340

660 58. Liebman M., Gibson L. Sundberg D., Heggenstaller A., Westerman P., Chase C., Hartzler R., Menalled H., Davis
661 M., and Dixon P., 2008. Agronomic and economic performance characteristics of conventional and low-external-input
662 cropping systems in the Central Corn Belt. *Agronomy Journal*, 100, 3: 600-610

663 59. Cavigelli M. A., Hima B.L., Hanson J.C., Teasdale J.R., Conklin A.E., and Lu Y., 2009. Long-term economic
664 performance of organic and conventional field crops in the mid-Atlantic region. *Renewable Agriculture and Food
665 Systems*. 24, 2: 102-119

666 60. Le Gal P.Y., Dugué P., Faure G., and Novak S., 2011. How does research address the design of innovative
667 agricultural production systems at the farm level? A review *Agricultural Systems* 104 : 714–728

668 61. Stoorvogel J.J., Bouma J., and Orlich R.A., 2004. Participatory research for systems analysis: Prototyping for a
669 Costa Rican banana plantation. *Agronomy Journal*, 96, 2: 323-336

670 62. Shili-Touzi I., de Tourdonnet S., Nciri R., Le Floch D., Saulas P., and Doré T. (submitted). Is it possible to manage
671 competition between wheat and alfalfa grown as a service plant in an intercropping system in temperate conditions?
672 *Agriculture, Ecosystems & Environment*.

673 63. Rochette P. and Janzen H., 2005. Towards a revised coefficient for estimating N₂O emissions from legumes.
674 *Nutrient Cycling in Agroecosystems* 73, 171–179.

675 64. Alletto L., Coquet Y., and Roger-Estrade J., 2010. Two-dimensional spatial variation of soil physical properties in
676 two tillage systems. *Soil Use and Management*, 26, 432–444

677 65. Munier-Jolain N.M., Faloya V., Davaine J.B., Biju-Duval L., Meunier D., Martin C., and Charles R., 2004. A
678 cropping system experiment for testing the principles of integrated weed management: first results, in *Annales AFPP*,
679 XIIe colloque international sur la biologie des mauvaises herbes, Dijon, 147–156.

680 66. Tonitto C., David M. B., Li C. S., and Drinkwater L. E., 2007. Application of the DNDC model to tile-drained
681 Illinois agroecosystems: model comparison of conventional and diversified rotations. *Nutrient Cycling in*
682 *Agroecosystems* 78, 65–81.

683 67. Deike S., Pallutt B., Melander B., Strassemeier J., and Christen O., 2008. Long-term productivity and
684 environmental effects of arable farming as affected by crop rotation, soil tillage intensity and strategy of pesticide use:
685 A case-study of two long-term field experiments in Germany and Denmark. *European Journal of Agronomy*, 29: 191–
686 199

687 68. Nemecek T., von Richthofen J.F., Dubois G., Casta P., Charles R., and Pahl H., 2008. Environmental impacts of
688 introducing grain legumes into European crop rotations, *European Journal of Agronomy* 28: 380–393

689 70. XueLi Fu; Zhang Hui, and Jia JiZeng., 2009. Yield performance and resources use efficiency of winter wheat and
690 summer maize in double late-cropping system. *Acta Agronomica Sinica*: 35, 9: 1708-1714

691 71. Freibauer A., Rounsevell M. D. A., Smith P., and Verhagen J., 2004. Carbon sequestration in the agricultural soils
692 of Europe. *Geoderma* 122, 1–23.

693 72. Mosier A. R., Halvorson A. D., Peterson G. A., Robertson G. P., and Sherrod L., 2005. Measurement of net global
694 warming potential in three agroecosystems. *Nutrient Cycling in Agroecosystems* 72, 67–76.

695 73. Lehuger S., 2009. Modélisation des bilans de gaz à effet de serre des agro-systèmes en Europe. Thèse de doctorat de
696 l'Institut des Sciences et Industries du Vivant et de l'Environnement AgroParisTech. p184

697 74. Pelster D.E., Larouche F., Rochette P., Chantigny M.H., Allaire S., and Angers D.A., 2011. N fertilization but not
698 soil tillage affects nitrous oxide emissions from a clay loam soil under a maize–soybean rotation. *Soil and Tillage*
699 *Research*. 115-116: 16-26

701 **Tables**

702 Table 1: Examples of published agronomic strategies and associated practices for achieving specific goals

Specific goal	Agronomic management strategies	Example of agronomic practices
To reduce pesticide use: (Munier-Jolain ⁶⁵ ; Aubertot ¹⁵ ; Chikowo ⁵¹)	To avoid the coincidence of the pest, disease and weed contamination periods and sensitive stages of crop development	Modification of sowing date, earlier for oilseed rape, later for winter wheat
	To reduce the density of pests	Use of mechanical weeding Use of <i>Trichogramma</i> parasitoid wasps against <i>Ostrinia nubilalis</i> on maize
	To reduce the impacts of pests, diseases and weeds on crops	Decrease in sowing density and N fertilization to decrease shoot biomass Choice of varieties with the highest resistance Maximization of competition against weeds, by sowing winter oilseed rape very early (mid-August in the Ile-de-France region)
	To decrease the pool of pathogenic fungi in the soil, diseases and weeds	Lengthening of the crop rotation Sowing of a wide range of crops, to decrease the pool of fungi in the soil Maximization of the use of stale seed beds to increase weed emergence before sowing
	To maintain beneficial insects	Shallow plowing to maintain populations of carabid beetles (slug predators)
To reduce fossil fuel energy use: (Tonitto ⁶⁶ ; Deick ⁶⁷ ; Nemecked ⁶⁸ ; XueLi ⁷⁰ ; Gelfand ⁴⁹)	Direct energy: To decrease the use of the most power- consuming farm machinery Indirect energy: To decrease chemical N fertilization	Reducing or eliminating the use of deep plowing Introduction of large numbers of legumes and species with the highest N use efficiency into the crop rotation
To enhance C sequestration in the soil: (Arrouays ⁹ ; Freibauer ⁷¹ ; Mosier ⁷² ; Beheydt ¹⁷ ; Smith ⁷ ; Lehuger ⁷³)	To decrease organic matter mineralization To allow high levels of C accumulation, depending on the nature of the crop residues	Avoidance of plowing, replaced by direct drilling (conversion to no-tillage practices) Sowing of maize and cereals in rotation
To reduce N₂O emissions: (Rochette ⁶³ ; Pelster ⁷⁴)	To decrease anaerobic conditions, to decrease denitrification mechanisms To decrease the amount of available mineral N in the soil	Decreasing compaction and soil moisture content without deep tillage Optimization of N fertilization, according to soil nitrogen content Increasing the N use efficiency of crops
To decrease nitrogen leaching	To decrease soil nitrate content during autumn and winter	Sowing of catch crops Sowing of oilseed rape after legumes Banning of N fertilization during autumn and winter

703

704

705 Table 2: Crop rotations of the innovative cropping systems designed, and a current cropping system, defined by experts
 706 on the basis of data from the Ile-de-France region in 2006 (Agreste ⁴⁵)

Cropping system	Crop rotation
Productive high environmental performance (PHEP)	Winter field bean - Winter wheat - Winter oilseed rape - Winter wheat – (with mustard as a catch crop) - Spring barley
Without pesticide (No-Pest)	Triticale – (species mixture as a catch crop) Maize – Winter wheat – (species mixture as a catch crop) Spring field bean - Winter wheat - (species mixture as a catch crop) - Hemp
Low energy use: (L-EN)	Winter field bean - Winter wheat - Winter oil flax - Winter wheat-white clover mixture – (white clover as a catch crop) Spring oat
Low greenhouse gas emissions (L-GHG).	Triticale – (frost-sensitive species mixture as a catch crop) - Spring field bean - Winter oilseed rape (volunteers) - Winter wheat – (legumes as a cover crop) - Winter barley – (legume-oat mixture as catch crop) Maize
Ile-de-France (IdF)	Winter oilseed rape – Winter wheat – Winter barley – Maize – Winter wheat - Winter wheat

707

708

709 Table 3: Targeted yields (t ha⁻¹) for different species sown in the innovative cropping systems and comparison with
 710 current yields. Cropping systems: PHEP (productive high environmental performance), No-Pest (no pesticide use), L-
 711 EN (low energy use), L-GHG (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).

Cropping systems / Crops (t ha ⁻¹)	PHEP	No-Pest	L-EN	L-GHG	IdF Mean from 1998 to 2007 (Agreste)
Spring barley	6.2	-	-	-	
Winter barley	-	4.6	-	6.6	6.6
Spring field bean	-	4.7	-	4.7	
Winter field bean	3.4	-	3.4		
Hemp	-	8	-	-	
Maize	-	7.3	-	9.1	9.1
Winter oil flax	-	-	1.8	-	
Winter oilseed rape	3.1	-	-	3.1	3.4
Triticale	-	4.2	-	6.1	
Winter wheat	7.9	5.5	6.3	7.9	8.0

712

713

714 Table 4: Comparisons of total fossil energy consumption, expressed in MJ ha⁻¹, MJ t⁻¹ and kJ jCal⁻¹ (INDIGO® tool,
 715 v.1.9), between cropping systems. Cropping systems: PHEP (productive high environmental performance), L-EN (low
 716 energy use) and IdF (current system in Ile-de-France region).

	Total fossil energy consumption (MJ ha ⁻¹)	Total fossil energy consumption (MJ t ⁻¹)	Total fossil energy consumption (kJ kCal ⁻¹)
<u>L-EN - PHEP</u> PHEP	-49%	-24%	-29%
<u>PHEP - IdF</u> IdF	-31%	-21%	-30%
<u>L-EN - IdF</u> IdF	-65%	-40%	-50%

718 Table 5: C sequestration (t CO₂.eq ha⁻¹) simulated over 25-year and 50-year periods for the different cropping systems
 719 with the ROTH C model (V.26-3) and the SIMEOS® tool (2010), for mean soil organic matter content in the Ile-de-
 720 France region (1.6%. SOM = 123.3 t CO₂.eq ha⁻¹). Cropping systems: PHEP (productive high environmental
 721 performance), L-GHG (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).

C sequestration (t CO ₂ .eq ha ⁻¹)		Assessment period	Assessment period
		25 years	50 years
Cropping systems	Tool or model		
PHEP	SIMEOS® tool	15.1	19.7
PHEP	ROTH C model	85.2	116.0
L-GHG	SIMEOS® tool	32.0	40.5
L-GHG	ROTH C model	102.9	147.8
IdF	SIMEOS® tool	17.4	22.1
IdF	ROTH C model	87.1	129.2
<u>(L-GHG – PHEP)</u>			
PHEP SOC	SIMEOS® tool	12%	15%
<u>(L-GHG – PHEP)</u>			
PHEP SOC	ROTH C model	9%	13%

723 Table 6: Comparison of total simulated greenhouse gas emissions, expressed in t CO₂-eq ha⁻¹ and t CO₂-eqt⁻¹, between
 724 cropping systems. Cropping systems: PHEP (productive high environmental performance), L-GHG (low greenhouse
 725 gas emissions) and IdF (current system in Ile-de-France region).

	Total greenhouse gas emissions (t CO ₂ -eq ha ⁻¹)	Total greenhouse gas emissions (t CO ₂ -eq t ⁻¹)
<u>L-GHG - PHEP</u> PHEP	-13%	-22%
<u>PHEP - IdF</u> IdF	-46%	-32%
<u>L-GHG - IdF</u> IdF	-53%	-47%

727 Table 7: Greenhouse gas balances (t CO₂-eq ha⁻¹), for different cropping systems, determined with the SIMEOS® tool
 728 (2010) and the ROTH C model (V.26-3), for a current soil in the Ile-de-France region (SOM = 1.6%) and the
 729 GES'TIM³⁷ database, for 25-year and 50-year periods. The GHG balance is positive when the amount of carbon emitted
 730 in greenhouse gases exceeds that sequestered. Cropping systems: PHEP (productive high environmental performance),
 731 L-GHG (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).

Greenhouse gas balances (t CO ₂ -eq ha ⁻¹),		Assessment period 25 years	Assessment period 50 years
Cropping system	Tool or model		
PHEP	SIMEOS® tool	16.7	44.0
PHEP	ROTH C model	-53.4	-52.3
L-GHG	SIMEOS® tool	-4.4	14.4
L-GHG	ROTH C model	-75.3	-92.6
IdF	SIMEOS® tool	41.9	96.4
IdF	ROTH C model	-39.7	-10.8
L-GHG – PHEP	SIMEOS® tool	-21.1	-29.3
L-GHG – PHEP	ROTH C model	-21.9	-40.3

733 Table 8: Greenhouse gas balance ratio of the various cropping systems, expressed in t CO₂-eq ha⁻¹ , simulated with the
734 ROTH C (V.26-3) model over two different periods (25 and 50 years). Cropping systems: PHEP (productive high
735 environmental performance), L-GHG (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).


	GHG balance: 25 years	GHG balance: 50 years
L-GHG / PHEP	1.51	1.76
PHEP /IdF	1.39	4.95
L-GHG / IdF	2.10	8.72

736


737

738 **Figures**

739 Figure 1: Cropping system prototypes assessment with GESTIM database: Mean total fossil energy consumption
740 (expressed in MJ ha⁻¹ year⁻¹) calculated over one rotation period for the different cropping systems, and total fossil
741 energy consumption for each crop of the various cropping systems (i.e.: indirect energy – d.e.: direct energy - w: winter
742 - s: spring). Cropping systems: PHEP (productive high environmental performance), L-EN (low energy use) and IdF
743 (current system in Ile-de-France region).



745 Figure 2: Cropping system prototypes assessment with GESTIM database: Mean total greenhouse gas emissions
 746 (expressed in kg CO₂-eq ha⁻¹ year⁻¹) calculated over one rotation period for the different cropping systems, and total
 747 greenhouse gas emissions for each crop of the various cropping systems (i.ghg: indirect greenhouse gas – d.ghg: direct
 748 greenhouse gas- w: winter - s: spring). Cropping systems: PHEP (productive high environmental performance), L-GHG
 749 (low greenhouse gas emissions) and IdF (current system in Ile-de-France region).



751 Figure 3: Environmental criteria assessment of the different cropping system prototypes with the INDIGO® tool
 752 (v.1.9). Cropping systems: PHEP (productive high environmental performance), L-GHG (low greenhouse gas
 753 emissions), L-EN (low energy), No-Pest (No pesticides) and IdF (current system in Ile-de-France region). Minimum
 754 values to satisfy environmental targets.

