

Origins of the performance gaps in innovative cropping systems under experimental assessment.

Caroline C. Colnenne-David, Gilles Grandeau, Véronique Tanneau, Marie-Helene M.-H. Jeuffroy, Thierry Doré

► To cite this version:

Caroline C. Colnenne-David, Gilles Grandeau, Véronique Tanneau, Marie-Helene M.-H. Jeuffroy, Thierry Doré. Origins of the performance gaps in innovative cropping systems under experimental assessment.. The 5th International Symposium for Farming Systems Design "Multi-functional farming systems in a changing world", Sep 2015, Montpellier, France. hal-01357467

HAL Id: hal-01357467 https://agroparistech.hal.science/hal-01357467v1

Submitted on 30 Aug2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Origins of the performance gaps in the innovative cropping systems under experimental assessment

Colnenne-David C.^{1,2}, Grandeau G.^{1,2}, Tanneau V.^{1,2}, Jeuffroy M.H.^{1,2}, Doré T.^{2,1}

¹ INRA, UMR 211, 78850 Thiverval-Grignon, France (caroline.colnenne@grignon.inra.fr), ² AgroParisTech, UMR 211, 78850 Thiverval-Grignon, France

Context and Objectives

To meet new agricultural issues and make agriculture more sustainable, innovative cropping systems (ICSs) targeting a multiplicity of purposes need to be designed. Four ICSs were designed by prototyping (Reau and Doré, 2008¹) and assessed in a long-term field system experiment.

Our objective was to analyze the results after the first complete rotation, particularly the bad performances compared to the targeted aims.

Materials and Methods

Multiple goals of the four designed ICSs (Colnenne-David and Doré, 2014²)

The PHEP ICS goals:

- 1. To satisfy multiple environmental criteria:
- *low pesticide use → high crop diversity, highly resistant varieties
- *low direct energy consumption \rightarrow only 1 ploughing within the
- rotation *low indirect energy **consumption** \rightarrow legumes in
 - the rotation
- *low nitrogen leaching → catch crop (CC) before spring crops and no N fertilization during autumn and winter
- *stabilization and/or to enrich **soil organic matter** → burying residues of all crops

2. To reach yield targets matching the Ile-de-France yields **Crop sequence**: winter faba

agrosystem occurred

• • • • • • • • • •

The L-GHG ICS goals:

1. <u>50% GHG emissions compared to the PHEP ICS</u>

- (i) increase soil C sequestration \rightarrow many cereals, continuous soil cover, high yield targets, no ploughing
- (ii) decrease N₂O emissions \rightarrow high number of legume crops in the rotation, improvement of N fertilization management, crops with taproots in order to reduce soil compaction
- 2. To satisfy multiple environmental criteria: idem PHEP ICS
- 3. To reach yield targets matching the Ile-de-France yields

Crop sequence: catch crop (CC), maize, triticale, CC, spring faba bean, winter oilseed rape, winter wheat, CC, winter barley

The L-EN ICS goals:

1. <u>50% fossil energy consumption</u> compared to the PHEP ICS

- (i) Low direct energy consumption \rightarrow no ploughing and using direct sowing machine (ii) Low indirect energy consumption → high number of legume species in the rotation, species with high N efficiency use, decrease N fertilization by reducing yield objectives
- 2. To satisfy multiple environmental criteria: idem PHEP ICS
- **3.** To reach yield targets: 20% lower than the lle-de-France yields
- **Crop sequence**: winter faba bean, winter wheat, winter flax, winter wheat-*trifolium* mixture, *Trifolium* as CC, spring oat


The No-Pest ICS goals:

- **1. No pesticide is allowed**
- \rightarrow long rotation including a wide diversity of species (e.g. hemp), alternate sowing dates, different

Main characteristics of the field

trial

✓ located in Grignon (78, France) \checkmark 6.2 ha (surface plot: ≈ 4000 m²) ✓ 3 blocks ✓ deep loamy soil ✓ Beginning of field assessment: 2008



bean, winter wheat, winter oilseed rape, winter wheat, mustard as CC and spring barley 3. To	s and densities of sowing, highly resistant varieties or mixtures, ploughing and mechanical weeding satisfy multiple environmental criteria: idem PHEP ICS reach yield targets: higher than organic systems in the IIe-de-France sequence: triticale, CC, maize, winter wheat, CC, spring faba bean, winter wheat, CC, hemp
	Results: Classification of the major disparities
Classification	Examples collected in the ICSs
Some agronomical strategies w no suitable to reach the goa Some practices were not adapted to satisfy a multiplic objectives	Is C sequestration evolution = -149kgCO ₂ ha ⁻¹ year ⁻¹ (+87kgCO ₂ ha ⁻¹ year ⁻¹ expected) In the No-Pest ICS: No possible to satisfy both the pesticide constraint and the S.O.M. criteria.
Some practices were not appropriate in the context of field-trial conditions	In the L-GHG ICS: Very dry conditions in summer 3 years / 6 → Low amount of aerial biomass of cover crops
An unpredicted evolution of t	he In both the L-EN and the L-GHG ICSs:

High weed development \rightarrow to mow oilseed rape plots in 2014

Discussions – Conclusion

- > After the first complete rotation the major sources of disparities were classified
- > Nevertheless, a more complete agronomic diagnosis is necessary to identify and to rank all the causes of **bad performances**
- > This knowledge allowed us to improve the innovative cropping systems through a new design step > This experiment contributes to the learning design processes and cropping system management

REFERENCES

¹Reau R. and Doré T. (2008). "Systèmes de culture innovants et durables : quelles méthodes pour les mettre au point et les évaluer ? "Educagri Editions, Dijon, France, pp 175 ² Colnenne-David C. and Doré T., 2014. Designing innovative productive cropping systems with quantified and ambitious environmental goals. "Renewable Agriculture and Food Systems". doi:10.1017/S1742170514000313.

Farming System Design, Conference 2015 (September) in Montpellier (France)