

# CO2 and O2 solubility and diffusivity data in food products stored in data warehouse structured by ontology

Valérie V. Guillard, Patrice Buche, Juliette Dibie-Barthelemy, Stéphane Dervaux, Filippo Acerbi, Estelle Chaix, Nathalie N. Gontard, Carole Guillaume

# ▶ To cite this version:

Valérie V. Guillard, Patrice Buche, Juliette Dibie-Barthelemy, Stéphane Dervaux, Filippo Acerbi, et al.. CO2 and O2 solubility and diffusivity data in food products stored in data warehouse structured by ontology. Data in Brief, 2016, 7, pp.1556-1559. 10.1016/j.dib.2016.04.044. hal-01357730

# HAL Id: hal-01357730 https://agroparistech.hal.science/hal-01357730

Submitted on 27 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Contents lists available at ScienceDirect

# Data in Brief





## Data Article

# CO<sub>2</sub> and O<sub>2</sub> solubility and diffusivity data in food products stored in data warehouse structured by ontology



Valérie Guillard <sup>a,\*</sup>, Patrice Buche <sup>a</sup>, Juliette Dibie <sup>b</sup>, Stéphane Dervaux <sup>b</sup>, Filippo Acerbi <sup>a</sup>, Estelle Chaix <sup>a</sup>, Nathalie Gontard <sup>a</sup>, Carole Guillaume <sup>a</sup>

#### ARTICLE INFO

Article history: Received 19 March 2016 Received in revised form 15 April 2016 Accepted 19 April 2016 Available online 26 April 2016

Keywords: Diffusivity Solubility Data Data warehouse Ontology Food  $O_2$  $CO_2$ 

#### ABSTRACT

This data article contains values of oxygen and carbon dioxide solubility and diffusivity measured in various model and real food products. These data are stored in a public repository structured by ontology. These data can be retrieved through the @Web tool, a user-friendly interface to capitalise and guery data. The @Web tool is accessible online at http://pfl.grignon.inra.fr/atWeb/.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license

(http://creativecommons.org/licenses/by/4.0/).

# Specifications Table

Subject area Biochemistry

More specific Food science and food engineering subject area

Type of data Table, links

E-mail address: guillard@univ-montp2.fr (V. Guillard).

<sup>&</sup>lt;sup>a</sup> UMR IATE, University of Montpellier – INRA, 2 place Pierre Viala, F-34060 Montpellier Cedex 1. France

b AgroParisTech & INRA UMR MIA 518, 16, rue Claude Bernard, F-75 231 Paris Cedex 05, France

<sup>\*</sup> Corresponding author.

How data was	Chemical titration (for CO <sub>2</sub> quantification) and luminescence-based detection (for
acquired	$O_2$ detection) implemented in dedicated experimental set-ups
Data format	Analyzed, ready to use
Experimental	Samples considered are model and real food products without any pre-treatment
factors	except addition of sodium azide to avoid microbial growth
Experimental	Solubility is measured by quantifying the concentration of dissolved gas in a
features	sample in equilibrium with a fix and controlled partial pressure.
	Diffusivity is identified from an experimental diffusion kinetic curve by using a
	mathematical model and appropriate numerical treatment (algorithm of
	optimization).
Data source	University of Montpellier, FR-34060, France
location	
Data accessibility	Data is within this article.

# Value of the data

- A unique set of CO<sub>2</sub> solubility and diffusivity data indispensable in food engineering to model CO<sub>2</sub> gas transfer in food.
- A unique set of O<sub>2</sub> diffusivity values within synthetic oils as a function of temperature.
- O<sub>2</sub> diffusivity data could be used to predict oxidation of O<sub>2</sub>-sensitive compounds in foods.
- These data could serve as benchmark for other researchers coping with research on gas transfer in food for numerous simulation.

#### 1. Data

Data shared with this article are more than 100 data of solubility and diffusivity of gases ( $O_2$  and  $CO_2$ ) in food samples. These data are stored in a data warehouse called @Web in which the data management is guided by ontology.

All data are available for uploading at the URL specified below and recalled in the table hereafter with the details about the nature and amount of data available at each URL.

Data type	Table URL (copy/paste the URL in your Internet browser)	Amount of data
$CO_2$	pfl.grignon.inra.fr/atWeb/TableServlet?	34
solubility	<u>viewTable = 2775&amp;idDoc = 1335&amp;id = 35272672</u>	
•	pfl.grignon.inra.fr/atWeb/TableServlet?	21
	viewTable = 2776&idDoc = 1335&id = 35305550	
	pfl.grignon.inra.fr/atWeb/TableServlet?	48
	viewTable = 2773&idDoc = 1335&id = 35245144	
	pfl.grignon.inra.fr/atWeb/TableServlet?	3
	viewTable = 2732&idDoc = 1332&id = 34354344	
CO <sub>2</sub> diffu-	pfl.grignon.inra.fr/atWeb/TableServlet?	12
sivity	<u>viewTable=2780&amp;idDoc=1346&amp;id=35361064</u>	
-	pfl.grignon.inra.fr/atWeb/TableServlet?	11
	<u>viewTable = 2826&amp; idDoc = 1346&amp; id = 36350532</u>	
	pfl.grignon.inra.fr/atWeb/TableServlet?	11
	<u>viewTable=2779&amp;idDoc=1346&amp;id=35346176</u>	
	pfl.grignon.inra.fr/atWeb/TableServlet?	12
	<u>viewTable = 2778&amp;idDoc = 1346&amp;id = 35333312</u>	
	pfl.grignon.inra.fr/atWeb/TableServlet?	16
	<u>viewTable</u> =2777&idDoc=1346&id=35320430	

	pfl.grignon.inra.fr/atWeb/TableServlet?	11
	<u>viewTable</u> =2733&idDoc=1332&id=34367074	
O <sub>2</sub> diffusiv-	pfl.grignon.inra.fr/atWeb/TableServlet?	24
ity	<u>viewTable</u> = <u>2764</u> & <u>idDoc</u> = <u>1342</u> & <u>id</u> = <u>35103704</u>	
	pfl.grignon.inra.fr/atWeb/TableServlet?	30
	<u>viewTable</u> = <u>2755</u> & <u>idDoc</u> = <u>1342</u> & <u>id</u> = <u>34742706</u>	
	pfl.grignon.inra.fr/atWeb/TableServlet?	10
	<u>viewTable</u> = <u>2754</u> & <u>idDoc</u> = <u>1342</u> & <u>id</u> = <u>34730102</u>	
	pfl.grignon.inra.fr/atWeb/TableServlet?	44
	<u>viewTable</u> = <u>2765</u> & <u>idDoc</u> = <u>1342</u> & <u>id</u> = <u>35116534</u>	
	pfl.grignon.inra.fr/atWeb/TableServlet?	2
	<u>viewTable</u> = <u>2910</u> & <u>idDoc</u> = <u>1342</u> & <u>id</u> = <u>40225446</u>	

# 2. Experimental design, materials and methods

**O**<sub>2</sub>. Oxygen optical sensors (Presens GmbH, Regensberg, Germany) were used to monitor O<sub>2</sub> partial pressure. This measurement is based on dynamic luminescence quenching. Due to an excitation flash emitted through an optical fibre, the luminophore contained in the sensor goes into an excited state and thus emits fluorescence backscatter signal, which is detected by the optical fibre. If the luminophore is in contact with an oxygen molecule, the backscatter signal is changed due to a dynamic quenching of luminescence. The change in the backscatter signal permits to detect the O<sub>2</sub> partial pressure in the medium. Two different set-ups exist (1) an invasive O<sub>2</sub>-sensitive optical sensor made of a syringe probe (micro-sensors, Presens GmbH, Regensburg, Germany) connected to the optical fibre and oxygen metre (Oxy-4 micro, Presens) and (2) a non-invasive oxygen sensor made of a dot of 5 mm of diameter that can be stuck on the wall of a transparent container and measurement is then made through the transparent container.

Oxygen sorption kinetics were measured at fixed temperature value when imposing a controlled partial pressure of  $O_2$  in the surrounding of the sample. The mono-directional  $O_2$  ingress into the sample was measured locally at the bottom or in the middle of the thin layer of food material previously free of  $O_2$  using one of the aforementioned sensors. More details on the experimental set-up could be found in [1–3].

 $\mathbf{CO_2}$ . The solubility of  $\mathbf{CO_2}$  was measured at equilibrium by quantification of the gas dissolved in the sample using chemical titration [4,5]. This measurement was done in a set-up where the sample is in a controlled chamber (controlled temperature, relative humidity,  $\mathbf{CO_2}$  gas composition).

The diffusion of  $CO_2$  was characterised by (1) imposing a gradient of  $CO_2$  to a piece of material of simple geometry (cylinder or plane sheet), (2) measuring the  $CO_2$  sorption kinetic in the sample and (3) identifying diffusivity values by adjusting a dedicated mathematical model to the experimental kinetic. Two types of kinetic could be obtained: (1)  $CO_2$  space-dependent profile in the cylindrical sample after its slicing and  $CO_2$  quantification in each slice or (2)  $CO_2$  time-dependent profile after  $CO_2$  quantification in each thin slice (one slice corresponding at one time of kinetic) [4,6].

Numerical treatment. For both O<sub>2</sub> and CO<sub>2</sub>, diffusivities are identified by fitting a dedicated mathematical model to the experimental kinetic curve (space-dependent profile or time-dependent profile). This identification step is performed using a routine ("Isqnonlin") of Matlab<sup>®</sup> software.

# Acknowledgements

Part of the data presented here were acquired in the framework of the Map'Opt project (ANR-10-ALIA-002 2011 to 2015) funded by the French National Research Agency, whose title is "Equilibrium gas composition in modified atmosphere packaging and food quality".

# Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at http://dx.doi. org/10.1016/j.dib.2016.04.044.

## References

- [1] E. Chaix, C. Guillaume, N. Gontard, V. Guillard, Performance of a non-invasive methodology for assessing oxygen diffusion in liquid and solid food products, J. Food Eng. 171 (2016) 87–94. http://dx.doi.org/10.1016/j.jfoodeng.2015.10.014.
- [2] C. Pénicaud, S. Guilbert, S. Peyron, N. Gontard, V. Guillard, Oxygen transfer in foods using oxygen luminescence sensors: influence of oxygen partial pressure and food nature and composition, Food Chem. 123 (2010) 127–1281 (http://www.sciencedirect.com/science/article/B6T6R-5051PS2-2/2/923ab815d46f428626b86c9b60f335b5).
- [3] E. Chaix, C. Guillaume, V. Guillard, Oxygen and carbon dioxide solubility and diffusivity in solid food matrices: a review of past and current knowledge, Compr. Rev. Food Sci. Food Saf. 13 (2014) 261–286. http://dx.doi.org/10.1111/1541-4337.12058.
- [4] E. Chaix, C. Guillaume, N. Gontard, V. Guillard, Diffusivity and solubility of CO<sub>2</sub> in dense solid food products, J. Food Eng. 166 (2015) 1–9. http://dx.doi.org/10.1016/j.jfoodeng.2015.05.023.
- [5] F. Acerbi, V. Guillard, C. Guillaume, N. Gontard, Impact of selected composition and ripening conditions on CO<sub>2</sub> solubility in semi-hard cheese, Food Chem. 192 (2016) 805–812. http://dx.doi.org/10.1016/j.foodchem.2015.07.049.
- [6] F. Acerbi, V. Guillard, C. Guillaume, M. Saubanere, N. Gontard, An appraisal of the impact of compositional and ripening parameters on CO<sub>2</sub> diffusivity in semi-hard cheese, Food Chem. 194 (2016) 1172–1179. http://dx.doi.org/10.1016/j. foodchem.2015.08.020.