

## Ambitious environmental and economic goals for the future of agriculture are unequally achieved by innovative cropping systems

Caroline Colnenne-David, Gilles Grandeau, Marie-Hélène Jeuffroy, Thierry

Doré

### ▶ To cite this version:

Caroline Colnenne-David, Gilles Grandeau, Marie-Hélène Jeuffroy, Thierry Doré. Ambitious environmental and economic goals for the future of agriculture are unequally achieved by innovative cropping systems. Field Crops Research, 2017, 210, pp.114 - 128. 10.1016/j.fcr.2017.05.009 . hal-01540437

### HAL Id: hal-01540437 https://agroparistech.hal.science/hal-01540437

Submitted on 2 Aug 2019  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Ambitious environmental and economic goals for the future of agriculture are unequally achieved
 by innovative cropping systems

3 <u>https://dx.doi.org/10.1016/j.fcr.2017.05.009</u>

- 4 Colnenne-David C., Grandeau G., Jeuffroy M.H., Doré T.
- 5

#### 6 Abstract

7 Agriculture has to face huge challenges in the decades ahead. Four innovative cropping systems were assessed in 8 a "cropping system experiment" in the Ile-de-France region (France) from 2009 to 2014. Three were designed to 9 meet ambitious goals: the total elimination of pesticides (No-Pest), reducing fossil energy consumption by 50% 10 (L-EN), or decreasing greenhouse gas (GHG) emissions by 50% (L-GHG). They were also required to satisfy a 11 wide range of environmental criteria and to maximize yields whilst respecting the major constraint on the system 12 and the environmental targets set. A fourth system (PHEP), in which the environmental and yield targets were 13 achieved with no major constraint, was also assessed. After completion of the first full crop sequence for these 14 innovative systems, the results obtained indicated that it was possible to design and implement innovative systems 15 achieving multiple goals. In our field trial conditions, the pesticide and energy constraints were almost satisfied, 16 whereas the GHG target was missed by a considerable margin. All four innovative systems satisfied environmental 17 criteria in terms of N management, pesticide use, energy consumption and crop diversity. However, herbicide 18 treatment frequency index (TFIH) was higher than expected in the two systems with no-plow practices, L-EN and 19 L-GHG. In the pesticide-free system, soil organic matter content was lower than expected, due to frequent plowing 20 (every 2 years) and low residue levels as a result of the lower yields obtained. Yields were lower for the L-EN 21 system than for the reference system, and yield was variable in the L-GHG system. These innovative systems had 22 better environmental performances than the systems currently used in the Ile-de-France region, with no decrease 23 in gross margins.

- 24
- 25 Key words: cropping system experiment, field assessment, greenhouse gas emissions, pesticide, energy.
- 26
- 27

#### 28 <u>1. Introduction</u>

29 New challenges are continually arising in agriculture, necessitating profound breakthrough innovations in 30 agricultural practices. The most serious issues faced concern: (1) the loss of biodiversity in agroecosystems, (2) 31 the need to reduce chemical inputs, which are known to be harmful to the environment and human health, and (3) 32 the need to decrease the impact of agriculture on climate change, by decreasing greenhouse gas emissions and 33 promoting carbon storage in the soil. Current arable cropping systems are of questionable sustainability, and 34 alternative cropping systems must therefore be designed, to meet the goals of a more sustainable agriculture. 35 Agronomists design and assess innovative cropping systems to tackle a wide range of issues (Doré et al., 2011; 36 Blazy et al., 2009; Sadok et al., 2009). Moreover, given that global food security has become a primary concern 37 (Godfray et al., 2010), there is a need for innovative cropping systems that increase agricultural resource use 38 efficiency (Foley et al., 2011).

39

40 New strategies for crop management and new cropping systems have been designed in recent years. Many have 41 targeted a single principal goal, such as enhancing C sequestration through changes in crop management (e.g., 42 Freibauer et al., 2004; Dimissi et al., 2014), reducing pesticide use (Aubertot et al., 2005; Chikowo et al., 2009), 43 decreasing energy consumption (Singh et al., 2008; Khakbazan et al., 2009), or improving the yield of a single 44 crop (Tapia et al., 2014). However, some studies were "innovation-pushed": the authors compared cropping 45 systems on the basis of the combination of agricultural practices used (Kulak et al., 2015), rather than on the 46 achievement of target performances with the most appropriate practices. For example, they compared organic and 47 conventional systems (Panasiewiez et al., 2010; Nemecek et al., 2011a), or no-tillage and conventional tillage 48 systems (Abdi et al., 2014; Dimissi et al., 2014), without providing any further information about the objectives 49 to be reached. In most of these examples, only a few criteria were assessed in field trials: the distribution of 50 phosphorus species in the soil profile (Abdi et al., 2014), changes in soil structure and yield performances 51 (Abdollahi et al., 2015), soil biological properties (Ingle et al., 2014), ecophysiological characteristics of spring 52 barley and genotypes under various systems (Panasiewiez et al., 2010), and weed infestation under different long-53 term tillage systems (Chikowo et al., 2009). However, in some cases, multi-criteria analyses were performed, with 54 various methodologies (Nemecek et al., 2011a, 2011b; Loyce et al., 2012; Kulak et al., 2015). These multi-criteria 55 assessments made it possible to analyze combinations of agricultural practices with opposite impacts on specific 56 criteria, and to consider trade-offs. For example, no-till systems decrease energy consumption, but increase 57 herbicide use (Zentner et al., 2004).

58

59 To our knowledge, no study has yet both (i) designed in silico innovative and consistent cropping systems 60 addressing a multiplicity of current issues, and (ii) assessed them in a cropping system experiment involving the 61 analysis of multiple performances. We designed in silico innovative cropping systems addressing multiple issues 62 of importance (Colnenne-David and Doré, 2015a), and conducted system experiments to assess their ability to 63 achieve several goals. Four innovative cropping systems targeting various environmental goals and yield 64 objectives were designed by the prototyping method described by Vereijken (1997). Their performances were 65 assessed ex ante with various tools and models: the Indigo® method (www7.inra.fr/indigo) for environmental 66 performances, the Simeos® tool (using the AMG model, Andriulo et al., 1999) and the Roth C model for carbon 67 sequestration, as in the study by Colnenne-David and Doré, 2015a. For each combination of objectives, the most 68 promising candidate system was then implemented in a cropping system experiment.

69

We present here the cropping system experiment results for these four innovative cropping systems, for the first full crop sequence. We analyzed the performance of the cropping systems in several different ways: (1) we compared the innovative cropping systems implemented in the field trial with the prototypes (Colnenne-David and Doré, 2015a); (2) we compared the three innovative systems designed to meet particular constraints with a constraint-free innovative system used as the reference system and (3) we compared the innovative systems and the current system in the Ile-de-France region, where the field trial took place.

76

#### 77 2. Materials and methods

#### 78 **2.1.** General description of the four innovative cropping systems

79 Four innovative cropping systems with quantified constraints, and environmental and yield targets were designed 80 jointly with various stakeholders, including farmers, in 2008 (table 1, Colnenne-David and Doré, 2015a). The 81 "productive with high environmental performance" (PHEP) system was designed to minimize environmental 82 impact (decreasing nitrate and pesticide pollution, enhancing crop diversity or reducing fossil energy consumption 83 relative to current cropping systems) and to reach the maximum possible yield given the environmental targets, as 84 described by Colnenne-David and Doré, 2015a. This cropping system, which was designed without major 85 environmental constraints, was used as the reference system for comparisons with the other systems. Each of the 86 other three systems was designed to meet an additional environmental constraint, constituting a major 87 breakthrough in terms of the objectives for current cropping systems: the elimination of pesticide use (No-Pest), 88 reducing fossil energy consumption by 50% relative to the PHEP system (L-EN), or halving greenhouse gas 89 emissions relative to the PHEP system (L-GHG). These cropping systems were also designed to minimize 90 environmental impact whilst providing the maximum possible yield under the constraint imposed and respecting 91 the environmental targets. During the design step, the constraints and targets were prioritized as follows: the 92 environmental constraint had to be satisfied first, the set of other environmental targets then had to be attained, 93 and, finally, yield had to be maximized. The systems retained for field assessment corresponded to the combination 94 of agricultural practices resulting in the highest yields in silico among the candidate systems both satisfying 95 environmental constraints and meeting environmental targets.

96

#### 97 2.2. Main agronomic characteristics of the four innovative cropping systems

98 The four cropping systems were based on the agronomic strategies described in table 1 (Colnenne-David and Doré,
99 2015a).

100

#### 101 **2.3. Experimental trial**

102 Since 2008, the innovative cropping systems have been implemented in a cropping system experiment, located at 103 the AgroParisTech experimental farm at Grignon, in the Ile-de-France region (*i.e.* Paris Basin, N 48.84°, E 1.95°). 104 This site has a deep, homogeneous loamy clay soil (FAO, 1998). Mean annual rainfall, calculated over a 20-year 105 period was about 650 mm per year at this site. The crop immediately preceding this experiment was winter barley 106 and the field had been plowed (30 cm depth). The trial covered a total area of 6.2 ha, divided into large plots 107 (almost 4000 m<sup>2</sup>) to facilitate the rational use of farm machinery in conditions representative of those on farms. 108 Due to both the limited area available for the trial and the need for large plots, each system was randomly 109 distributed in a block design with only three replicates. The size of the trial was such that we were unable to grow 110 all of the crops of each crop sequence in each innovative system each year. The interannual variability results were 111 taken into account by sowing three different crops from the crop sequence of each system in the three replicates 112 for the year concerned, for each of the innovative systems (e.g. in 2009, winter wheat, winter oilseed rape and 113 spring barley were sown in the three different replicates of the PHEP system). The first full crop sequence covered 114 the 2009-2014 period: five successive crops for the PHEP and L-EN systems (2009-2013), and six for the No-Pest 115 and L-GHG systems (2009-2014).

116

#### 117 **2.4. Measurements**

#### 118 **2.4. 1. Calculation of indicators**

Assessment of the environmental performance of the cropping systems was based on energy consumption, GHG emissions, C sequestration and various environmental criteria, for real practices in the cropping system experiment. Each environmental indicator was calculated over an entire crop sequence, and expressed on a per hectare and per year basis. Criter® software (V4.0.), based on the Indigo® method and easy to manage, was used to calculate a set of environmental indicators taking values of 1 (worst) to 10 (best), with 7 selected as the target value for the entire crop sequence (Bockstaller *et al.*, 2009; Reau *et al.*, 2012).

125

#### 126 2.4.2. Pesticide indicators

127 Three pesticide indicators provided qualitative information about the volatilization, runoff and leaching into 128 groundwater of pesticides, thereby providing an indication of potential environmental damage. The treatment 129 frequency index (TFI), developed by Gravesen (2003) and widely used to assess cropping systems in France 130 (Ecophyto R&D, 2011; Jacquet et al., 2011), was also calculated, to assess the intensity of pesticide (fungicides, 131 herbicides, insecticides, molluscicides) use. This index takes into account the number of pesticide applications and the amounts applied. For each crop, TFI was calculated as follows:  $TFI = \sum_{T} {}^{AD_{T}}/{}_{RD_{T}}$ , where T is the pesticide 132 133 application, AD is the amount applied per hectare (l.ha<sup>-1</sup> or kg.ha<sup>-1</sup>) and RD is the amount authorized per hectare 134 (l.ha<sup>-1</sup> or kg.ha<sup>-1</sup>) (OECD http://www.oecd.org/site/worldforum/33703867.pdf; Pingault et al., 2009). The 135 recommended doses were those indicated in the E-phy database of the French Ministry of Agriculture (Ephy 136 website, 2014). This indicator describes pesticide use through a single synthetic variable, facilitating comparison 137 between systems. TFI, TFIH and "TFI others" correspond to overall pesticide use, herbicide use and the use of 138 fungicides plus insecticides plus molluscicides, respectively. Neither growth regulators nor nematicides were 139 sprayed on crops.

140

#### 141 **2.4.3.** Energy consumption, energy output and energy use efficiency

Energy consumption was assessed with the GES'TIM database (2010). Direct and indirect non-renewable energy consumption (*i.e.* energy inputs, EI, expressed in MJ.ha<sup>-1</sup>.year<sup>-1</sup>) corresponded to the fuel, lubricants and electricity used to power farm machinery and tractors. Indirect energy consumption was defined as the energy used in the manufacture, formulation, packaging and maintenance of inputs, such as machinery, fertilizers and pesticides.

146

Energy outputs (EO, expressed in MJ.ha<sup>-1</sup>.year<sup>-1</sup>) were calculated as the gross energy content of the harvested products. This indicator was calculated for each crop in each year, as follows: EO = Y \* CV, where Y is the yield of the harvested crop (t.ha<sup>-1</sup>), and CV is its calorific value (MJ.t<sup>-1</sup>). Yield values were calculated as the mean of six samples (each from an area of 75 to 140 m<sup>2</sup>, depending on the length of the plot harvested) collected at maturity with a combine harvester from each plot. CV was assessed with the GES'TIM database (2010). Energy use efficiency (EUE) was calculated by dividing EO by EI for the whole cropping system.

153

#### 154 **2.4.4.** Carbon balance

155 Carbon balance (kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup>) was calculated taking both C sequestration in the soil and total GHG 156 emissions into account. C sequestration in the soil (kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup>) was assessed with the Simeos® tool 157 (2014), as recommended by Saffih-Hdadi and Mary (2008), and climatic data from a meteorological station located 158 150 m from the trial. The soil characteristics of the plowed layer (0-30 cm) used for the calculations were as 159 follows: clay content 20.6%, silt content 71.9%, sand content 7.4%, bulk density 1.4, initial C content 13 g.kg<sup>-1</sup> 160 dry matter, typical of soils in the Ile-de-France region. Annual yields, calculated from our experimental data, were 161 used to estimate the expected annual biomass separately for the residues above and below the ground. Direct and 162 indirect GHG emissions were estimated with the GES'TIM database (2010), with Intergovernmental Panel on 163 Climate Change coefficients, focusing on two main greenhouse gases: nitrous oxide (N<sub>2</sub>O) and carbon dioxide 164 (CO<sub>2</sub>). Direct emissions included N<sub>2</sub>O emissions from N fertilizers and the CO<sub>2</sub> produced by the combustion of 165 fossil fuels by farm machinery. The CO<sub>2</sub> respired by soil organisms was not taken into account in these 166 assessments. Indirect emissions corresponded to the use of fossil energy in the manufacture and maintenance of 167 farm inputs.

168 Carbon balance was calculated over a period of 50 years, in accordance with Intergovernmental Panel on Climate 169 Change proposals and current knowledge of C sequestration kinetics in the soil. In this calculation, any GHG 170 entering the cropping system was attributed a negative value, whereas GHG leaving the system took a positive 171 value. The overall balance was therefore positive if more GHGs were emitted than sequestered in the system.

172

#### 173 2.4.5. Nitrogen indicators

Three nitrogen indicators were calculated with the Criter® tool (v. 4.0.). Two of these indicators provided qualitative information about ammonia (NH<sub>3</sub>) volatilization and N<sub>2</sub>O emissions. NH<sub>3</sub> volatilization was assessed for each fertilizer type, set of soil chemical characteristics (specifically calcium content) and fertilizer burial status. 177 N<sub>2</sub>O emissions were calculated as described by Bouwman *et al.* (1996): the emission factor was 1.25% N<sub>2</sub>O-N per 178 kg N of spread mineral fertilizer. The target value for this indicator (*i.e.* 7) corresponds to 20 kg of NH<sub>3</sub> volatilized 179 per hectare and per year, and 3 kg of N<sub>2</sub>O emissions per hectare and per year. Nitrogen leaching into groundwater 180 was also assessed with the Criter® tool (v. 4.0.), and expressed as a quantitative value (kgNO<sub>3</sub><sup>-</sup>.ha<sup>-1</sup>.year<sup>-1</sup>). The 181 assessment took into account both the amount of fertilizer applied and the date of the application, together with 182 rainfall over the leaching periods, from the end of winter until summer and during the winter season after crop 183 harvest (i.e. from 01/08 to 31/03 at Grignon).

184

#### 185 2.4.6. Crop diversity indicator

186 This indicator takes into account both the number of different species sown in the crop sequence, and the number 187 of genotypes for each species included in the crop sequence. It is calculated at the scale of a full crop sequence. 188 The contribution of catch crops is halved, as their growth period is shorter than that of the main crop (Criter® 189 software, V4.0).

190

#### 191 **2.4.7.** Economic indicators

192 We took the variability of prices and costs over time into account, by calculating mean values for France for the 193 2005-2012 period (INSEE). Changes in CAP (Common Agricultural Policy) directives resulted in CAP subsidies 194 being based on a shorter period in 2010-2012. These subsidies averaged €325 ha<sup>-1</sup>.year<sup>-1</sup> in the Yvelines, the area 195 in which this trial was located. Gross outputs ( $\pounds$ .ha<sup>-1</sup>) were calculated by multiplying yield (t.ha<sup>-1</sup>) by the farm-gate 196 price  $(\pounds, t^{-1})$  received for harvest products. Total variable costs  $(\pounds, ha^{-1})$  included total input costs (e.g. mineral 197 fertilizer, seeds, pesticides) and machinery costs (e.g. machinery maintenance, fuel, labor for operations). The 198 costs per hectare of different operations were determined from the data in a published database specific to North-199 Eastern France in 2013. Price variability was taken into account by calculating mean fuel price (€0.8 l<sup>-1</sup>) over the 200 2008-2013 period. Gross margins (€.ha<sup>-1</sup>) were calculated as the difference between "gross outputs plus CAP 201 subsidies" and total variable costs.

202

#### 203 2.5. Three comparisons of cropping system performances

The performances of the innovative cropping systems implemented in the field trial were first compared with that of the prototype (the prototype characteristics were described by Colnenne-David and Doré, 2015a) in a multicriteria analysis for each innovative system (*i.e.* for each innovative system and for each performance, ratios were 207 calculated as follows: ex post performance lex ante performance). The performances of the three innovative 208 systems subject to constraint (the No-Pest, L-EN and L-GHG systems) were also compared to those of the PHEP 209 system, by calculating ratios as follows: for each innovative system under constraint and for each performance, 210 performance in the innovative system under constraint /performance of the PHEP system. Finally, performance 211 ratios were calculated for the four innovative systems relative to the current system in the Ile-de-France region, 212 defined on the basis of the data collected in 2006 (Agreste, https://agreste.agriculture;gouv.fr/; Colnenne-David 213 and Doré, 2015a) (i.e. for each innovative system (the PHEP, No-Pest, L-EN and L-GHG systems) and for each 214 performance, ratios were calculated as follows: performance of the innovative system /performance of the current 215 system in the Ile-de-France region).

216

#### 217 2.6. Statistical and multi-criteria analyses

The performance and yield data were analyzed with by comparing means and carrying out analysis of variance (ANOVA) with R statistical core software (R Development Core Team R, 2014). If the result was significant (p < 0.05), the Tukey test for multiple comparisons was performed, for means with a *p*-value of 0.05 or less. When the variance was zero (*e.g.* the TFI values of all replicates of the No-Pest system were zero), only the confidence intervals (p < 0.05) were calculated.

223

#### 224 <u>3. Results</u>

#### 225 **3.1.** Assessment of the environmental performance of cropping systems

226 3.1.1. Pesticide use

# 3.1.1.1 The pesticide constraint in the No-Pest cropping system: comparison between the No-Pest and PHEP systems

229 The pesticide constraint was satisfied because no pesticides were applied in the No-Pest cropping system.

#### 230 **3.1.1.2.** Pesticide use in the four innovative cropping systems

The values of zero obtained for TFI, TFIH and TFIothers in the No-Pest system were significantly lower (p < 0.05) than those calculated for the other three innovative systems (table 2). TFI values were not significantly different (p < 0.05) between the three systems using pesticides (*i.e.* the PHEP, L-GHG and L-EN systems). In our experimental conditions, the association of no-plow practices with flax crops resulted in the highest levels of herbicide use, with significantly higher TFIH values for the L-EN system than for the other three systems (TFIH values for the various systems: L-EN=2.03; L-GHG=1.67; PHEP=1.23; No-Pest=0). Moreover, TFIothers was 237 significantly higher in the L-GHG system than in the other three systems (TFIothers values for the various systems: 238 L-GHG=1.01; PHEP=0.70; L-EN=0.35; No-Pest=0). Crop residues were not buried, and molluscicides were more 239 frequently required for slug control than in the other systems (0.5 treatments per year in the L-GHG system, versus 240 0.2 and 0.1 treatments per year in the PHEP and L-EN systems, respectively). The inclusion of winter oilseed rape 241 in the crop sequence resulted in higher levels of fungicide use: 0.3 treatments per year were applied in both the L-242 GHG and PHEP systems, whereas no fungicide was applied in either the L-EN or the No-Pest system (details in 243 table 3). Overall, "TFIothers" values, which included data for fungicides, were low, due to climatic conditions 244 unfavorable for disease development over the 2009-2014 period (Agreste, 2014).

245

#### 246 **3.1.2.** Energy use

# 3.1.2.1. The energy constraint in the L-EN cropping system: comparison between the L-EN and PHEP systems

249 Mean total fossil energy consumption (direct and indirect energy) was  $7755 \pm 711$  MJ.ha<sup>-1</sup>.year<sup>-1</sup> for the PHEP 250 system and 5201  $\pm$  502 MJ.ha<sup>-1</sup>.year<sup>-1</sup> for the L-EN system; energy consumption was thus 33% lower for the L-251 EN system (table 4). The energy constraint target (half the energy consumption of the PHEP system) was therefore 252 not met, although the decrease was nevertheless considerable. Indirect energy consumption, which accounted for 253 almost 50% of total energy consumption in both cropping systems, was 37% lower in the L-EN system (2584  $\pm$ 254 479 MJ.ha<sup>-1</sup>, year<sup>-1</sup>) than in the PHEP system (4090  $\pm$  489 MJ.ha<sup>-1</sup>, year<sup>-1</sup>). The mean amounts of N fertilizer, the 255 largest contributor to indirect energy consumption, were 19 kgN.ha<sup>-1</sup>.year<sup>-1</sup> for the L-EN system and 56 256 kgN.ha<sup>-1</sup>.year<sup>-1</sup> for the PHEP system (table 3). Direct energy consumption, defined as energy used exclusively by 257 farm machinery (i.e. for plowing, tillage, sowing, fertilization, crop protection and harvest, table 3), was 29% 258 lower in the L-EN system (2618  $\pm$  171 MJ.ha<sup>-1</sup>.year<sup>-1</sup>) than in the PHEP system (3665  $\pm$  223 MJ.ha<sup>-1</sup>.year<sup>-1</sup>), 259 mostly due to direct drilling and the absence of tillage.

260

#### 261 **3.1.2.2.** Energy performance of the four innovative cropping systems

Total energy consumption was not significantly different in the PHEP (7755  $\pm$  711 MJ.ha<sup>-1</sup>.year<sup>-1</sup>), No-Pest (7604  $\pm$  517 MJ.ha<sup>-1</sup>.year<sup>-1</sup>) and L-GHG (7459  $\pm$  793 MJ.ha<sup>-1</sup>.year<sup>-1</sup>) systems and was significantly higher than that in the L-EN system (5201  $\pm$  502 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, p<0.05). A similar pattern was observed for both indirect and direct energy consumption (*i.e.* lowest values for the L-EN system, p<0.05, table 4). An analysis of energy components revealed differences between systems. In the No-Pest system, direct energy consumption was significantly higher 267 (4417  $\pm$  425 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, p < 0.05) than that in the other systems, due to the large number of plowings (four 268 plowings over the six-year crop sequence, table 3). The indirect energy consumption linked to fertilization (table 269 3) was significantly greater in the PHEP (4090  $\pm$  489 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, p < 0.05) and L-GHG (4897  $\pm$  568 270 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, p < 0.05) systems than in the other two systems (the No-Pest system: 3187  $\pm$  99 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, the 271 L-EN system: 2584  $\pm$  479 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, table 4).

272

The L-EN system generated significantly less energy than the other systems (70997  $\pm$  9991 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, p < 0.05, table 4). The high degree of variability of the energy output of this system was linked to the low winter wheat yield in 2012 (yield of 0.75 t.ha<sup>-1</sup>, replicate 3, table 5), due to the development of highly competitive white clover. In the No-Pest system, despite low yields for most crops, energy output was high (103323  $\pm$  3629 MJ.ha<sup>-1</sup>.year<sup>-1</sup>), due to the production of hemp (mean yield value of 11.23 t.ha<sup>-1</sup>, table 5, with a calorific value of 1.65 MJ.t<sup>-1</sup>). However, it was not significantly different from that calculated for the PHEP (95965  $\pm$  8397 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, p < 0.05) and L-GHG (90229  $\pm$  5572 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, p < 0.05) systems.

280

The higher energy use efficiency of the No-Pest system (13.61  $\pm$  0.54, ns) than of the L-GHG system (12.14  $\pm$ 0.74, ns) resulted principally from its higher energy output, with no significant difference in the total energy consumption of these two systems (table 4). The energy efficiency value of the L-EN system was high (13.71  $\pm$ 2.10), but not significantly different from the other systems, due to the high level of energy output variability for the L-EN system (70997  $\pm$  9991 MJ.ha<sup>-1</sup>.year<sup>-1</sup>, table 4).

- 286
- 287 **3.1.3.** Carbon balance performance

3.1.3.1. The carbon balance constraint in the L-GHG cropping system: comparison between the L-GHG
 and PHEP systems

The carbon balances of the L-GHG ( $1202 \pm 86 \text{ kgCO}_2\text{eq.ha}^{-1}$ .year<sup>-1</sup>) and PHEP ( $1188 \pm 270 \text{ kgCO}_2\text{eq.ha}^{-1}$ .year<sup>-1</sup>) systems were not significantly different (p < 0.05, table 6). The carbon balance constraint (halving the emissions relative to the PHEP system) was not, therefore, achieved. For both systems, total greenhouse gas emissions and C sequestration accounted for nearly 90% and 10% of the carbon balance, respectively. There was no significant difference between these two systems in terms of total, direct and indirect greenhouse gas emissions (p < 0.05, table 6), resulting in similar ratios of direct and indirect greenhouse gas emissions for the two systems. The difference in direct greenhouse gas emissions between the L-GHG ( $541 \pm 102 \text{ kgCO}_2\text{eq.ha}^{-1}$ .year<sup>-1</sup>) and PHEP ( $622 \pm 82$ 

kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup>) systems was linked to an absence of plowing and only a few shallow tillage operations in the L-GHG system, whereas the plot was plowed once and subjected to numerous shallow tillage operations over the course of the crop sequence in the PHEP system (table 3). In the L-GHG system, indirect greenhouse gas emissions  $(511 \pm 82 \text{ kgCO}_2\text{eq.ha}^{-1}.\text{year}^{-1})$  were higher due to the high seed requirement: (1) the number of seeds sown in noplow conditions was systematically greater than that sown in current systems, in accordance with technical references, (2) emergence failure was observed for winter rapeseed in 2009 and 2014, and for maize and spring field beans in 2011, leading to a second sowing, and (3) cover crops were sown systematically each year (table 3).

305 After the first crop sequence, C sequestration was  $-149 \pm 117$  kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup> for the L-GHG system and - $117 \pm 150 \text{ kgCO}_2\text{eq.ha}^{-1}$ .vear<sup>-1</sup> for the PHEP system (negative values indicate a decrease in CO<sub>2</sub> relative to initial 306 307 C content, *i.e.* 13 g.kg<sup>-1</sup> dry matter) and this difference between these two systems was not significant. Cover crop 308 biomasses were lower than expected (data not shown) in the L-GHG system, due both to the high frequency of 309 very dry summers (in 2009 and 2012, total rainfall in August was 7 mm and 29 mm, respectively, whereas the 20-310 year mean value for rainfall in August was 51 mm) and the high degree of competition with weeds (data not 311 shown). In addition, yields for spring field bean and winter oilseed rape (in 2014) were lower than expected (table 312 5, see explanations below). These crop residues did not, therefore, increase the C content of the soil.

313

#### 314 **3.1.3.2.** Carbon balance of the four innovative cropping systems

315 Carbon balance did not differ significantly between the four systems (p < 0.05, table 6). However, the similarities 316 in carbon balance resulted from very different combinations of the two components of this balance: total 317 greenhouse gas emissions and C sequestration. The proportions of the two components were almost identical for 318 the L-GHG and PHEP systems. In the L-EN system, total greenhouse gas emissions were significantly lower (554 319  $\pm$  107 kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup>, p < 0.05) than those of the L-GHG (1052  $\pm$  183 kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup>) and PHEP (1071 320  $\pm$  145 kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup>) systems, and were linked to significantly lower direct and indirect greenhouse gas 321 emissions than for the other two systems (p < 0.05, table 6). The significantly lower level of N fertilization in the 322 L-EN system (19  $\pm$  6 kgN.ha<sup>-1</sup>.year<sup>-1</sup>, p < 0.05, table 3) than in the L-GHG (57  $\pm$  13 kgN.ha<sup>-1</sup>.year<sup>-1</sup>) and PHEP 323  $(56 \pm 11 \text{ kgN.ha}^{-1}.\text{year}^{-1})$  systems led to low direct and indirect N<sub>2</sub>O emissions (*i.e.* use over input manufacture). 324 However, the low yields in the L-EN system resulted in small amounts of crop residues (table 5), leading, in turn, 325 to a sharp decrease in C sequestration. The performance of the L-EN system was thus poorer than that of the L-326 GHG and PHEP systems. The No-Pest system had intermediate total greenhouse emissions ( $844 \pm 46 \text{ kgCO}_2\text{eq.ha}$  <sup>1</sup>.year<sup>-1</sup> table 6). In this system, direct emission levels were high, due to four plowing and several tillage operations during the six-year crop sequence (table 3), but indirect emissions were low, due to the low N fertilizer requirements (low yield objectives close to those in organic systems, 4.7 t.ha<sup>-1</sup>, table 5). In this system, intensive plowing practices (table 3) and the small amounts of crop residues due to low yields (table 5) both resulted in much

- 331 lower levels of C sequestration (-560  $\pm$  49 kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup>).
- 332

#### **333 3.1.4. Other environmental performances**

The values obtained for the various qualitative indicators were lowest for crop diversity indicators (ranging from 6.8, for the PHEP system, to 7.8 for the L-EN system), whereas the other environmental indicators reached values of at least 8.4 (table 7). The environmental targets may therefore be considered to have been achieved. These findings varied little between replicates.

338 The PHEP system generated significantly less N<sub>2</sub>O (8.69  $\pm$  0.16, p < 0.05) than the L-EN system (9.17  $\pm$  0.06), 339 due to differences in the amounts of N fertilizer applied (see explanation above). The amount of nitrogen leached 340 was very small in all systems (less than 10 kgN.ha<sup>-1</sup>.year<sup>-1</sup>), due to the small amounts of N fertilizer applied (table 341 3). In the cropping system currently used in Ile-de-France, the mean amount of N fertilizer applied was about 110 342 kgN.ha<sup>-1</sup>.year<sup>-1</sup>, whereas the mean amount of fertilizer applied in the PHEP system was  $56 \pm 11$  kgN.ha<sup>-1</sup>.year<sup>-1</sup>. 343 Furthermore, careful adjustment of N application dates according to plant N requirements (not shown in table 3) 344 and/or regular soil cover with plants or crop residues over time (i.e. catch or cover crops present most of the time 345 between main crops, resulting in only short periods of bare soil) could explain these results.

346

The values of all pesticide indicators were greater than 8. These results were generally consistent with the TFI values obtained. However, it is difficult to explain the small differences between the innovative systems. The main findings were the significantly higher scores for the No-Pest system (p < 0.05) and the low level of variability between the replicates of each innovative system.

351

352 **3.2. Yield** 

#### 353 **3.2.1.** The PHEP system

354 Yield objectives (table 5) were regularly achieved, for all crops except winter faba bean (mean decrease in yield 355 of almost 50%:  $1.45 \pm 0.31$  t.year<sup>-1</sup> versus 3.0 t.year<sup>-1</sup> expected), and were even higher than expected for winter 356 oilseed rape (higher yields than expected:  $3.64 \pm 0.29$  t.year<sup>-1</sup> versus 2.8 t.year<sup>-1</sup> expected). During the first three 357 years of the field assessment, very long cold winter periods destroyed many legume plants and delayed growth in 358 the spring, thereby decreasing potential yield (i.e. in 2009, 2010 and 2011 10-day minimum temperatures from the 359 beginning of December to the end of February, were -7.23°C, -5.2°C and -2.4°C respectively, whereas the mean 360 10-day minimum temperature calculated over a 20-year period was systematically above 0°C. In 2009, 2010 and 361 2011, 10-day minimum temperatures below 0°C were observed from 1/12/2008 to 20/02/2009, from 10/12/2009 362 to 10/02/2010 and from 01/12/2010 to 30/01/2011). No-till practices may also decrease potential yield, consistent 363 with the results obtained for the L-EN system (3.11  $\pm$  0.97 t.year<sup>-1</sup>). In this system, winter faba bean yields in 364 replicates 1 (2.88 t, year<sup>-1</sup>) and 3 (2.28 t, year<sup>-1</sup>) were much lower than those in replicate 2 (4.16 t, year<sup>-1</sup>), in which 365 plowing took place (in 2009, all the plots of the L-EN system were plowed, to homogenize soil structure in the 366 trial).

367

#### 368 3.2.2. The No-Pest system

Winter wheat yields were systematically higher than expected  $(6.38 \pm 1.47 \text{ t.year}^{-1} \text{ and } 6.40 \pm 0.36 \text{ t.year}^{-1} \text{ rather}$ than the 4.7 t.year<sup>-1</sup> expected), due to the low pest pressure over this period as a result of specific climatic conditions (*i.e.* very long cold winters in four of the six years and very dry conditions in spring in 2009 and 2011), resulting in an absence of disease outbreaks in spring (http://agriculture.gouv.fr/bulletins-de-sante-du-vegetal). Hemp yields were both very high and variable (11.23 ± 2.65 t.year<sup>-1</sup> versus 8.0 t.year<sup>-1</sup> expected), highlighting the underestimation of the target to be attained in a region without relevant references, and improvements in crop management over time.

376

#### **377 3.2.3. The L-EN system**

378 In the L-EN system, the yields of winter wheat, sown after winter faba bean, were higher than expected (6.46  $\pm$ 379 0.51 t.year<sup>-1</sup> versus 5.4 t.year<sup>-1</sup> expected), due to optimal use of the N provided by this legume (the objective yields 380 for faba bean were achieved, see above), in conditions in which small amounts of N fertilizer were applied. Yield 381 varied considerably between replicates for winter wheat following flax  $(4.40 \pm 1.45 \text{ t.year}^{-1})$ . The lowest yield 382 obtained for winter wheat, sown after flax (yield of 0.8 t.ha<sup>-1</sup> in replicate 3, table 5), resulted from high levels of 383 competition with white clover and weeds (data not shown). The high levels of herbicide use on both flax and winter 384 wheat (six and three applications on these two crops, respectively, table 3) reflect the high degree of weed 385 development.

386

#### **387 3.2.4.** The L-GHG system

388 Yield goals were not always reached, but the results obtained differed between crops. Winter wheat yields were 389 higher than expected (7.38  $\pm$  015 t.year<sup>-1</sup> versus 6.7 t.year<sup>-1</sup> expected), whereas spring faba bean yields were much 390 lower than anticipated (*i.e.* 66% lower than the target yield on average,  $1.36 \pm 0.71$  t.year<sup>-1</sup> versus 4.1 t.year<sup>-1</sup> 391 expected). These low yields reflected severe black aphid attacks in 2011 (i.e. 0.61 t.ha<sup>-1</sup>, replicate 3, 392 http://agriculture.gouv.fr/bulletins-de-sante-du-vegetal). Moreover, no-till practices are also known to reduce 393 yield. In the L-GHG and No-Pest systems (*i.e.* without and with plowing, respectively, table 3), spring faba bean 394 yields were  $1.36 \pm 0.71$  t.year<sup>-1</sup> and  $2.41 \pm 1.98$  t.ha<sup>-1</sup>, respectively. Winter oilseed rape yields varied considerably 395 between years  $(2.43 \pm 2.14 \text{ t.year}^{-1})$ , with the lowest value obtained for 2014 (*i.e.* 0 t.ha<sup>-1</sup>, replicate 1). In our trial 396 conditions, no-plow practices over a six-year period led to a gradual increase in the weed population (data not 397 shown), resulting in an increase in herbicide use (one, two, two, three, and four herbicides used per year in 2009 398 to 2014, respectively; table 3). In the face of such weed competition, winter rapeseed was cut at the flowering stage 399 in 2014.

400

#### 401 **3.2.5.** Impact of particular annual weather conditions and role of the crop preceding the trial

402 Weather conditions explained some low yields in the innovative systems: in 2009. Low yields for maize in both 403 the No-Pest (replicate 1, 3.81 t.year<sup>-1</sup> versus 5.6 t.year<sup>-1</sup> expected) and L-GHG (replicate 3, 5.27 t.year<sup>-1</sup> versus 7.0 404 t.year<sup>-1</sup> expected) systems were linked to a very dry summer period (*i.e.* in July and August 2009, 44 mm of rainfall: 405 calculated during a period for which the 20-year mean was 114 mm; https://donneespubliques.meteofrance.fr/); in 406 2012, the lowest yield of flax (replicate 2, 0.86 t.year<sup>-1</sup> versus 1.6 t.year<sup>-1</sup> expected) resulted from a very cold period 407 in February (*i.e.* during the first 10 days of February 2012, the mean minimum temperature was -5.5°C; the 20-408 year mean minimum temperature for the corresponding period was 4.8°C) that required a second sowing (*i.e.* of 409 spring flax). In the L-EN system, the high spring oat yield in replicate 2 (6.06 t.year<sup>-1</sup> versus 3.2 t.year<sup>-1</sup> expected) 410 resulted from combination of good weather conditions, low pest а very pressure 411 (http://agriculture.gouv.fr/bulletins-de-sante-du-vegetal) and high nitrogen availability due to the earlier sowing 412 of a legume catch crop (*i.e.* data not shown). Variability may also be linked to features specific to the trial: the 413 sowing of winter barley as the prior crop in 2008 led to the development of winter wheat root disease (i.e. 414 *Gaeumannomyces graminis*, data not shown) in the No-Pest system, replicate 3 ( $6.38 \pm 1.47 \ 1.6 \ t.year^{-1}$ ), in 2009.

415

#### 416 **3.2.6.** Variability over time

417 Yields were fairly stable in the PHEP, L-EN and No-Pest systems. By contrast, in the L-GHG system, yields 418 reached expected levels in the first four years of the crop sequence, but were lower in the last two years (*e.g.* for 419 oilseed rape in replicate 1: 0.00 t.year<sup>-1</sup> versus 2.8 t.year<sup>-1</sup> expected). Both large increases in the weed population 420 and changes in soil structure (data not shown) due to an absence of plowing gradually reduced crop yields in our 421 field conditions.

422

#### 423 **3.3. Economic results**

424 Gross margins were highest for the PHEP (757.0 ± 88.7 €.ha<sup>-1</sup>.year<sup>-1</sup>), No-Pest (701.4 ± 48.4 €.ha<sup>-1</sup>.year<sup>-1</sup>) and L-425 GHG (619.6 ± 77.3 €.ha<sup>-1</sup>.year<sup>-1</sup>) systems (table 8). Furthermore, gross margins were significantly higher for the 426 PHEP system than for the L-EN system (606.4  $\pm$  56.1  $\in$ .ha<sup>-1</sup>.year<sup>-1</sup>, p < 0.05). The similar results obtained for the 427 L-GHG and L-EN systems resulted from different combinations of gross outputs and total variable costs. In the L-428 GHG system, high total variable costs (567.7 ± 29.7 €.ha<sup>-1</sup>.year<sup>-1</sup>) counteracted the high gross output (861.9 ± 429 84.0 €.ha<sup>-1</sup>.year<sup>-1</sup>). In the L-EN system, both gross output (696.0  $\pm$  74.5 €.ha<sup>-1</sup>.year<sup>-1</sup>) and total variable costs (415.0 430  $\pm$  46.7 €.ha<sup>-1</sup>.year<sup>-1</sup>) were lower than in any other system because (i) less N fertilizer was applied than in the PHEP 431 and L-GHG systems, (ii) the no-till practices resulted in lower levels of fuel consumption and machinery use than 432 for the No-Pest system, which was characterized by several plowing and tillage operations over the course of the 433 crop sequence (table 3). The gross margin of the No-Pest system was one of the highest (701.4 ± 48.4 €.ha<sup>-1</sup>.year<sup>-1</sup> 434 <sup>1</sup>), due to hemp and winter wheat yields both being higher than expected (table 5).

435

#### 436 **3.4.** Performance comparisons between the prototype systems and the field trial assessments

For the four innovative systems, most environmental performance indicators (total GHG emissions, total energy consumption, energy output, energy efficiency) and gross margins were close to the predictions of *ex ante* assessments (figure 1). The specific environmental constraints of the No-Pest and L-EN systems were almost satisfied. TFI, TFIH and "TFIothers" in the No-Pest system, and total energy consumption in the L-EN system closely matched expectations. For the L-GHG system, total greenhouse gas emissions were as projected, whereas C sequestration levels were much lower than expected.

443

444 A comparison of *ex ante* and *ex post* assessments showed large differences for TFI, TFIH and "TFIothers" (figure

445 1). In the L-GHG and L-EN systems, herbicide applications were underestimated in the prototype systems, and

446 TFIH was four times higher for the L-GHG, and two times higher for the L-EN in the field assessments than

447 estimated for the prototypes. TFIH was also higher than expected in the PHEP system, but to a lesser extent.

448 During the design process, "TFIothers" was systematically overestimated because it could not take into account

the specific low pest pressures occurring over the 2009-2014 period. For each innovative system, the energy output

450 results measured in field conditions were very close to the expected values.

451

# 452 3.5. Performance comparisons between the innovative cropping systems subject to constraints and the 453 PHEP system, taken as the reference system

454 The PHEP system performed particularly well, so the three constraint-limited systems performed poorly by 455 comparison (figure 2). In both the L-GHG and L-EN systems, TFI, TFIH and "TFIothers" were higher than those 456 calculated for the PHEP system (see the explanations above). However, these two systems under constraints 457 differed for other performances. Most environmental performances were similar for the L-GHG and PHEP 458 systems, whereas the L-EN system outperformed the reference system. In both the L-GHG and L-EN systems, 459 gross margins were lower than those in the PHEP system, due to lower yields (table 5). This was unexpected for 460 the L-GHG system, but was anticipated at the design step for the L-EN system (*i.e.* this system was designed with 461 a target yield 20% lower than that of the PHEP system, to satisfy the energy constraint; Colnenne-David and Doré, 462 2015a). For a similar gross margin, pesticide indicator performances in the No-Pest system were much better than 463 those in the PHEP system, but were associated with poor direct energy and C balance performances.

464

### 465 **3.6.** Performance comparisons between the innovative systems implemented in the field trial and the current

466 system in the Ile-de-France region

467 Comparison between the four innovative systems and the current system in the Ile-de-France region (figure 3) 468 demonstrated that all environmental performances were better in the innovative systems (*i.e.* all ratios below 1) 469 than in the current system. Moreover, despite the lower energy outputs of the new systems than of the current 470 system, gross margins were similar or slightly higher in the new systems than in the current Ile-de-France system. 471 However, in the L-EN and No-Pest systems, TFIH and direct energy consumption, respectively, were similar to 472 those for the current system (*i.e.* ratio values close to 1).

473

#### 474 **<u>4. Discussion</u>**

#### 475 **4.1.** Achievement of a multiplicity of objectives

476 We were able to design and implement the PHEP system, the environmental performances of which were better 477 than those of the current system in the region, with no decrease in gross margin. The absence of pesticide use in 478 the No-Pest system did not reduce gross margin either (the lower target yield resulted in an absence of impact on 479 yield performance in our trial), but improved environmental performance (low greenhouse gas emissions, high 480 energy use efficiency, low nitrate leaching). However, higher levels of direct energy consumption, linked to the 481 high frequency of tillage practices, resulted in lower levels of C sequestration. It was possible to decrease energy 482 consumption in the L-EN system only with a decrease in yield, resulting in a lower gross margin, and low levels 483 of C storage in the soil. However, with the exception of the herbicide indicator, most of the environmental 484 performances were fine. The management of agronomic strategies in the L-GHG system led to high yield 485 variability, with a low economic impact. All environmental performances were satisfactory, with the exception of 486 the herbicide use indicator, which was similar to that for the current system in the region.

487

488 As discussed in previous studies (Colnenne-David and Doré, 2015a), the various targets set for innovative systems 489 can be antagonistic. The imposition of strong environmental constraints modified the performances of the 490 constrained systems. Some performances deteriorated. In both the L-GHG and L-EN systems, no-plow practices 491 led to higher levels of herbicide use to destroy cover crops and weeds (high TFIH), as previously reported by 492 Zetner et al. (2004), Moreno et al. (2011), and Soane et al. (2012). In the L-EN system, lower levels of energy 493 consumption, due to both no-till practice and low levels of N fertilization, were associated with 20% lower yields. 494 In the No-Pest system, the absence of pesticide use had an adverse effect on SOM and yield. Decreases in the 495 frequency of tillage and target yields resulted in much lower levels of C sequestration. Conversely, some 496 environmental performances were significantly improved by the imposition of a severe environmental constraint. 497 In both the L-EN and No-Pest systems, gas balance and energy efficiency were as high as those in the reference 498 PHEP system. Economic comparisons with published findings were difficult, because the prices of both inputs 499 and outputs depend on the country concerned, the period analyzed and the cropping system used (organic farm 500 produce is sold at higher prices than the products of conventional agriculture). We then compared the gross margins 501 of the innovative systems and the current system in Ile-de-France, in one price context: gross margins were slightly 502 lower than those of the regional system for the L-GHG and L-EN systems, and slightly higher for the other new 503 systems. However, this initial assessment did not take into account the contribution of product quality to farm-gate 504 price, which is potentially higher for free-pesticide seeds, and the existence of specific markets for crops such as 505 hemp.

506

507 It was difficult to meet the energy constraint in the L-EN system and environmental performances were less 508 satisfactory (specifically for herbicide use) than in the PHEP system. It was not possible to satisfy the greenhouse 509 gas constraint in the L-GHG system. For this system, during the design step, a clear hierarchy between the two 510 sub-objectives (*i.e.* to enhance carbon sequestration first and then to reduce N<sub>2</sub>O emissions) were defined. In our 511 field conditions, this strategy was not effective. Biomass production was low (see the above comments for yields) 512 and resulted in lower levels of carbon storage than expected. Moreover, the amount of N fertilizer required to 513 produce the expected biomass did not differ between the L-GHG and PHEP systems (*i.e.* total greenhouse gas 514 emissions did not differ significantly between these two systems, table 6). After the first crop rotation, another 515 design step was required to improve the L-GHG system, and a new combination of agricultural practices is 516 currently being assessed in the field. The environmental results of the PHEP system were also very good, making 517 it difficult to achieve both the energy goal in the L-EN system and the greenhouse gas target in the L-GHG system.

518

#### 519 4.2. Difficulties implementing innovative systems with multiple goals in the field

520 Overall, the predictive capacity of ex ante assessment was good. However, discrepancies between the estimated 521 performance of prototype systems and trial results, with some goals not achieved or the occurrence of unexpected 522 environmental conditions, highlighted the difficulties involved in managing such systems in the field. We 523 investigated the reasons for these differences, by analyzing agronomic practices, which we classified into four 524 groups. Group 1: the chosen agronomic strategies were unsuitable for achieving the goals set. For example, in the 525 L-GHG system, the absence of plowing did not lead to an increase in C sequestration. Group 2: some practices 526 were unable to satisfy multiple goals simultaneously. For example, in the No-Pest system, the restitution of small 527 amounts of organic matter, due to low yields, combined with regular plowing, which was required to manage weed 528 populations, had an adverse effect on C sequestration. Group 3: some of the planned practices may not have been 529 appropriate in field trial conditions. For example, despite the setting of TFI targets based on local experimental results obtained over a 10-year period, the "TFIothers" and TFIH values obtained did not match expectations. 530 531 During the design process, pest occurrence rates were overestimated, resulting in higher levels of pesticide use 532 estimated for the prototypes than actually applied in the field, except for the No-Pest system 533 (http://agriculture.gouv.fr/bulletins-de-sante-du-vegetal, see the explanations above). Group 4: an unpredicted 534 change occurred in the agrosystem. For example, weed levels were higher after flax in the L-EN system, resulting 535 in higher levels of herbicide use than anticipated in several years (table 3). Similarly, weed populations increased throughout the crop sequence in the L-GHG system, resulting in larger slug populations (data not shown). Despite two molluscicide applications in 2014, oilseed rape was sown twice with no final yield (*i.e.* in 2014, yield was 0.00 t.ha<sup>-1</sup>; in replicate 1, table 5). This classification highlighted the need for more time to eliminate technical uncertainties and to improve the management of innovative systems, to prevent the technical problems observed here (sowing failure, bad weed management in no-till systems). Moreover, the use of a broad range of tools should make it possible to improve the predictive capacity of *ex ante* assessment.

542

#### 543 **4.3.** Comparison of performances with published results

#### 544 **4.3.1.** Energy consumption

545 The total energy consumption per hectare of the innovative systems was similar to that reported by Zentern et al. 546 (2004) for different winter wheat-based crop sequence plow practices, and by Planche et al. (2015) for different 547 cropping systems designed to meet specific environmental goals and assessed in France. As shown by Zentern et 548 al. (2004) and regularly confirmed by different authors (Dumaski et al., 2006; Rothke et al., 2007; Morano et al., 549 2011), the reduction of energy consumption due to no-till practices was generally offset by an increase in herbicide 550 use. A similar pattern was observed when the energy performances of the PHEP and L-GHG systems were 551 compared. The contribution of N fertilization to the overall energy consumption of the new systems was similar 552 to that calculated for conventional, minimum tillage and no-till systems by Zentner et al. (2004), Rothke et al. 553 (2007) and Moreno et al. (2011). Nevertheless, the energy consumption of an "integrated" system, such as that 554 described by Nemecek et al. (2011b), with similar amounts of applied nitrogen to the PHEP system, was 555 significantly greater than that calculated for the PHEP system. However, more details of the practices used in the 556 Swiss "integrated" system, and of the references used for energy calculations, are required to analyze this 557 discrepancy. By contrast, the energy consumption of the No-Pest system due to chemical fertilization was greater 558 than that for organic systems using organic fertilizers (Morano et al., 2011; Nemecek et al., 2011a).

559

#### 560 **4.3.2.** GHG emissions

561 Goglio *et al.* (2014) used a combination of LCA and ecosystem modeling to assess GHG emissions in innovative 562 systems. Over the 2009-2012 period, global warming potential (GWP) was 1.36 to 4.25 kgCO<sub>2</sub>eq.ha<sup>-1</sup> in the PHEP 563 system. Brentrup *et al.* (2004b) reported GWP ranges of 0.29 to 4.10 kgCO<sub>2</sub>eq.ha<sup>-1</sup> for wheat with different 564 amounts of N fertilizer, and Charles *et al.* (2006) reported a value of 2.42 kgCO<sub>2</sub>eq.ha<sup>-1</sup> for the same crop. With a 565 range of 2.15 to 5.03 kgCO<sub>2</sub>eq.ha<sup>-1</sup>, the estimates for the Swiss "integrated" and organic systems involving cereals (Nemecek *et al.*, 2011b) were slightly higher than those for the PHEP system. Despite the high variability of these results, all the GWP results obtained were of the same order of magnitude. Since 2013, new cropping systems with multiple goals, including lower levels of tillage, have been assessed in field trials (Planche *et al.*, 2015). The annual GHG results calculated with the GES'TIM database (2010) ranged from 1340 to 2060 kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup>. Despite the use of different methodologies (calculation of GHG emissions at the crop sequence scale in the innovative systems), the lowest value was close to that for the PHEP system (1071 kgCO<sub>2</sub>eq.ha<sup>-1</sup>.year<sup>-1</sup>).

572

573 There is some debate about the degree to which no-till practices can increase soil organic carbon (SOC) 574 sequestration relative to conventional tillage. Conservation tillage practices, with an absence of tillage and 575 permanent soil cover, are adopted to limit the decline in SOC levels (Jonhson et al., 2007; Smith et al., 2008; Luo 576 et al., 2010). However, Dimassi et al. (2014) and Virto et al. (2012) have shown that the C input from crop residues 577 is the major factor significantly correlated with differences in SOC levels between no-till and inversion tillage 578 systems. Moreover, the SOC initially present modifies the rate of mineralization of soil biogeochemical 579 components. Initial SOC content was 31 tC.ha<sup>-1</sup> in the trials managed by Li et al. (2005), 35 tC.ha<sup>-1</sup> in the study by 580 Andriulo et al. (1999) and 42-45 tC.ha<sup>-1</sup> in that by Dimassi et al. (2014). The C sequestration process was complex, 581 due to interaction between tillage practices and the amount of crop residue present. In the No-Pest system, low 582 levels of C sequestration may be linked to both the small amounts of crop residues left on the soil and intensive 583 plowing practices during the crop sequence (table 3), consistent with current scientific knowledge. However, in 584 the L-EN system, in which only small amounts of crop residues were present, the no-till practices did not prevent 585 C sequestration from being very low. Likewise, C sequestration did not differ significantly between the PHEP and 586 L-GHG systems, despite large differences in tillage practices (table 3). Moreover, in the other studies, assessments 587 were carried out over longer periods than this study. Dimassi et al. (2014) analyzed SOC evolution after 12 years 588 of no-till practice. Bremer et al. (2008) measured changes in SOC 12 years after the introduction of fallow. The 589 impacts of different tillage practices on C sequestration were assessed over 27- and 30-year periods by Liu et al. 590 (2009) and Ghangsen Li et al. (2005), respectively. However, our results, simulated with the Simeos® tool, require 591 validation with trial measurements. They were obtained after the first complete crop sequence (*i.e.* 5 to 6 years), 592 which may be too short for the analysis of C sequestration. At least another full crop sequence may be required for 593 a reliable analysis of changes in SOC. In organic systems, SOC content is generally reported to be higher than that 594 in conventional systems, due to the use of organic fertilizers (Clark et al., 1998; Wells et al., 2000; Azeez G., 2008; 595 Mancinelli et al., 2010). The significant difference, by a factor of about five, between the No-Pest and PHEP

596 systems, may be explained by the many plowing operations and lower yields in the No-Pest system than in the

597 PHEP system, and by the removal of hemp straw.

598

#### 599 4.3.3. Yield performances

600 The target yields of the innovative systems were lower than those of conventional systems in the Ile-de-France 601 region, to make it possible to satisfy environmental targets. Yields in the PHEP system were 5% to 10% lower 602 than those of current systems, depending on the species considered, but gross margins were similar. Over the 2009-603 2013 period, mean winter wheat yield was 9.77 t.ha<sup>-1</sup> for a conventional system (Colnenne-David et al., 2015b) 604 assessed in a field trial located near Grignon (Debaeke et al., 2009), whereas mean winter wheat yield was 8.56 605 t.ha<sup>-1</sup> in the PHEP system. The corresponding TFIs were 4.64 and 1.85 and the amount of N fertilizer applied was 606 147 kgN.ha<sup>-1</sup>.year<sup>-1</sup> and 56 kgN.ha<sup>-1</sup>.year<sup>-1</sup>, respectively. The energy output of the innovative systems, with a range 607 of 71 to 103 GJ.ha<sup>-1</sup>.year<sup>-1</sup> for the L-EN and No-Pest systems, respectively, was lower than that of conventional 608 systems in the Ile-de-France region (114 GJ.ha<sup>-1</sup>.year<sup>-1</sup>). Variable energy output results have been reported for 609 conventional and organic systems (Klimekova et al., 2007; Moreno et al., 2011) from different locations and with 610 different methodologies (different ways of taking straw energy into account). In all studies, energy output was 611 systematically higher in conventional than in organic systems, contrasting with the results for the PHEP and No-612 Pest systems (96 and 103 GJ.ha<sup>-1</sup>.year<sup>-1</sup>, respectively). The high score of the No-Pest system resulted from both 613 high crop productivity, particularly for hemp, which had a mean yield of 11.23 t.ha<sup>-1</sup>, and high calorific value. 614 Without hemp in the crop sequence, energy output would probably reach about 87 GJ.ha<sup>-1</sup>.year<sup>-1</sup>. The energy use 615 efficiency of the new systems ranged from 12.1 to 13.7, and was thus much higher than published values. The 616 EUE of the current Ile-de-France system was 7.75; those for conventional and organic systems were 6.55 and 6.41, 617 respectively, over an 11-year period in Bulgaria (Bochu et al., 2008) and 7.77 and 10.57, respectively, over a six-618 year period in Poland (Klimekova et al., 2007). These high performances reflect a significant optimization of 619 agronomic practices, in terms of both plowing and N fertilizer management. Moreover, the specific climatic 620 conditions prevailing in the 2009-2014 period resulted in high yields with little or no pesticide application.

621

#### 622 <u>5. Conclusion</u>

We show here that it is possible to design and implement innovative cropping systems with multiple goals combining environment performance and economic results. However, some of these goals appear to be more easily attainable than others. In our conditions, and during the first full crop sequence in the innovative systems, the 626 application of a constraint imposing an absence of pesticide use did not result in poorer environmental and 627 economic results that were obtained with the PHEP system, despite the strong performance of the PHEP system. 628 However, our efforts to halve GHG emissions failed, due to the use of an inadequate strategy, which was 629 nevertheless based on the knowledge available at the design stage. Increasing numbers of studies of the effects of 630 agricultural practices on L-GHG emissions and carbon storage are being published, and their findings should make 631 it possible to refine our strategy on the basis of cutting-edge knowledge. The L-EN system was moderately 632 successful. It performed well, but did not quite achieve the targets set, and environmental performances were 633 declining over time, suggesting a need for adaptation of the strategy. We are currently carrying out assessments 634 for the second complete crop sequence in the same field trial (1) to validate the preliminary results for the PHEP 635 and No-Pest systems, for which agricultural practices have been kept the same as in the first crop sequence, and 636 (2) to assess the performances of new prototypes of the L-EN and L-GHG systems, which have been modified to 637 decrease herbicide use, and to make it easier to satisfy the GHG constraint of the L-GHG system. We believe that 638 such agronomic studies, combining in silico loops with field trials, are important and will facilitate the design of 639 new innovative cropping systems to deal with the range of issues faced by agriculture.

640

#### 641 Acknowledgments

Financial support was provided by the PURE project, supported by the European Commission through the Seventh Framework Programme (FP7/2011-2014) under contract number 265865. We thank Veronique Tanneau for her technical work in the field and the technical team of the "Versailles-Grignon" experimental unit for the management of agricultural practices, M. Valantin-Morison for her useful comments on a previous version of the paper and D. Makowski for his statistical advice.

#### 647 **References**

- Abdi D., Cade-Menun BJ., Ziadi, N., Parent, L. E., 2014. Long-term impact of tillage practices and phosphorus
- 649 fertilization on soil phosphorus forms as determined by 31p nuclear magnetic resonance spectroscopy. Journal of
- 650 Environmental Quality, 43, 4, 1431-1441, DOI 10.2134/jeq2013.10.0424.
- 651 Abdollahi L., Hansen E.M., Rickson R. J., Munkhohn L. J., 2015. Overall assessment of soil quality on humid
- sandy loams: Effects of location, rotation and tillage. Soil & Tillage Research, 145, 29-36, DOI:
  10.1016/j.still.2014.08.009.
- Andriulo, A., Mary, B., and Guerif, J. 1999. Modelling soil carbon dynamic with various cropping sequences on
  the rolling pampas. Agronomie 19:365–379.
- Aubertot, J.N., Barbier, J.M., Carpentier, A., Gril, J.J., Guichard, L., Lucas, P., Savary, S., Savini, I., and Voltz,
- 657 M. 2005. Pesticides, agriculture and the environment: reducing the use of pesticides and limiting their
- 658 environmental impact. Collective Scientific Expert Report. INRA CEMAGREF. INRA edition. p. 63.
- Azeez G.S.E & Hewlett K.L., 2008. The Comparative Energy Efficiency of Organic Farming. 16th IFOAM
- 660 Organic World Congress, Modena, Italy, June 16-20, 2008 http://orgprints.org/view/projects/conference.htlm.
- 661 Blazy, J.M., Ozier-Lafontaine, A., Doré, T., Thomas, A., and Wery, J. 2009. A methodological framework that
- 662 accounts for farm diversity in the prototyping of crop management systems. Application to banana-based systems
- 663 in Guadeloupe. Agricultural Systems 101:30–41.
- 664 Bochu JL., Risoud B., Mousset J., 2008. Consommation d'énergie et émissions de GES des exploitations en
- 665 agriculture biologique : synthèse des résultats PLANETE 2006. International conference Organic agriculture and
- 666 climate change, Enita de Clermont Ferrand; France, April 17-18th 2008.
- 667 Bockstaller C., Guichard L., Keichinger O., Girardin P., Galan M.B., and Gaillard G., 2009. Comparison of
- methods to assess the sustainability of agricultural systems. A review. Agronomy of Sustainable Development
  29:223–235.
- Bouwman AF., 1996. Direct emission of nitrous oxide from agricultural soils. Nutrition Cycling Agroecosystem,
  46, 53–70.
- Bremer E., Janzen H., Ellert B., McKenzie R., 2008. Soil organic carbon after twelve years of various crop
  rotations in an aridic boroll. Soil and Water Management and Conservation, SSAJ: Volume 72: Number 4.
- 674 Brentrup F, Küsters J, Lammel J, Barraclough P, Kuhlmann H. Environmental impact assessment of agricultural
- 675 production systems using the life cycle assessment (LCA) methodology II. The application to N fertilizer use in
- 676 winter wheat production systems. Eur J Agron 2004b; 20: 265-279.

- 677 Charles R, Jolliet O, Gaillard G, and Pellet D, 2006. Environmental analysis of intensity level in wheat crop
- 678 production using life cycle assessment. Agric Ecosyst Environ; 113(1–4):216–225.
- 679 Charles H., Godfray J., and Garnett T., 2016. Food security and sustainable intensification.
  680 http://rstb.royalsocietypublishing.org/.
- 681 Chikowo, R., Faloya, V., Petit, S., and Munier-Jolain, N. 2009. Integrated weed management systems allow
- reduced reliance on herbicides and long-term weed control. Agriculture, Ecosystems and Environment 132:237–
- 683 242.
- 684 Colnenne-David C., Doré T., 2015a. Designing innovative productive cropping systems with quantified and
- ambitious environmental goals. "Renewable Agriculture and Food Systems". doi:10.1017/S1742170514000313.
- 686 Colnenne-David C., Grandeau G., Tanneau V., Doré T., 2015b. Innovative IPM for winter wheat-based rotations:
- 687 first results of ex post sustainability assessment of cropping systems tested at INRA (France). PURE Congress:
- 688 IPM innovation in Europe, Poznan (Poland) (poster).
- 689 Debaeke, P., Munier-Jolain, N., Bertrand, M., Guichard, L., Nolot, J.M., Faloya, V., and Saulas, P. 2009. Iterative
- design and evaluation of rule-based cropping systems: methodology and case studies. Agronomy of Sustainable
  Development 29:73–86.
- 692 Dimassi B., Mary B., Wylleman R., Labreuche J., Couture D., Piraux F., Cohan JP., 2014. Long-term effect of
- 693 contrasted tillage and crop management on soil carbon dynamics during 41 years. Agriculture, Ecosystems and
- 694 Environment, 188, 134–146.
- 695 Dumanski J., Peiretti R., Benetis J., McGarry D., Pieri C., 2006. The paradigm of conservation tillage. Proceeding
  696 World Association. Soil and Water Conservation, 1, 58–64.
- 697 Ecophyto R&D, 2011. Which options to reduce pesticide use? INRA, 8 pp. http://www.international.inra.fr/the
- 698 institute/advanced studies/ecophyto r d.
- 699 FAO 1998, Topsoil characterization for sustainable land management, draft. Land and Water Division, Soil
- 700 Resources, Management and Conservation Service, Rome.
- 701 Freibauer, A., Rounsevell, M.D.A., Smith, P., and Verhagen, J. 2004. Carbon sequestration in the agricultural soils
- 702 of Europe. Geoderma 122:1–23.
- Foley, J.A., Ramankutty, N., Brauman, K.A., Cassidy, E.S., Gerber, J.S., Johnston, M., Mueller, N.D., O'Connell,
- 704 C., Ray, D.K., West, P.C., Balzer, C., Bennett, L.M., Carpenter, S.R., Hill, J., Monfreda, C., Polasky, S.,
- Rockström, J., Sheehan, J., and Siebert, S., 2011. Solutions for a cultivated planet. Nature 478:337–342.

- 706 Li C., Frokling S., Butterbach-Bahl, K., 2005. Carbon sequestration in arable soils is likely to increase nitrous
- 707 oxide emissions, offsetting reductions in climate radiative forcing. Climatic Change, 72, 321-338, DOI:
- 708 10.1007/s10584-005-6791-5.
- 709 GES'TIM 2010. Ministère de l'agriculture, de l'alimentation, de la pêche, de la ruralité et de l'aménagement du
- 710 territoire. Réf. 0933103.
- 711 Gravesen, L., 2003. The treatment frequency index: an indicator for pesticide use and dependency as well as overall
- 712 load on environment. In: Reducing Pesticide Dependency in Europe, pp. 28–30.
- 713 Goglio P., Grant B., Smith W., Desjardins R., Worth D., Zentner R., and Malhi S., 2014. Impact of management
- strategies on the global warming potential at the cropping system level. Science of the Total Environment, 490,
- 715 921-933. 10.1016/j.scitotenv.2014.05.070.
- 716 Ingle SS., Jadhao SD., Kharche VK., Sonune BA. and Mali DV., 2014. Soil biological properties as influenced by
- 717 long-term manuring and fertilization under sorghum (Sorghum bicolor) -wheat (Triticum aestivum) sequence in
- vertisols. Indian Journal of Agricultural Sciences, 84, 4, 452-457. WOS:000343587000003.
- Jacquet, F., Butault, J.P., Guichard, L., 2011. An economic analysis of the possibility of reducing pesticides in
  French field crops. Ecol. Econ. 70, 1638–1648.
- 721 Khakbazan M., Mohr R.M., Derksen D.A., Monreal M.A., Grant C.A., Zentner R.P., Moulin A.P., McLaren D.L.,
- 722 Irvine R.B., Nagy C.N., 2009. Effects of alternative management practices on the economics, energy and GHG
- emissions of a wheat-pea cropping system in the Canadian prairies. Soil & Tillage Research, 104, 30–38.
- 724 Klimeková M., Lehocká Z., 2007. Comparison of organic and conventional farming system in terms of energy
- 725 efficiency. Wissenschaftstagung Ökologischer Landbau. Beitrag archiviert unter
  726 http://orgprints.org/view/projects/wissenschaftstagung-2007.html.
- Kulak M., Nemecek T., Frossard E., Gaillard G., 2013. How eco-efficient are low-input cropping systems in
  Western Europe, and what can be done to improve their eco-efficiency. Sustainability, 5, 3722-3743.
- 729 doi:10.3390/su5093722.
- 730 Lescourret F., le Magda D., Richard G., Adam-Blondon AF., Bardy M., Baudry J., Doussan I., Dumont B., Lefevre
- 731 F., Litrico I., Martin-Clouaire R., Montuelle B., Pellerin S., Plantegenest M., Tancoigne E., Thomas A., Guyomard
- H. and Soussana JF., 2015. A social-ecological approach to managing multiple agro-ecosystem services. Current
- 733 Opinion in Environmental Sustainability 2015, 14:68–75.
- Liu D., Chan K., Conyers M., 2009. Simulation of soil organic carbon under different tillage and stubble
  management practices using the Rothamsted carbon model. Soil & Tillage Research, 104, 65–73.

- 736 Loyce C., Meynard J.M., Bouchard C., RollandB., Lonnet P., Bataillon P., Bernicot M.H., Bonnefoy M., Charrier
- X., Debote B., Demarquet T., Duperrier B., Félix I., Heddadj D., Leblanc O., LeleuM., Mangin P, Méausoone M.,
- 738 Doussinault G., 2012. Growing winter wheat cultivars under different management intensities in France: A
- multicriteria assessment based on economic, energetic and environmental indicators. Field Crops Research, 125,
- 740 167–178.
- Luo Y., Wang Z., Gao M., Wei C., 2011. Effects of conservation tillage on organic carbon, nitrogen and enzyme
  activities in a hydragric anthrosol of Chongqing, China. Energy Procedia, 5, 30–36.
- Mäder P., Fliebbach A., Dubois D., Gunst L., Fried P., Niggli U., 2002. Soil fertility and biodiversity in organic
  farming. Science, 296, 1694-1697.
- 745 Mancinelli R., Campiglia E., Di Tizio A., Marinari S., 2010. Soil carbon dioxide emission and carbon content as
- affected by conventional and organic cropping systems in Mediterranean environment. Applied Soil Ecology, 46,
- 747 64–72.
- 748 Moreno M., Lacasta C., Meco R., Moreno C., 2011. Rainfed crop energy balance of different farming systems and
- rop rotations in a semi-arid environment: Results of a long-term trial. Soil & Tillage Research, 114, 1, 18–27.
- Nemecek T., Dubois D., Huguenin-Elie O., and Gaillard G. 2011a. Life cycle assessment of Swiss farming
  systems: I. Integrated and organic farming. Agricultural Systems, 104, 217–232.
- 752 Nemecek T., Huguenin-Elie O., Dubois D., Gaillard G, Schaller B, and Chervet A., 2011b. Life cycle assessment
- of Swiss farming systems: II. Extensive and intensive production. Agricultural Systems, 104, 233–245.
- Panasiewiez K., Koziara W., and Krawczyk R., 2010. Comparison of grain sowing quality and vigor of spring
- barley grown in accordance with organic and conventional system. Journal of Research and Applications in
- 756 Agricultural Engineering, 55, 4, 42-45.
- 757 Pingault N., Pleyber E., Champeaux C., Guichard L., and Omon B., 2009. Produits phytosanitaires et protection
- 758 intégrée des cultures: l'indicateur de fréquence de traitement (IFT). Notes et Etudes Economiques, Ministère de
- 759 l'agriculture et de la pêche, 32.
- 760 Planche R., 2014. Evaluation *ex post* d'un système de culture multiobjectif sous contraintes de réduction du travail
- 761 du sol. Mémoire de fin d'études pour l'obtention du diplôme d'ingénieur AgroParisTech.
- 762 R Development Core Team, R version 3.1.2 (2014-10-31) -- "Pumpkin Helmet" Copyright (C) 2014. The R
- 763 Foundation for Statistical Computing, Platform: i386-w64-mingw32/i386 (32-bit).

- Reau R., Monnot L.A., Schaub A., Munier-Jolain N., Pambou I., Bockstaller C., Cariolle M., Chabert A., Dumans
- P., 2012. Les ateliers de conception de systèmes de culture pour construire, évaluer et identifier des prototypes
- 766 prometteurs. Innovations Agronomiques, 20, 5-33.
- Rathke G., Wienhold B., Wilhelm W., Diepenbrock W., 2007. Tillage and rotation effect on corn–soybean energy
- balances in eastern Nebraska. Soil & Tillage Research, 97, 60–70.
- 769 Saffih-Hdadi K. and Mary B.,2008. Modeling consequences of straw residues export on soil organic carbon. Soil
- 770 Biology & Biochemistry 40, 594–607.
- 571 Singh K.P., Prakash V., Srinivas K., Srivastva A.K., 2008. Effect of tillage management on energy-use efficiency
- and economics of soybean (Glycine max) based cropping systems under the rainfed conditions in North-West
- Himalayan Region. Soil & Tillage Research, 100, 78–82.
- Smith P., 2008. Land use change and soil organic carbon dynamics. Nutr. Cycl. Agroecosyst, 81, 169–178. DOI
  10.1007/s10705-007-9138-y.
- 576 Soane BD., Ball BC., Arvidsson J., Basch G., Moreno F., and Roger-Estrade J., 2012. No-till in northern, western
- and south-western Europe: A review of problems and opportunities for crop production and the environment. Soil
- 778 & Tillage Research 118,66-87.
- Tapia S., Villamil MB., Grabau M., 2014. Evaluation of N sources, cover crops, and tillage systems for corn grown
  under organic management. Phyton-International Journal of Experimental Botany, 83, 71-81.
- 781 Vaca Garcia, VM., Dominguez Lopez A., Gonzalez Huerta A. Morales Rosales EJ., Mora OF., Gutierrez
- 782 Rodriguez F., 2014. Assessment of soil compaction under different management regimes using double-cycle
- viaxial compression test. Terra Latino Americana, 32, 2, 119-126.
- 784 Vereijken, P. 1997. A methodological way of prototyping integrated and ecological arable farming systems
- 785 (I/EAFS) in interaction with pilot farms. European Journal of Agronomy 7:235–250.
- Virto I., Barre P., Burlot A., Chenu C., 2012. Carbon input differences as the main factor explaining the variability
- in soil organic C storage in no-tilled compared to inversion tilled agrosystems. Biogeochemistry, 108, 17–26.
- 788 Wells A., Chan K., Cornish P., 2000. Comparison of conventional and alternative vegetable farming systems on
- the properties of a yellow earth in New South Wales. Agriculture, Ecosystems and Environment, 80, 47–60.
- 790 Zentner, R.P., Lafond, G.P., Derksen, D.A., Nagyd, C.N., Wall, D.D., and May, W.E. 2004. Effects of tillage
- 791 method and crop rotation on non-renewable energy use efficiency for a thin Black Chernozem in the Canadian
- 792 Prairies. Soil and Tillage Research 77: 125-136

793

**Figure 1.** Multicriteria assessment of the four innovative cropping systems. Comparisons between *ex post* and *ex ante* assessments (the gray area corresponds to the *ex post / ex ante* ratio). Cropping systems: A: PHEP (productive with high environmental performance), B: L-GHG (low greenhouse gas emissions), C: L-EN (low energy use), and D: No-Pest (no pesticide use). Cg balance: carbon gas balance. C seq: carbon sequestration. Tot GHG: total greenhouse gas emissions. Tot En: total energy consumption. En op: energy output. En eff: energy efficiency. TFI: treatment frequency index. TFIH: TFI for herbicides. TFIothers: TFI for all pesticides other than herbicides. G margin: gross margin. Dotted lines indicate a score of 1: *ex post* system = *ex ante* system.



**Figure 2.** Multi-criteria assessment of the four innovative cropping systems. Comparisons between the three constraint-limited innovative systems and the PHEP system (the gray area corresponds to the constrained system/PHEP system ratio). Cropping systems: PHEP (productive with high environmental performance), L-GHG (low greenhouse gas emissions), L-EN (low energy use) and No-Pest (no pesticide use). Cg balance: carbon gas balance. Tot GHG: total greenhouse gas emissions. D GHG: direct greenhouse gas emissions. Ind GHG: indirect greenhouse gas emissions. Tot En: total energy consumption. D En: direct energy consumption. Ind En: indirect energy consumption. En op: energy output. EN eff: energy efficiency. TFI: treatment frequency index. TFIH: TFI for herbicides. TFIothers: TFI for all pesticides other than herbicides. G margin: gross margin. Dotted lines indicate a score of 1: constrained system = PHEP system.





**Figure 3.** Multi-criteria assessment of the four innovative cropping systems. Comparisons between the four innovative systems and the current system in the Ile-de-France region (the gray area corresponds to the innovative system/current Ile-de-France system ratio. Cropping systems: A: PHEP (productive with high environmental performance), B: L-GHG (low greenhouse gas emissions), C: L-EN (low energy use) and D: No-Pest (no pesticide use). Cg balance: carbon gas balance. Tot GHG: total greenhouse gas emissions. D GHG: direct greenhouse gas emissions. Ind GHG: indirect greenhouse gas emissions. Tot En: total energy consumption. D En: direct energy consumption. Ind En: indirect energy consumption. EN op: energy output. TFI: treatment frequency index. TFIH: TFI for herbicides. TFIothers: TFI for all pesticides other than herbicides. G margin: gross margin. Dotted lines indicate a score of 1: innovative system = current Ile-de-France system.





Table 1. Main crop management strategies used in the four cropping systems to meet constraints and environmental objectives. Cropping systems: PHEP (productive with high environmental performance), L-GHG (low greenhouse gas emissions), L-EN (low energy use), No-Pest (no pesticide use). In bold: constraints set for the innovative systems.

| Cropping | Constraints and    | Specific agronomic practices managed in the innovative systems to reach the combination of targets       | Common agronomic practices       |
|----------|--------------------|--|----------------------------------|
| systems  | objectives of the  | specific to the system   | managed in the four systems      |
|          | systems            |  |                                  |
| PHEP     | - No constraint    | Earlier sowing of oilseed rape to maximize competition against weeds, and use of stale seed-bed          | Lengthening of the crop          |
|          | - Environmental    | techniques to increase weed emergence before sowing and to reduce herbicide use                          | sequence (five or six years)     |
|          | objectives         | Shallow plowing to maintain beneficial insects such carbides (slug predators) and to reduce molluscicide | and sowing a wide range of       |
|          | - High yield       | use  | crops to enhance crop            |
|          |                    | One plowing permitted during the five-year crop sequence, to reduce energy consumption                   | diversity and to reduce the      |
|          |                    | Target yield: similar to that of low-input cropping systems in the Ile-de France region                  | impact of pests on crops         |
| No-Pest  | - No pesticide use | Alternate sowing of host and non-host plants or of spring and winter crops, to decrease pest pressure    | Sowing of highly resistant       |
|          | - Environmental    | Sowing winter wheat later to reduce insect impact during autumn (aphids)                                 | varieties or variety mixtures to |
|          | objectives         | Sowing species with rapid shoot growth, such as hemp and triticale, to increase competitiveness          | reduce the impact of diseases    |
|          | - High yield       | Using stale seed-bed techniques to increase weed emergence before sowing                                 | on crops                         |
|          |                    | Shallow plowing to maintain beneficial insects such carbides (slug predators)                            | Lower sowing density and         |
|          |                    | Use of Trichogramma parasitoid wasps against Ostrinia nubilalis on maize                                 | levels of N fertilization to     |
|          |                    | Mechanical weeding   | decrease shoot biomass and       |
|          |                    | Lowering target yield and levels of N fertilizer to decrease pest impact                                 | disease developments             |
|          |                    | Target yield: lower than for the PHEP system, higher than those achieved in organic systems because      | Sowing of a legume to reduce     |
|          |                    | chemical fertilizers were allowed  | N fertilization needs (i.e. to   |

| L-EN  | - To halve energy      | Prohibition of plowing and use of a direct drilling system to reduce direct energy consumption            | reduce indirect energy           |
|-------|------------------------|---|----------------------------------|
|       | consumption relative   | Inclusion of legumes and high N use efficiency species in the crop sequence to reduce N fertilization     | consumption)                     |
|       | to the PHEP system     | requirements (i.e. indirect energy consumption)   | Sowing catch crops before        |
|       | - Environmental        | Target yield: 20% lower than for the PHEP system, to reduce N fertilization ( <i>i.e.</i> indirect energy | spring crops, oilseed rape after |
|       | objectives             | consumption)  | legumes and prohibition of N     |
|       | - High yield           |   | fertilization during the autumn  |
| L-GHG | - To halve greenhouse  | Sowing of many cereals and maintenance of continuous soil cover (with a cover crop), to generate large    | and winter, to decrease          |
|       | gas emissions relative | amounts of residues to increase soil organic matter content   | nitrogen leaching during these   |
|       | to the PHEP system     | Prohibition of plowing and use of a direct drilling system to reduce carbon mineralization                | seasons.                         |
|       | - Environmental        | Sowing of legumes to reduce N fertilization ( <i>i.e.</i> N <sub>2</sub> O emissions)                     | Non-removal of crop residues,    |
|       | objectives             | Systematic sowing of cover crop to reduce NO3 <sup>-</sup> availability and N2O emissions                 | to stabilize soil organic matter |
|       | - High yield           | Sowing of species with taproots to reduce soil compaction   | levels                           |
|       |                        | Target yield: similar to that of the PHEP system  |                                  |

**Table 2.** Mean annual treatment frequency indices (TFI: all pesticides; TFIH: herbicides; TFI others: pesticides than herbicides) for the four cropping systems, calculated at the crop sequence scale. Cropping systems: PHEP (productive with high environmental performance), L-GHG (low greenhouse gas emissions), L-EN (low energy use), No-Pest (no pesticide use). The values in brackets are the confidence intervals (p < 0.05) for the three replicates. For the No-Pest system, all values are zero.

| Performances                                      | PHEP        | L-GHG         | L-EN          | No-Pest |
|---|-------------|---------------|---------------|---------|
| TFI (ha <sup>-1</sup> .year <sup>-1</sup> )       | [1.73;2.15] | [2.56; 2.81]  | [1.83 ; 2.93] | 0.00    |
| TFIH (ha <sup>-1</sup> .year <sup>-1</sup> )      | [1.06;1.41] | [1.49 ; 1.85] | [1.71 ; 2.36] | 0.00    |
| TFIothers (ha <sup>-1</sup> .year <sup>-1</sup> ) | [0.47;0.94] | [0.78; 1.25]  | [0.03 ; 0.67] | 0.00    |

**Table 3.** Main agronomic practices of the four cropping systems, for each replicate and each crop. Bold characters correspond to the crops sown in 2009, *i.e.* the first crop of the crop sequence sown in each replicate. Rep = replicate. Nb = number. Catch or cover crops were systematically sown before main crops. W and S are winter and spring crops, respectively. MWheat or MBarley = mixture of varieties for wheat and barley, respectively. 20ilseed rape or 2Maize = two sowings of oilseed rape or maize, respectively, due to plant emergence failure. Flax(W)+Flax(S) = sowing of spring flax after winter flax was destroyed by frost. None = no cover or catch crop. Bmustard and Wmustard correspond to brown and white mustard, respectively. Wclover = white clover. IntWheat = intercropped winter wheat and white clover. Cropping systems: PHEP (productive with high environmental performance), L-GHG (low greenhouse gas emissions), L-EN (low energy use), No-Pest (no pesticide use).

|  |                                | Nb of         |             |                        | Species of cover/catch    | Nb of herbicides   |            |
|--|--------------------------------|---------------|-------------|------------------------|---------------------------|--|------------|
| Cropping systems                                       |                                | plowings      | Sowing      | Mineral                | crops before each crop or | (H), fungicides (F),   | Nb of      |
| Replication  |                                | (P) and       | density (kg | nitrogen rate          | associated crop in        | insecticides (I), and  | mechanical |
| Crop sequence  | Crop                           | tillage(T)    | seed.ha-1)  | (kg.ha <sup>-1</sup> ) | intercropping             | molluscicides (M)  | weedings   |
| PHEP   | Barley(S)                      | 1P+1T         | 154         | 40                     | Bmustard                  | 1H   | None       |
| Rep1: Barley(S)-Faba                                   | Faba bean(W)                   | 1T            | 162         | 0                      | None                      | None   | None       |
| bean(W)-Wheat(W)-                                      | Wheat(W)                       | 3T            | 110         | 90                     | None                      | 1H   | None       |
| Rape(W)-   | Rane(W)                        | 2Т            | 2           | 115                    | None                      | 2H+1F+2I+1M  | None       |
| MWheat(W)  | MWheat(W)                      | 1T            | 99          | 90                     | None                      | 3H   | None       |
| PHFP   | Wheat(W)                       | 1T            | 122         | 40                     | None                      | 1H   | None       |
| Ren? Wheat(W)-   | Barley(S)                      | 11            | 112         | 70                     | Wmustard                  | 111<br>1H  | None       |
| Barley(S)-Faba   | Eaba bean $(W)$                | 1 P           | 212         | 0                      | None                      | 311  | None       |
| been(W) Wheet(W)                                       | Wheat(W)                       | 11<br>4T      | 100         | 70                     | None                      | 2H   | None       |
| Bana(W)  | Pana(W)                        | 41<br>2T      | 2           | 100                    | None                      | 211<br>2U+1E+1101M   | None       |
|  |                                | 21            | 2.2         | 50                     | None                      | 111, 1E, 21, 1M  | None       |
| PHEP   | $2 \text{Kape}(\mathbf{W})$    | 11            | 2+3         | 50                     | INONE                     | 1H+1F+2I+1M  | None       |
| Rep3 : 2Kape(w)-                                       | wheat(w)                       | 21            | 100         | 120                    | None                      | 2H   | None       |
| Wheat(W)-Barley(S)-                                    | Barley(S)                      | IP+IT         | 127         | 60                     | Wmustard                  | 2H   | None       |
| Faba bean(W)-  | Faba bean(W)                   | None          | 342         | 0                      | Buckwheat                 | 2H   | None       |
| Wheat(W)   | Wheat(W)                       | 21            | 104         | 0                      | None                      | 2H+2F  | None       |
| L-GHG  | Wheat(W)                       | 1T            | 122         | 40                     | None                      | 1H   | None       |
| Rep1   | Barley(W)                      | None          | 127         | 80                     | Peas                      | 1H   | None       |
| Wheat(W)-  | 2Maize                         | 1T            | 190         | 80                     | Clover+Oat                | 2H+1M  | None       |
| Barley(W)-2Maize -                                     | Triticale                      | None          | 205         | 0                      | None                      | 2H+1F  | None       |
| Triticale-Faba   | Faba bean(S)                   | None          | 220         | 0                      | Wmustard                  | 3H   | None       |
| bean(S)-2Rape(W)                                       | 2Rape(W)                       | None          | 3+4         | 50                     | None                      | 4H+2M  | None       |
| L-GHG  | 2Rape(W)                       | 1T            | 2+3         | 50                     | None                      | 1H+1F+2I+1M  | None       |
| Rep2: 2Rape(W)-  | Wheat(W)                       | None          | 137         | 80                     | Peas                      | 2H   | None       |
| Wheat(W)-  | Barlev(W)                      | None          | 184         | 80                     | Peas                      | 4H   | None       |
| Barlev(W)-Maize-                                       | Maize                          | 1T            | 190         | 110                    | Clover+Oat                | 3H+1M  | None       |
| Triticale-Faba bean(S)                                 | Triticale                      | None          | 165         | 90                     | No                        | 1H   | None       |
|  | Faha hean(S)                   | None          | 248         | 0                      | Lentil+Oat+Wmustard       | 1H+1F+1M   | None       |
| L-GHG  | Maize                          | 1T            | 190         | 130                    | Bmustard                  | 1H+1I  | None       |
| Ren3: Maize  | Triticale                      | None          | 100         | 0                      | None                      | 111  | None       |
| Triticale-2Eaba  | 2Eaba bean(S)                  | None          | 220+73      | 0                      | None                      | 3H±1I±1M   | None       |
| hean(S) Pape (W)                                       | $\mathbf{P}_{aba}(\mathbf{W})$ | None          | 0           | 40                     | None                      | $2\mathbf{H}_{\perp}1\mathbf{E}_{\perp}1\mathbf{I}_{\perp}1\mathbf{M}$ | None       |
| Wheat(W)   | Wheat(W)                       | None          | 112         | 100                    | Fenugreek                 | $4H_{\pm}2H_{\pm}1M$   | None       |
| Wheat(W)-  | MDorlay(W)                     | None          | 112         | 100                    | Puelswheet                | 40+20+1M   | None       |
| Mbarley(w)   | Mibarley(w)                    | INOILE        | 145         | 90                     | Buckwheat                 | <u> 3П</u>   | None       |
| L-EN   | $\mathbf{Flax}(\mathbf{W})$    | 11            | 33          | 0                      | None                      | IH   | None       |
| $\operatorname{Rep1}: \operatorname{Flax}(\mathbf{w})$ | Intwheat(w)                    | None          | 125         | 40                     | wclover                   | IH   | None       |
| Int Wheat(W)-Oat(W)-                                   | Oat(W)                         | None          | 117         | 0                      | Wclover                   | 3H   | None       |
| Faba bean(W)-  | Faba bean(W)                   | None          | 342         | 0                      | None                      | 4H   | None       |
| Wheat(W)   | Wheat(W)                       | None          | 126         | 30                     | None                      | 3H+1M  | None       |
| L-EN   | Oat(S)                         | 1T            | 114         | 0                      | Wclover                   | 1H   | None       |
| Rep2: Oat(S)-Faba                                      | Faba bean(W)                   | None          | 159         | 0                      | None                      | None   | None       |
| bean(W)-Wheat(W)-                                      | Wheat(W)                       | None          | 142         | 40                     | None                      | 4H   | None       |
| Flax(W)+Flax(S)-                                       | Flax(W)+Flax(S                 | None          | 35+60       | 0                      | None                      | 5H   | None       |
| IntWheat(W)  | IntWheat(W)                    | None          | 124         | 90                     | Wclover                   | 4H+1M  | None       |
| L-EN   | Faba bean(W)                   | 1T            | 131         | 0                      | None                      | None   | None       |
| Rep3: Faba   | Wheat(W)                       | None          | 125         | 0                      | None                      | 2H   | None       |
| bean(W)-Wheat(W)-                                      | Flax(W)                        | None          | 40          | 0                      | None                      | 6H   | None       |
| Flax(W)-   | IntWheat(W)                    | None          | 173         | 80                     | Wclover                   | 3H   | None       |
| IntWheat(W)-Oat(S)                                     | Oat(S)                         | None          | 150         | 0                      | Wclover                   | 2H   | None       |
| No-Pest  | Maize                          | 1P+2T         | 190         | 80                     | Bmustard                  | None   | 2          |
| Rep1: Maize-   | MWheat(W)                      | None          | 235         | 70                     | None                      | None   | 1          |
| MWheat(W)-Faba   | Faba bean(S)                   | 1P+1T         | 220         | 0                      | Wmustard+Oat              | None   | 1          |
| bean(S)-MWheat(W)-                                     | MWheat(W)                      | 3T            | 160         | Ő                      | None                      | None   | 1          |
| Hemp-Triticale   | Hemp                           | 1P            | 57          | 30                     | Vetch                     | None   | 0          |
| Themp Thueure  | Triticale                      | 1P+1T         | 141         | 30                     | None                      | None   | 1          |
|  | Faha hoon(S)                   | 1P+7T         | 180         | 0                      | None                      | None   | 0          |
| No-Dect  | MWheat(W)                      | 11 72 I<br>1T | 156         | 0                      | Barley volunteers         | None   | 1          |
| Ren2 · Faba boon(S)                                    | Hemp                           | 1р_7т         | 55          | 0                      | Clover_Mustard            | None   | 1          |
| MWheat(W) Hame   | Triticala                      | 11 TZ 1<br>2T | 160         | 40                     | Nona                      | None   | 0          |
| wiwneau(w)-Hemp-                                       | Trucale                        | 31            | 100         | 40                     | inone                     | inone  | U          |

| Triticale-Maize- | Maize        | 1P+1T | 190     | 90  | Wmustard+Lentil | None | 2 |
|------------------|--------------|-------|---------|-----|-----------------|------|---|
| MWheat(W         | MWheat(W)    | 1P    | 160     | 70  | None            | None | 1 |
| No-Pest          | Wheat(W)     | 2T    | 174     | 0   | None            | None | 0 |
| Rep3: Wheat(W)-  | Faba bean(S) | 1P+2T | 196     | 0   | Buckwheat       | None | 1 |
| Faba bean(S)-    | MWheat(W)    | 3T    | 160     | 0   | None            | None | 1 |
| MWheat(W)-Hemp-  | Hemp         | 1P+3T | 55      | 0   | Peas            | None | 0 |
| Triticale-2Maize | Triticale    | 1T    | 150-    | 0   | None            | None | 1 |
|                  | 2Maize       | 1P+3T | 220+220 | 110 | Mustard+Lentil  | None | 5 |
|                  |              |       |         |     |                 |      |   |

**Table 4.** Energy (total, direct and indirect) consumption (MJ.ha<sup>-1</sup>.year<sup>-1</sup>), energy output (MJ.ha<sup>-1</sup>.year<sup>-1</sup>) and energy use efficiency of the four cropping systems. Cropping systems: PHEP (productive with high environmental performance), L-GHG (low greenhouse gas emissions), L-EN (low energy use), No-Pest (no pesticide use). The values shown are the means and the standard deviations for the three replicates. The same letters indicate homogeneous groups according to the Tukey test, p < 0.05 (ns: not significant).

| Performance  | PHEP                      | L-GHG                 | L-EN                          | No-Pest               |
|--|---------------------------|-----------------------|-------------------------------|-----------------------|
| Total energy consumption (MJ.ha <sup>-1</sup> .year <sup>-1</sup> )    | 7755 ± 711 a              | 7459 ± 793 a          | 5201 ± 502 b                  | 7604 ± 517 a          |
| Direct energy consumption (MJ.ha <sup>-1</sup> .year <sup>-1</sup> )   | 3665 ± 223 b              | 2562 ± 235 c          | 2618 ± 171 c                  | 4417 ± 425 a          |
| Indirect energy consumption (MJ.ha <sup>-1</sup> .year <sup>-1</sup> ) | $4090 \pm 489 \text{ ab}$ | 4897 ± 568 a          | 2584 ± 479 c                  | $3187 \pm 99$ bc      |
| Ratio: Indirect energy consumption/<br>Total energy consumption        | 52.7%                     | 65.7%                 | 49.7%                         | 41.9%                 |
| Energy output (MJ.ha <sup>-1</sup> .year <sup>-1</sup> )               | 95965 ± 8397 a            | 90229 ± 5572 a        | 70997 ± 9991 b                | 103323 ± 3629 a       |
| Energy use efficiency  | $12.41 \pm 1.07$ (ns)     | $12.14 \pm 0.74$ (ns) | $13.71 \pm 2.10 \text{ (ns)}$ | $13.61 \pm 0.54$ (ns) |

**Table 5.** Annual yield (t.ha<sup>-1</sup>) values (0% humidity) from 2009 to 2014, for each crop of the four cropping systems. Results in bold characters correspond to the crops sown in 2009 (i.e. the first crop of the crop sequence sown in each replicate). W and S are winter and spring crops, respectively. Cropping systems: PHEP (productive with high environmental performance), L-GHG (low greenhouse gas emissions), L-EN (low energy use), No-Pest (no pesticide use).

#### Cropping system

Successive crops in the crop sequence

Replication

| PHEP: species                         | S barley        | W faba bean     | W wheat         | W rape          | W wheat         |                 |
|---------------------------------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| calorific value (MJ.t <sup>-1</sup> ) | 14.5            | 14.4            | 14.5            | 24.6            | 14.5            |                 |
| PHEP: target yield                    | 5.3             | 3.0             | 6.7             | 2.8             | 6.7             |                 |
| PHEP: Replicate 1                     | 6.81            | 1.80            | 8.13            | 3.49            | 7.78            |                 |
| PHEP: Replicate 2                     | 5.32            | 1.38            | 6.65            | 3.48            | 6.13            |                 |
| PHEP: Replicate 3                     | 4.78            | 1.18            | 6.73            | 3.96            | 8.17            |                 |
| PHEP: mean yield                      |                 |                 |                 |                 |                 |                 |
| value and standard                    |                 |                 |                 |                 |                 |                 |
| deviation                             | 5.64 ± 1.04     | $1.45 \pm 0.31$ | $7.17 \pm 0.81$ | $3.64 \pm 0.29$ | 7.36 ± 1.12     |                 |
| L-GHG: species                        | W wheat         | W barley        | Maize           | Triticale       | S faba bean     | W rape          |
| calorific value (MJ.t <sup>-1</sup> ) | 14.5            | 14.5            | 14.5            | 14.6            | 14.4            | 24.6            |
| L-GHG: target yield                   | 6.7             | 6.1             | 7.0             | 6.0             | 4.1             | 2.8             |
| L-GHG: Replicate 1                    | 7.40            | 6.15            | 7.64            | 5.82            | 1.95            | 0.00            |
| L-GHG: Replicate 2                    | 7.51            | 4.94            | 7.46            | 5.53            | 1.53            | 4.04            |
| L-GHG: Replicate 3                    | 7.24            | 4.96            | 5.27            | 6.99            | 0.61            | 3.26            |
| L-GHG: mean yield                     |                 |                 |                 |                 |                 |                 |
| value and standard                    |                 |                 |                 |                 |                 |                 |
| deviation                             | $7.38 \pm 0.15$ | $5.35 \pm 0.72$ | $6.79 \pm 1.30$ | $6.11 \pm 0.79$ | $1.36 \pm 0.71$ | $2.43 \pm 2.14$ |
| L-EN: species                         | S oat           | W faba bean     | W wheat         | W flax          | W wheat         |                 |
| calorific value (MJ.t <sup>-1</sup> ) | 15.8            | 14.4            | 14.5            | 21.2            | 14.5            |                 |
| L-EN: target yield                    | 3.2             | 3.0             | 5.4             | 1.6             | 5.4             |                 |
| L-EN: Replicate 1                     | 3.69            | 2.88            | 6.07            | 1.72            | 6.00            |                 |
| L-EN: Replicate 2                     | 6.06            | 2.28            | 6.33            | 0.86            | 6.51            |                 |
| L-EN: Replicate 3                     | 3.46            | 4.16            | 6.98            | 1.29            | 0.75            |                 |

#### L-EN: mean yield

#### value and standard

| deviation                             | $4.40 \pm 1.45$ | $3.11 \pm 0.97$ | $6.46 \pm 0.51$ | $1.29 \pm 0.40$ | $4.42 \pm 3.16$ |           |   |
|---------------------------------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------|---|
| No-Pest: species                      | Maize           | W wheat         | S faba bean     | W wheat         | Hemp            | Triticale | - |
| calorific value (MJ.t <sup>-1</sup> ) | 14.5            | 14.5            | 14.4            | 14.5            | 16.5            | 14.6      |   |
| No-Pest: target yield                 | 5.6             | 4.7             | 3.1             | 4.7             | 8.0             | 4.3       |   |
| No-Pest: Replicate 1                  | 3.81            | 7.99            | 0.28            | 6.25            | 13.10           | 3.38      |   |
| No-Pest: Replicate 2                  | 5.24            | 6.09            | 4.19            | 6.82            | 8.20            | 4.97      |   |
| No-Pest: Replicate 3                  | 5.83            | 5.07            | 2.76            | 6.14            | 12.40           | 5.03      |   |
| No-Pest: mean yield                   |                 |                 |                 |                 |                 |           |   |
| value and standard                    |                 |                 |                 |                 |                 |           |   |

| deviation | 496 +  |
|-----------|--------|
| deviation | 4.90 1 |

1.03  $6.38 \pm 1.47$  2.41  $\pm 1.98$  6.40  $\pm 0.36$  11.23  $\pm 2.65$  4.46  $\pm 0.92$ 

**Table 6.** Carbon balance, C sequestration and greenhouse gas emissions (total, direct and indirect) of the four cropping systems, calculated over a 50-year period (C content of the soil = 13 g.kg<sup>-1</sup> dry matter). Cropping systems: PHEP (productive with high environmental performance), L-GHG (low greenhouse gas emissions), L-EN (low energy use), No-Pest (no pesticide use). The values shown are the means and standard deviations for the three replicates. Identical letters indicate homogeneous groups according to the Tukey test, p < 0.05 (ns: not significant).

| Performances   | PHEP                      | L-GHG                    | L-EN                     | No-Pest                  |
|--|---------------------------|--------------------------|--------------------------|--------------------------|
| Carbon balance (kgCO <sub>2</sub> eq.ha <sup>-1</sup> .year <sup>-1</sup> )                    | $1188 \pm 270 \text{ ns}$ | $1202 \pm 86 \text{ ns}$ | $1072 \pm 29 \text{ ns}$ | $1404 \pm 90 \text{ ns}$ |
| C sequestration (kgCO <sub>2</sub> eq.ha <sup>-1</sup> .year <sup>-1</sup> )                   | -117 ± 150 a              | -149 ± 117 a             | -518 ± 92 b              | -560 ± 49 b              |
| Total greenhouse gas emissions (kgCO <sub>2</sub> eq.ha <sup>-1</sup> .year <sup>-1</sup> )    | 1071 ± 145 a              | 1052 ± 183 a             | 554 ± 107 b              | 844 ± 46 ab              |
| Direct greenhouse gas emissions (kgCO <sub>2</sub> eq.ha <sup>-1</sup> .year <sup>-1</sup> )   | 622 ± 82 a                | 541 ± 102 a              | 311 ± 40 b               | 509 ± 26 a               |
| Indirect greenhouse gas emissions (kgCO <sub>2</sub> eq.ha <sup>-1</sup> .year <sup>-1</sup> ) | 449 <u>±</u> 64 ab        | 511 ± 82 a               | $243 \pm 67 c$           | $335 \pm 20 \text{ bc}$  |

**Table 7.** Environmental performances of the various innovative cropping systems calculated with the Criter<sup>®</sup> tool, at the crop sequence scale. Cropping systems: PHEP (productive with high environmental performance), L-GHG (low greenhouse gas emissions), L-EN (low energy use), No-Pest (no pesticide use). For the NH<sub>3</sub> volatilization, N<sub>2</sub>O emissions and NO<sub>3</sub><sup>-</sup> leaching indicators, the values shown are the means and standard deviations for the three replicates. Identical letters indicate homogeneous groups according to the Tukey test, p < 0.05 (ns: not significant). The Pesticide volatilization, Pesticide leaching, and Pesticide runoff indicators, all take a value of 10 for the No-Pest system. We therefore show confidence intervals in italic brackets (p < 0.05). For the Crop diversity indicator, no standard deviations were calculated because the three replicates of each system had the same crop sequence.

| Indicators  | PHEP          | L-GHG                     | L-EN               | No-Pest            |
|---|---------------|---------------------------|--------------------|--------------------|
| Qualitative indicators                              |               |                           |                    |                    |
| NH <sub>3</sub> volatilization                      | 9.84 ± 0.03 b | 9.85 ± 0.04 b             | 9.94 ±.0.02 a      | $9.91 \pm 0.01$ ab |
| N <sub>2</sub> O emissions                          | 8.69 ± 0.16 b | $8.80 \pm 0.14$ ab        | 9.17 ± 0.06 a      | $9.10 \pm 0.13$ ab |
| Pesticide volatilization                            | [8.52 ; 9.72] | [8.22 ; 8.78]             | [8.39;9.37]        | 10.00              |
| Pesticide leaching                                  | [8.37 ; 8.43] | [8.74 ;8.78]              | [8.38;8.76]        | 10.00              |
| Pesticide runoff                                    | [8.59;9.10]   | [8.69 ;8.90]              | [8.72 ;9.02]       | 10.00              |
| Crop diversity                                      | 6.8           | 7                         | 7.8                | 7.5                |
| Quantitative indicator                              |               |                           |                    |                    |
| NO <sub>3</sub> <sup>-</sup> leaching               |               |                           |                    |                    |
| $(\text{kg NO}_3^-\text{ha}^{-1}.\text{year}^{-1})$ | 8.93 ± 2.24 a | $4.53 \pm 0.56 \text{ b}$ | $6.25 \pm 0.67$ ab | $7.83 \pm 0.80$ ab |

**Table 8.** Economic results (gross margin, gross output and total variable costs, all expressed in  $\in$ .ha<sup>-1</sup>.year<sup>-1</sup>) for the various innovative cropping systems. CAP = Common Agricultural Policy. Cropping systems: PHEP (productive with high environmental performance), L-GHG (low greenhouse gas emissions), L-EN (low energy use), No-Pest (no pesticide use). The values shown are the means and standard deviations for the three replicates. Identical letters indicate homogeneous groups according to the Tukey test, p < 0.05.

| Performances   | PHEP                | L-GHG           | L-EN           | No-Pest             |
|--|---------------------|-----------------|----------------|---------------------|
| Gross margin (€.ha <sup>-1</sup> .year <sup>-1</sup> )         | 757.0 ± 88.7 a      | 619.6 ± 77.3 ab | 606.4 ± 56.1 b | $701.4 \pm 48.4$ ab |
| Gross output (€.ha <sup>-1</sup> .year <sup>-1</sup> )         | 929.1 ± 71.6 a      | 861.9 ± 84.0 a  | 696.0 ± 74.5 b | 879.0 ± 48.9 a      |
| Total variable costs (€.ha <sup>-1</sup> .year <sup>-1</sup> ) | 497.5 $\pm$ 43.8 ab | 567.7 ± 29.7 a  | 415.0 ± 46.7 b | $503.0 \pm 54.6$ ab |
| CAP subsidies (€.ha <sup>-1</sup> .year <sup>-1</sup> )        | 325                 | 325             | 325            | 325                 |