



HAL
open science

Contribution of exudates, arbuscular mycorrhizal fungi and litter depositions to the rhizosphere priming effect induced by grassland species

Tanvir Shahzad, Claire Chenu, Patricia Genet, Sébastien Barot, Nazia Perveen, Christian Mougin, Sébastien Fontaine

► To cite this version:

Tanvir Shahzad, Claire Chenu, Patricia Genet, Sébastien Barot, Nazia Perveen, et al.. Contribution of exudates, arbuscular mycorrhizal fungi and litter depositions to the rhizosphere priming effect induced by grassland species. *Soil Biology and Biochemistry*, 2015, 80, pp.146–155. 10.1016/j.soilbio.2014.09.023 . hal-01567145

HAL Id: hal-01567145

<https://agroparistech.hal.science/hal-01567145v1>

Submitted on 3 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Contribution of exudates, arbuscular mycorrhizal fungi and litter depositions to the rhizosphere priming effect induced by grassland species

Tanvir Shahzad^{1,2}, Claire Chenu³, Patricia Genet^{4,5}, Sébastien Barot⁶, Nazia Perveen¹, Christian Mougin⁷, Sébastien Fontaine¹

¹ INRA, UR 874 Unité de Recherche sur l'Ecosystème prairial, 5, Chemin de Beaulieu 63039 Clermont Ferrand cedex 2, France.

² Department of Environmental Sciences, Government College University Faisalabad, Allama Iqbal Road, 38000 Faisalabad, Pakistan

³ AgroParisTech, UMR BioEMCo (Univ. Paris 6, Univ. Paris 12, AgroParisTech, ENS, CNRS, INRA, IRD), 78850, Thiverval-Grignon, France

⁴ UPMC Paris 6 – CNRS UMR 7618, 46 rue d'Ulm, 75230, Paris Cedex 05, France

⁵ Université Paris Diderot – Paris 7, 75205, Paris Cedex 13, France

⁶ IRD-Bioemco, UMR 7618, Ecole Normale Supérieure, 46 Rue d'Ulm, F-75230 Paris Cedex

⁷ INRA UR 251 PESSAC, Centre de Versailles-Grignon RD 10- 78026 Versailles Cedex

Keywords: SOM mineralization, Continuous ¹³C labeling, Microbial Community, Priming effect, Root exudates, PLFA, Grasslands.

Abstract

The presence of plants induces strong accelerations in soil organic matter (SOM) mineralization by stimulating soil microbial activity - a phenomenon known as the rhizosphere priming effect (RPE). The RPE could be induced by several mechanisms including root exudates, arbuscular mycorrhizal fungi (AMF) and root litter. However the contribution of each of these to rhizosphere priming is unknown due to the complexity involved in studying rhizospheric processes. In order to determine the role of each of these mechanisms, we incubated soils enclosed in nylon meshes that were permeable to exudates, or exudates & AMF or exudates, AMF and roots under three grassland plant species grown on sand. Plants were continuously labeled with ^{13}C depleted CO_2 that allowed distinguishing plant-derived CO_2 from soil-derived CO_2 . We show that root exudation was the main way by which plants induced RPE (58-96% of total RPE) followed by root litter. AMF did not contribute to rhizosphere priming under the two species that were significantly colonized by them i.e. *Poa trivialis* and *Trifolium repens*. Root exudates and root litter differed with respect to their mechanism of inducing RPE. Exudates induced RPE without increasing microbial biomass whereas root litter increased microbial biomass and raised the RPE mediating saprophytic fungi. The RPE efficiency (RPE/unit plant-C assimilated into microbes) was 3-7 times higher for exudates than for root litter. This efficiency of exudates is explained by a microbial allocation of fresh carbon to mineralization activity rather than to growth. These results suggest that root exudation is the main way by which plants stimulated mineralization of soil organic matter. Moreover, the plants through their exudates not only provide energy to soil microorganisms but also seem to control the way the energy is used in order to maximize soil organic matter mineralization and drive their own nutrient supply.

1. Introduction

Rhizosphere processes contribute almost half of the total CO₂ emissions from the terrestrial ecosystems to the atmosphere at global level (Schimel, 1995). They are also suggested to play a significant role in mediating ecosystem feedbacks to climate change through their effects on net primary productivity, organic matter decomposition, nutrient cycling and carbon (C) storage (Grayston et al., 1997; Kaiser et al., 2010; Cheng et al., 2013). This has led to studies to better understand the magnitude, controls and direction of rhizosphere processes over soil organic matter (SOM) dynamics (Dijkstra et al., 2006a,b; Dijkstra and Cheng, 2007; Cheng et al., 2013; Dijkstra et al., 2013; Drake et al., 2013). Despite this increased interest, there are still many uncertainties regarding which processes strongly accelerate SOM turnover in the rhizosphere thereby making the predictions of soils' feedback to climate change contradictory (Davidson and Janssens, 2006).

Rhizodeposition, i.e. the release of root exudates, mucilage and sloughed-off root border cells, constitute significant inputs of labile carbon into soil (Paterson, 2003; Nguyen, 2009). Overall, up to 20% of the net C fixed by plants is released into soil in the form of rhizodeposition during vegetative period (Hütsch et al., 2002). The main component of rhizodeposition is root exudates through which 10-100 times more carbon is released than mucilage and sloughed-off border cells (Nguyen, 2009). The exudates have been found to play important role in soil ecology and plant nutrition. For example, they have been found to enhance mycorrhizal fungal growth (Elias and Safir, 1987; Ratnayake et al., 1978; Tawarayaya et al., 1996) thereby helping the plant to explore larger volumes of soil in search of nutrients. Moreover, organic acids present in exudates, help solubilize the insoluble phosphorus (P) in rhizosphere (Moghimi et al., 1978; Lipton et al., 1987; Saleque and Kirk, 1995). This knowledge has even led to the development of

novel plant varieties more efficient in using soil P through genetically-engineered enhanced release of organic acids by roots (López-Bucio et al., 2000). Finally exudates have also been suggested to increase the SOM turnover by promoting the microbial activity in the rhizosphere (Hamilton and Frank, 2001; Phillips et al., 2011) although the direct evidence is still lacking (Jones et al., 2004).

The mineralization of SOM is accelerated under living plants when compared to unplanted controls due to stimulation of soil microbes (Helal and Sauerbeck, 1984; Liljeroth et al., 1994; Kuzyakov and Cheng, 2001) - a phenomenon known as the rhizosphere priming effect (RPE). It has been suggested that exudates from roots and root-associated mycorrhizae provide energy-rich substrates to rhizosphere microbes thereby enabling them to secrete extracellular enzymes responsible for the accelerated SOM decomposition (Clarholm, 1985; Hamilton and Frank, 2001). However, lab incubations testing the effect of various components of exudates on SOM mineralization have reported contradictory results. Briefly, the additions of isotopically labeled sugars, amino acids and organic acids induced positive or negative priming effects (Hamer and Marschner, 2005a, 2005b; Blagodatskaya et al., 2007; Ohm et al., 2007) or did not have any effect even if the microbial activity was stimulated (De Nobili et al. 2001; Salomé et al. 2010). This lack of effect is explained by *r*-strategist microorganisms that only use these easily degradable substrates (Fontaine et al., 2003; Blagodatskaya et al., 2007; Ohm et al., 2007). Despite their association with roots of about 80% of terrestrial plant species, the role of arbuscular mycorrhizal fungi (AMF) in the RPE remains unknown. The catabolic capabilities of root-associated AMF are generally considered low compared to those of soil decomposers (Read and Perez-Moreno, 2003; Talbot et al., 2008), suggesting they probably play a minor role in rhizosphere priming. However, they have been found to participate in the degradation of plant

litter (Hodge et al., 2001; Leigh et al., 2009; Cheng et al., 2012), suggesting that their catabolic capability has been underestimated. Finally, the RPE induced by plants may arise from the supply of litter to soil decomposers since this type of organic matter systematically induces the priming effect in incubation studies (Nottingham et al. 2009; Pascault et al. 2013). Knowledge of the contribution of exudates, AMF and root litter depositions to RPE is fundamental to predicting plant effects on soil C cycling under changing climates.

Plant species differ in terms of total labile C inputs through rhizodeposition into the soil thereby inducing varying stimulation of soil microorganisms and RPE (Dijkstra et al., 2006a,b). Moreover, a specific plant species can shape a specific structure of microbial community (Grayston et al., 1997; Germida et al., 1998; Broeckling et al., 2008) by controlling the quality and quantity of rhizodeposition into the rhizosphere (Grayston et al., 1997; Broeckling et al., 2008). Therefore it is important to study the mechanisms of rhizosphere priming under different plant species and linking the variation in RPE with the soil microbial community structure shaped by a certain plant species.

The aim of this study was to determine the role of exudates and AMF in rhizosphere priming, the relative importance of exudates, AMF and roots in determining the RPE and mechanisms by which each of these induce changes in the RPE. The effect of exudates on SOM dynamics was disentangled from that of mycorrhizae and root-litter deposition by using meshes of different pore sizes under monocultures of three grassland species namely *Lolium perenne* (Lp), *Poa trivialis* (Pt) and *Trifolium repens* (Tr). Continuous ^{13}C labeling of plants was used to distinguish soil-derived (Rs) and plant-derived respiration (Rp). The rhizosphere priming effect was calculated as the difference between Rs from planted soils and from control bare soil. We

hypothesized that exudates and root litter would induce strong priming effects by favoring the growth of microbes and AMF would not have any positive effect on RPE.

2. Materials and Methods

2.1. Soil sampling and conditioning

The soil was sampled from an upland grassland located in the environmental research observatory (ORE) established by French National Institute for Agricultural Research (INRA) in central France in 2003 (Theix, 45° 43'N, 03°01'E). The soil is a drained Cambisol developed from a granitic rock. The soil was taken from 10-40 cm soil profile. The upper 10 cm that is rich in fresh C was removed given that respiration of this pre-existing unlabeled fresh C cannot be separated from that of recalcitrant SOM. Moreover the presence of plants can modify the decomposition of fresh C (Personeni and Loiseau, 2004). The soil properties were: pH 6.3 ± 0.23 , clay 21 ± 2.1 %, soil organic carbon (SOC) 17 ± 0.28 g kg⁻¹ soil and SOC $\delta^{13}\text{C}$ -26.4 ± 0.02 ‰. This soil was enclosed in small PVC cylinders (height 1.5 cm, diameter 5 cm) whose sides were sealed by the mesh of three different pore sizes. The pore size 0.45 μm was meant to only allow the entry of exudates excluding the mycorrhizae and roots in a living rhizosphere (exudates treatment). Whereas, the pore sizes 30 and 1000 μm would also permit the entry of mycorrhizae and roots (mycorrhizae & roots) respectively. From now on, these soil-containing cylinders will be called soil compartments.

2.2. Establishment of monocultures

Three grassland species i.e. *Lolium perenne* (Lp), *Poa trivialis* (Pt) and *Trifolium repens* (Tr), that were previously found to induce variable priming effects (Shahzad et al., 2012), were selected for this experiment. PVC pots (20 cm high, 7.8 cm internal diameter) were filled with sand (pH 7) whereas four soil compartments each containing about 15g of equivalent dry soil were placed vertically in each pot (Fig. 1). In August 2010, twelve pots were sown by each of the

three plant species representing four replicates for each pore size of mesh (0.45, 30, 1000 μm). Four pots containing soil compartments (1000 μm mesh) were kept bare as control soil. An automated drip irrigation method was used for water supply and all pots were water-saturated whenever the soil moisture decreased to $75 \pm 5 \%$ of the soil field capacity. The near-field capacity conditions were maintained to avoid the artifact of drying-rewetting cycles that may increase CO_2 release from soil (Schimel et al., 2011).

2.3. Labeling system

The detailed description of the labeling system has previously been given in Shahzad et al., 2012 (Materials & Methods and Supplementary material). Briefly, a compressor injected ambient air into a molecular sieve to remove CO_2 , H_2O and all particles. The CO_2 -free air was then mixed with ^{13}C - depleted CO_2 of fossil fuel origin ($\delta^{13}\text{C}$: $-38.55 \pm 0.07 \text{‰}$) and passed through a humidifier. The water flow in humidifier was regulated such that the relative humidity of the labeled air reaching the chamber, containing all the microcosms (soil, plants) used in this experiment, was around 50-60%. The air injected in the chamber had a CO_2 concentration of 400 ± 20 ppm. The chamber consisted of an iron box with a plexiglass screen mounted on it (dimensions: $350 \times 140 \times 140$ cm). All planted and bare pots were placed in the chamber and continuously ventilated with air produced by labeling system. The volume of the air in the chamber was renewed twice a minute. This quick renewal of the air was used to maintain constant concentration (400 ppm) and isotopic composition (-38.55‰) of CO_2 in the chamber. The ventilation of air did also not let the temperature difference exceed $2 \text{ }^\circ\text{C}$ between inside and outside of chamber. All the planted and bare pots were placed in a continuous ^{13}C - CO_2 labeling system throughout the experiment.

2.4. *Mycorrhizal colonization*

Since the plants were grown in sand which is an unfavorable medium for the spontaneous development of mycorrhizal colonization of roots, an inoculum of endomycorrhizae was applied to plant pots. Viable fungal spores were extracted in 1% CaCl₂ on a filter paper using fifty grams of fresh soil, taken from the same field from where experimental soil was taken. The number of viable fungal spores was found sufficient during trials of extractions i.e. at least 200 per filter paper. The filters containing spores were gently washed with distilled water at the base of the one week-old plants. It should be noted that all the plants irrespective of mesh treatments were inoculated. Murashige and Skoog (1962) nutritive solution (MS0, without sucrose) was applied thrice during the experiment: first application was done without nitrogen salts one day after the inoculation. Briefly 100 ml of salt concentrations namely calcium chloride (400 mg/L), magnesium sulphate (370 mg/L), Potassium phosphate (170 mg/L) and Sodium molybdate (0.25 mg/L) were added to each pot. The following applications were done two and five days after inoculation and included the nitrogen concentrations i.e. Ammonium nitrate (1650 mg/L) and Potassium nitrate (1900 mg/L). The nutritive solutions were applied to all the pots irrespective of mesh treatments.

Mycorrhizal colonization was measured using the method of McGonigle et al., (1990). Briefly, roots of 5 plants at the end of experiment were cut into smaller pieces of about 1 cm each, cleared in 10% KOH during 24 h, rinsed with distilled water, acidified with 1% HCl during 15 minutes, stained overnight with 0.1% Trypan blue at ambient temperature, rinsed with distilled water and destained in 50% glycerol. About 200 fragments of one sample were mounted on microscopic slides and examined with intersection method at × 200 magnification. The

parameters measured were arbuscular colonization, vesicular abundance, hyphal colonization and the root fragments non-colonized by mycorrhizae.

2.5. Respiration measurements

The evening before each respiration measurement, the pots were irrigated to field capacity thus ensuring that all the respiration measurements were done in similar soil moisture conditions. The next day pots were taken out of the chamber and sealed in air-tight PVC chambers (height 100 cm, diameter 15 cm) for 24 h. Absence of light stopped photosynthesis consequently stopping the plant absorption of soil-respired CO₂. The CO₂ released by soil plant system was trapped in soda lime trap of 100 ml of 1M NaOH that was placed in the respiration chambers. By conducting additional measurements in the respiration chambers using chromatography, we found that the soda lime trap fixed more than 99% of the CO₂ released over 24 hours by soil-plant system. Total C trapped in NaOH was measured using total inorganic C analyzer. Carbonates in NaOH of a subsample were precipitated using an excess BaCl₂ and filtered. The ¹³C abundance of carbonates (trapped CO₂) was measured with an elemental analyzer coupled to an Isotope-ratio mass spectrometer (IRMS). The soil-derived CO₂ (R_s , mg C-CO₂ kg⁻¹ dry soil day⁻¹) was separated from plant-derived CO₂ (R_p , mg C-CO₂ kg⁻¹ dry soil day⁻¹) using mass balance equations:

$$R_t = R_s + R_p \quad \text{Eq. 2.5.1}$$

$$R_t \times A_{tr}^{13} = R_s \times A_s^{13} + R_p \times A_p^{13} \quad \text{Eq. 2.5.2}$$

Where R_t was the total CO₂ emitted by the plant-soil system, R_s was the soil-derived CO₂ released as result of microbial mineralization of SOM, R_p was the CO₂ coming from plant,

mycorrhizae and microbial respiration of rhizodeposits and plant litter, A_s^{13} was the ^{13}C abundance (% atom) of soil-derived carbon, A_p^{13} was the ^{13}C abundance of respective plant root C and A_{tr}^{13} was the ^{13}C abundance of total CO_2 emitted from the plant-soil system. Equations 2.5.1 and 2.5.2 were resolved to calculate soil-derived $\text{CO}_2\text{-C}$ from total CO_2 emitted from soil-plant system as follows:

$$R_s = R_t \times (A_{tr}^{13} - A_p^{13}) / (A_s^{13} - A_p^{13}) \quad \text{Eq. 2.5.3}$$

The rhizosphere priming effect (RPE, $\text{mg CO}_2\text{-C kg}^{-1}$ dry soil day^{-1}) induced by the plants was calculated as:

$$RPE = (R_s, \text{planted soil}) - (R_s, \text{control soil}) \quad \text{Eq. 2.5.4}$$

Where R_s , control soil ($\text{mg C-CO}_2 \text{ kg}^{-1}$ dry soil day^{-1}) was CO_2 emitted by bare control soil.

2.6. Plant & soil analyses

Fifty one days after sowing, pots were destructively sampled for plant and soil analysis. Soil from the four compartments of each pot was taken out and mixed homogenously. Plant roots were washed to remove the sand and the soil (for $>1000 \mu\text{m}$ mesh) attached to them. Roots and shoots were then oven dried for 48 h at 60°C and finely ground. The dried plant material was analyzed on an elemental analyzer coupled to an isotope ratio mass spectrometer (IRMS) for total C and N content and ^{13}C abundance. Microbial biomass was measured using a modified version (Fontaine et al., 2011) of the fumigation-extraction method proposed by Vance et al., (1987). Briefly 5 g of soil was extracted with 20 mL of 30mM K_2SO_4 after 1 hour shaking. Another 5 g of soil sample was fumigated with alcohol free chloroform under vacuum conditions

in a glass desiccator for 24 hours. Chloroform was removed from the soil by ventilation and soils were extracted with 20 mL of 30mM K₂SO₄. The extracts were filtered (0.45 μm) and then lyophilized. The recovered crystals were analyzed for C content and δ¹³C. Total microbial biomass (MB_t) was calculated as:

$$MB_t = \frac{1}{k}(C_f - C_{nf}) \quad \text{Eq. 2.6.1}$$

Where C_f and C_{nf} were the carbon content of crystals obtained from extraction of fumigated and non-fumigated soil samples, respectively, and k is extraction yield of microbial biomass ($k = 16\%$, Fontaine et al., 2004b). The soil-derived (MB_s) and plant-derived (MB_p) microbial biomasses were determined as:

$$MB_t = MB_p + MB_s \quad \text{Eq. 2.6.2}$$

$$MB_t \times A_{t\text{bm}}^{13} = MB_p \times A_p^{13} + MB_s \times A_s^{13} \quad \text{Eq. 2.6.3}$$

Where $A_{t\text{bm}}^{13}$ was the ¹³C abundance of total microbial biomass. The $A_{t\text{bm}}^{13}$ was calculated as:

$$A_{t\text{bm}}^{13} = (C_f \times A_f^{13} - C_{nf} \times A_{nf}^{13}) / (C_f - C_{nf}) \quad \text{Eq. 2.6.4}$$

Where A_f^{13} and A_{nf}^{13} were the ¹³C abundances of C_f and C_{nf} respectively.

In order to determine if root exudates really penetrated in soil compartments, the amount of plant-derived C incorporated in soil was quantified for the three pore sizes (0.45, 30, 1000 μm) at the end of the experiment. The soil organic C content and its δ¹³C were determined with an elemental analyzer coupled to an IRMS. By adopting the same approach used for CO₂ and

microbial biomass, the amount of plant-derived C incorporated in soil C (C_p) was calculated as follows:

$$C_p = C_t \times (A_t^{13} - A_s^{13}) / (A_p^{13} - A_s^{13}) \quad \text{Eq. 2.6.5}$$

Where C_t was the total soil carbon content and A_t^{13} its ^{13}C abundance.

2.7. PLFA measurements

A soil sample of 2 g was freeze-dried and ground for each replicate after remaining plant materials were picked out. Phospholipids fatty acids (PLFA) were extracted using a modified method of Bligh and Dyer (1959) (Frostegård et al., 1991). Briefly, PLFA were extracted in a single-phase mixture of chloroform methanol: citrate buffer (1:2:0.8, v:v:v, pH 4.0) shaken at 400rpm for 1h. Phase splitting was done by adding equal volume of chloroform and citrate buffer. The organic phase was then submitted to a solid phase extraction on silica gel extraction cartridges (Discovery® DSC-Si SPE Tube bed wt. 500mg, volume 3mL from Supelco). Neutral lipids, glycolipids and PLFA were eluted by chloroform, acetone and methanol respectively. Methyl nonadecanoate (fatty acid methyl ester 19:0) was added as an internal standard and PLFA were trans-methylated under mild alkaline conditions to yield fatty acid methyl esters (FAMES), (Dowling et al., 1986). FAMES were then analyzed by GC/MS (4000 GC/MS, Varian) in split-less mode (1 mL, injector temperature: 250 °C) equipped with a BPX70 column (60m, 0.25 mm i.d., 0.25 mm df., SGE), and helium as a carrier gas. The temperature program was 50 °C for 5 min, raised to 165 °C at 15 C/min, followed by increases of 2 C/min up to 225 °C. This temperature was held for 15 min. To identify the FAMES, the retention times and mass spectra were compared with those obtained from standards (Bacterial Acid Methyl Ester Mix from Supelco and 11 Hexadecenoic acid (92 % cis. 8% trans) from Matreva). The PLFAs i15:0, a15:0,

i16:0, and i17:0 were designated as derived from gram positive bacteria while 17:0cy, 19:0cy & 16:1ω9c were designated as derived from gram negative bacteria (Frostegård and Bååth, 1996; Zelles, 1997). The PLFAs 18:1ω9c, 18:2ω9t and 18:2ω6c were designated as representatives of saprophytic fungi (Frostegård and Bååth, 1996; Zelles, 1997). The PLFA 16:1ω5c are often considered to represent the arbuscular mycorrhizal fungi (Olsson et al., 1995) although they can also be found in bacteria (Nichols et al., 1986). For this study this biomarker is being considered a representative of arbuscular mycorrhizal fungi (AMF) albeit with caution as suggested recently by Frostegård et al., (2011).

2.8. Statistical analyses

A two way analysis of variance (ANOVA) (95% CI) was used to determine the significant effect of plant species and mesh treatment on soil organic matter mineralization i.e. soil-derived CO₂-C and rhizosphere priming effect. The relationship between plant-derived CO₂-C or plant biomass and the rhizosphere priming effect across the plant species and mesh treatments was assessed by simple regression analysis. The effect of mesh treatment on total soil microbial biomass, soil-derived microbial biomass, plant-derived microbial biomass and different microbial groups was determined using one-way ANOVA. One-way ANOVA was also used to determine the significant effect of mesh treatment on RPE efficiency i.e. RPE induced per unit of plant-derived C assimilated in microbial biomass. All statistical analyses were performed with *Statgraphics Plus* (Manugistics, USA).

3. Results

All the three grassland species induced strong rhizosphere priming effects across all the mesh treatments ($P < 0.05$, Fig. 2). The SOM mineralization (R_s) in planted soils remained consistently higher throughout the incubation, representing between 118 % and 640 % of respiration observed in bare soil. The three plant species significantly varied in terms of the RPE induced ($P < 0.05$, Fig. 2), with the highest RPE induced by *Trifolium repens* (Tr), followed by *Lolium perenne* (Lp) and *Poa trivialis* (Pt) respectively. A strong correlation was found between the RPE and plant-derived $\text{CO}_2\text{-C}$ (R_p , $r^2 = 0.79$) or total plant biomass across all the species ($r^2 = 0.59$, Fig. 3).

Initially the mesh pore size (0.45, 30 or 1000 μm) had no effect on RPE (Fig. 2) indicating that all the RPE across the three plant species was induced by the root exudates. The roots might not have yet grown enough to reach in soil compartments. However during later stages of the experiment, the presence of roots (1000 μm mesh) in soil compartments significantly increased the total RPE when compared to that due to exudates alone (0.45 μm mesh) ($P < 0.05$, Fig. 2). The contribution of roots in induced RPE (% of the total) was 35-42 % for Lp, 20 to 28 % for Pt and 26% for Tr. The 30 μm mesh treatment, in which AMF could pass through linking roots and soil compartments, showed no effect on SOM mineralization in addition to that by exudates under all the plant species throughout the experiment ($P > 0.05$ Fig. 2).

Significant amounts of plant-derived C (^{13}C labeled C) were added to total soil organic carbon in mesh treatments 0.45 μm and 30 μm under all the three plant species. The amount of plant C incorporated in soil organic carbon in mesh treatments 0.45 μm and 30 μm was 153 ± 19

and 150 ± 8 , 147 ± 14 and 159 ± 27 and 169 ± 27 and 163 ± 22 mg C kg⁻¹ soil for Lp, Pt and Tr respectively. However the presence of roots induced incorporation of highest amounts of plant-derived C in soil organic C in the presence of roots (Fig. S2) that were 1892 ± 256 , 681 ± 120 and 1463 ± 229 mg C kg⁻¹ soil for Lp, Pt and Tr respectively.

The three plant species showed varying degree of arbuscular mycorrhizal root colonization (%) which also varied with mesh pore size (Table S1, Supplementary Material). No colonization was found for Lp for all mesh pore sizes and Pt for 0.45 μ m mesh. The Pt showed a root colonization of 4.56 % (Coefficient of variation, CV = 0.32 %) and 6.68 % (CV = 0.46 %) for 30 μ m and 1000 μ m mesh pore size respectively. The overall mycorrhizal root colonization was higher in Tr than the other plant species with percentage colonization of 8.55 % (CV = 0.85 %), 16.98 % (CV = 0.46 %) and 17.54 % (CV = 0.67) for 0.45 μ m, 30 μ m and 1000 μ m mesh pore sizes respectively.

Total microbial biomass (MB_{tot}) remained unchanged under 0.45 μ m and 30 μ m mesh treatments under all plant species ($P > 0.05$) except Pt where MB_{tot} increased by 27.9 (± 24.5) % in the presence of exudates i.e. 0.45 μ m mesh ($P < 0.05$, Fig. 4). In contrast, the presence of roots in soil compartments (1000 μ m mesh) significantly increased the MB_{tot} under all species ($P < 0.05$). The MB_{tot} was almost doubled in the presence of roots of Lp and Tr. The soil C-derived microbial biomass (MB_{soc}), remained unchanged under 0.45 μ m and 30 μ m mesh treatments under all plant species when compared to control soils ($P > 0.05$) except Pt where it significantly increased in the presence of exudates ($P < 0.05$). However, the presence of roots in soil compartments significantly increased MB_{soc} in comparison to control soils under all the plant species ($P < 0.05$) except Pt where it remained unchanged ($P > 0.05$). The assimilation of plant carbon (¹³C labeled C) i.e. plant-derived microbial biomass (MB_{plant}) was significant in all mesh

treatments under all the plant species ($P < 0.05$, Fig. 4). In mesh treatments 0.45 μm and 30 μm , 23 \pm 1.6 and 45.1 \pm 9.2, 21.9 \pm 2.6 and 32.1 \pm 7.6 and 25.7 \pm 3.4 and 28.2 \pm 8.5 mg plant-derived C kg^{-1} soil was assimilated by microbial biomass under Lp, Pt and Tr respectively. The plant-derived microbial biomass was significantly higher in 30 μm than 0.45 μm mesh treatment in Lp and Pt. However the presence of roots (1000 μm mesh) increased MB_{plant} by 2 (Pt) to 9 times (Tr) when compared to 0.45 μm and 30 μm mesh treatments ($P < 0.05$).

The plant C derived microbial biomass (MB_{plant}) was used to determine RPE efficiency i.e. the amount of RPE induced per unit of plant-derived C assimilated into microbial biomass. The RPE efficiency was highest in the presence of exudates or root-associated AMF while it was lowest in the presence of roots for all the species (Fig. 5). For example, root exudates were 3 (Pt), 6 (Lp) or 7 (Tr) times more efficient than the roots in inducing the RPE. Under Lp, significantly lower RPE efficiency was found in 30 μm mesh treatment than that in the presence of exudates (0.45 μm mesh) but it was still higher than that in the presence of roots.

The AMF biomarker (PLFA 16:1 ω 5c) in the soils showed no change under 0.45 μm (exudates) and 30 μm (permeable to root colonizing mycorrhizae) mesh treatments under all the plant species ($P > 0.05$). The presence of roots (1000 μm mesh) significantly increased the concentration of AMF in soil but only under Tr (Fig. 6). The concentrations of all saprophytic microbial groups remained unchanged in 0.45 μm and 30 μm mesh treatments under all the plant species ($P > 0.05$). However, the presence of roots increased their concentrations by large amounts across all the species ($P < 0.05$). The response of Gram negative bacteria to various types of plant C deposition was inconsistent under the three plant species (Fig. 6). The concentrations of Gram negative bacteria, when compared to that in control soils, were significantly increased in the presence of exudates under Lp and Tr but it remained unchanged

under Pt. However, their concentrations remained unchanged for 30 μm mesh. Moreover, the presence of roots stimulated concentrations of Gram positive bacteria only under Pt and Tr but not Lp.

4. Discussion

The three perennial plant species induced substantial over-production of unlabeled CO₂. The extent of this rhizosphere priming is similar to other studies (Cheng, 2009; Dijkstra and Cheng, 2007; Zhu and Cheng, 2011). The RPE can have two origins: i/ an increase in SOM mineralization and ii/ an acceleration of microbial turnover (Bingeman et al., 1953; Dalenberg and Jager, 1989, 1981). Indeed, the supply of labeled C may activate dormant microbes which renew their metabolites and release unlabeled microbial C as CO₂. The over-production of unlabeled CO₂ in this case is often called apparent RPE because it comes from an acceleration of microbial turnover and not from mineralization of SOM (Bingeman et al., 1953; Dalenberg and Jager, 1989, 1981). The occurrence of apparent RPE can be detected by measuring the amount of unlabeled C in microbial biomass of control and C-amended soil; the apparent PE decreases the unlabeled microbial C in the C-amended soil (Wu et al., 1993; Fontaine et al., 2004b). Our results show that the presence of plants had no effect or increased the unlabeled microbial C (Fig 4), indicating that the observed RPE was mostly the result of an increase in SOM mineralization i.e. the apparent RPE is negligible compared to the real RPE.

Most of the root biomass for all the plants was concentrated around soil compartments covered by meshes (Fig. 1) suggesting that the roots were injecting carbon inside the soil compartments and benefiting from the nutrient release by soil microorganisms even in treatments where if the roots were excluded from soil (treatments 0.45 μ m and 30 μ m). Indeed, the root exudates are known to diffuse up to 12 mm from their point of release (Sauer et al., 2006) whereas the maximum distance exudates had to cover in this experiment from their entry through mesh of either side to the middle of the soil compartment was 7.5 mm (height of the PVC cylinder used was 1.5 cm) indicating that the whole soil present in a compartment was accessible

to exudates. This diffusion of exudates into soil compartments is confirmed by the significant amounts of plant-derived C present in soil organic carbon (Fig. S2) and microbial biomass (Fig. 4). Therefore the SOM mineralization and other soil attributes in 0.45 μm mesh treatment can safely be attributed to exudates.

Our findings suggest that root exudates are the most important means for a plant to induce an acceleration in SOM mineralization i.e. rhizosphere priming effect (Fig. 2). The result was common for one leguminous and two grasses although they have different growth strategies (Maire et al., 2009). The role of exudates in the stimulation of microbial enzyme activities and SOM mineralization has previously been suspected (Hamilton and Frank, 2001; Phillips et al., 2012). However, by disentangling the effect of exudates from other root-induced processes, the fact that the SOM mineralization is mainly driven by exudates has important consequences for our understanding of plant-soil interactions. The deposition of exudates is closely connected to the plant photosynthetic activity, with a transfer of photosynthates to rhizosphere microorganisms occurring in less than 24 hours (Johnson et al., 2002; Deneff et al., 2009; De Deyn et al., 2011). Given that these exudates drive the SOM mineralization and thereby the release of mineral N (Phillips et al., 2011; Dijkstra et al., 2013), the plant might finely adjust its own N supply to the potential growth offered by its environment (e.g. light, CO_2). The fine control of SOM mineralization by plants is supported by a recent study (Shahzad et al., 2012) showing that the decrease in plant photosynthesis induced by clipping resulted in a 20-56% decrease in RPE within 24 hours.

The plant biomass and plant-derived C respiration, two proxies of plant photosynthesis and exudation (Bahn et al., 2009), were positively linked with RPE across the three plant species (Fig. 2). Therefore, the interspecific differences in RPEs can be explained by the difference of

photosynthetic activity and exudation among the three plant species. The more a genotype is adapted to the environmental conditions, the more photosynthesis it carries out resulting in more C being exuded from roots thereby inducing higher RPE. These results suggest that the fertility of soils (defined here as SOM mineralization) not only depends on inherent properties of soils (i.e. SOM content) but also on plant-soil interactions allowing a fine tuning of SOM mineralization to plant demand which can vary with plant species.

Only *Trifolium repens* (Tr) among the three plant species showed mycorrhizal root-colonization (8.55 ± 0.85 %, 16.9 ± 0.5 % and 17.5 ± 0.7 % for mesh 0.45 μm , 30 μm and 1000 μm respectively, Table S1, Supplementary Material) comparable to previously reported value for this species (20 %, Medina et al., 2010). *Poa trivialis* (Pt) showed significant mycorrhizal root colonization in 30 μm (4.56 ± 0.32 %) and 1000 μm mesh (6.68 ± 0.46 %) treatments whereas no root-colonization was observed in 0.45 μm treatment. To our knowledge, there is no study that previously reported the mycorrhizal colonization in Pt. Both Pt and Tr showed significantly higher root colonization in 30 μm and 1000 μm mesh treatments compared to 0.45 μm mesh indicating that the root colonization was higher when the mycorrhizal hyphae could access the soil through 30 μm or 1000 μm mesh. The better mycorrhizal root colonization of Tr can be explained by the fact that legumes are generally better colonized by mycorrhizal fungi than grasses due to their higher need in P and thereby dependence on AMF for P acquisition (Chen et al., 2005; Eschen et al., 2013). No apparent reason was found for the complete failure of mycorrhizal colonization under Lp.

Under Tr, where root colonization was quantitatively very important, root associated AMF (30 μm mesh) did not affect RPE (Fig. 2). Similarly significant mycorrhizal colonization of Pt in the 30 μm mesh treatment did not induce any increase in the RPE compared to the 0.45

μm treatment, suggesting that AMF has no or negligible effect on SOM mineralization rate. It must be noted that the previous studies reporting the affirmation of catabolic capacities of AMF, albeit decomposition of litter only, did not disentangle the effect of exudates from AMF (Hodge et al., 2001; Cheng et al., 2012). It may be speculated that the AMF-induced decomposition may actually come from effect of exudates delivered through them since their ability to produce extra-cellular enzymes is very limited in comparison to ericoid and ecto-mycorrhizal fungi (Read and Perez-Moreno, 2003). However we suggest that this result must be verified with other plants although it supports the idea that AMF have limited decomposing capacity.

Root exudates and the presence of live and dead roots (1000 μm mesh) strongly differ with respect to their effect on SOM mineralization and microbial biomass (Figs. 2 & 4). The amount of RPE induced per unit of plant-derived C assimilated into microbial biomass is 3 to 7 times higher for exudates than for root litter (Fig. 5). This result indicates that the exuded C is used by microbes mostly to synthesize and release extra-cellular enzymes mineralizing SOM instead of promoting microbial growth (Fig 2) and N immobilization (Fig. 1, Supplementary material). This result supports the idea that the RPE is a sort of indirect co-evolved mutualism between plants and rhizosphere microbes (Cheng et al, 2013). The microbial preference for mineralizing activity over growth when relatively small amounts of resource (exudates) are available may have been in anticipation of bigger amounts of a resource (De Nobili et al, 2001). The bigger amounts become available when roots are also present (1000 μm mesh) allowing them to prefer growth over activity which is evident in little increase in the RPE in 1000 μm mesh treatment (Fig. 2). In addition, the microbes are less growth efficient when only exudates are available i.e. they uptake less plant carbon (Fig. 4). The exudates, though a labile and enriched source of energy like a 'fast-food', may soon render microbes nutrient-deficient thereby

driving them to accelerate SOM decomposition (RPE). Whereas root litter a relatively complex substrate provides complete diet more likely to support microbial growth. This indirect mutualistic relationship also enables plants to not only provide energy (i.e. exudates) to soil microorganisms but also control the way of energy is used in order to maximize SOM mineralization and their return on investment (greater nutrient availability).

The absence of a substantial shift in microbial community structure in the presence of exudates and the significant increase in saprophytic fungi in the presence of root litter (Fig 6, Table 2 Supplementary material) confirms differences between the two sources of C deposition *vis-à-vis* their use by the soil microbial community. It seems that exudates are good enough for only stimulating the *k*-strategist microbes (Fontaine et al., 2003) to produce and release extracellular enzymes decomposing SOM. Moreover, the growth of *r*-strategist microorganisms (Fontaine et al., 2003), which do not decompose SOM but can grow rapidly on substrates similar to exudates, is blocked by a factor that could not be identified in this study. In contrast, root litter raised saprophytic fungi specifically the biomarker 18:2w6c across all plants. These saprophytic fungi are considered the actors of the PE induced by plant litter (Fontaine et al., 2011; Shahzad, 2012; Shahzad et al., 2012).

In conclusion, our findings show that the plant strongly modulates SOM mineralization (rhizosphere priming effect) through their exudates. This modulation is likely the result of 100,000s of years of plant-microbe co-evolution which has been realized in an atmosphere where CO₂ concentration mostly fluctuated between 200 and 280 ppm (Barnola et al., 1987). However, the human activities have led to a rapid increase in atmospheric CO₂ concentration currently reaching 400 ppm (Mauna Loa Observatory, 2014), with the possibility of disturbing the exudation rate of plants (Phillips et al., 2011). The mineralization of SOM in ecosystems

exposed to elevated CO₂ is intensified (Langley et al., 2009; Phillips et al., 2012) leading, in several cases, to net decrease in SOM stock (Carney et al., 2007; Langley et al., 2009) and an increase in N leaching (Liu et al., 2008; Hungate et al., 2014). These findings suggest that an increase in CO₂ concentration, by increasing the exudation rate of plants and thereby the RPE leads to a rupture of the synchronization (Perveen et al., 2014) between microbial mineralization of SOM and plant uptake of nutrients. Further experiments are necessary to precise the role of exudates in the plant-microbe synchronization in a changing environment.

Acknowledgement

We thank S. Revaillet and O. Delfosse for chemical and isotopic analyses. Maurice Crocombette and Patrick Pichon are thanked for their valuable help in chamber construction and experiment establishment respectively. The research received funding from the European Community's 6th and 7th Framework Programs (FP6 and FP7) under grant agreement no. 017841 (Nitro-Europe) and no. 226701 (CARBO-Extreme). The funding from National Research Agency's projects BIOMOS and DIMIMOS is also acknowledged. TS & NP have been funded by Higher Education Commission of Pakistan (HEC) and TS European Community's 7th framework under grant agreement no. 226701 (CARBO-EXTREME).

References

- Bahn, M., Schmitt, M., Siegwolf, R., Richter, A., Brüggemann, N., 2009. Does photosynthesis affect grassland soil-respired CO₂ and its carbon isotope composition on a diurnal timescale? *New Phytologist* 182, 451–460.
- Barnola, J.M., Raynaud, D., Korotkevich, Y.S., Lorius, C., 1987. Vostok ice core provides 160,000-year record of atmospheric CO₂. *Nature* 329, 408–414.
- Bell, J.M., Smith, J.L., Bailey, V.L., H, B., 2003. Priming effect and C storage in semi-arid no-till spring crop rotations. *Biology and Fertility of Soils* 37, 237–244.
- Bingeman, C.W., Varner, J.E., Martin, W.P., 1953. The effect of the addition of organic materials on the decomposition of an organic soil. *Soil Science Society of America Journal* 17, 34–38.
- Blagodatskaya, E.V., Blagodatsky, S. a., Anderson, T.-H., Kuzyakov, Y., 2007. Priming effects in Chernozem induced by glucose and N in relation to microbial growth strategies. *Applied Soil Ecology* 37, 95–105.
- Bligh, E.G., Dyer, W.J., 1959. A rapid method of lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology* 37, 911–917.
- Broeckling, C.D., Broz, A.K., Bergelson, J., Manter, D.K., Vivanco, J.M., 2008. Root exudates regulate soil fungal community composition and diversity. *Applied and Environmental Microbiology* 74, 738–744.
- Carney, K.M., Hungate, B.A., Drake, B.G., Megonigal, J.P., 2007. Altered soil microbial community at elevated CO₂ leads to loss of soil carbon. *Proceedings of the National Academy of Sciences of the United States of America* 104, 4990–4995.
- Chen, X., Wu, C., Tang, J., Hu, S., 2005. Arbuscular mycorrhizae enhance metal lead uptake and growth of host plants under a sand culture experiment. *Chemosphere* 60, 665–671.
- Cheng, L., Booker, F.L., Tu, C., Burkey, K.O., Zhou, L., Shew, H.D., Ruffy, T.W., Hu, S., 2012. Arbuscular mycorrhizal fungi increase organic carbon decomposition under elevated CO₂. *Science* 337, 1084–1087.
- Cheng, W., 2009. Rhizosphere priming effect: Its functional relationships with microbial turnover, evapotranspiration, and C–N budgets. *Soil Biology and Biochemistry* 41, 1795–1801.
- Cheng, W., Parton, W.J., Gonzalez-meler, M.A., Phillips, R., Asao, S., McNickle, G.G., Brzostek, E., Jastrow, J.D., 2013. Synthesis and modeling perspectives of rhizosphere priming. *New Phytologist* 201, 31–44.

- Clarholm, M., 1985. Interactions of bacteria, protozoa and plants leading to mineralization of soil nitrogen. *Soil Biology & Biochemistry* 17, 181–187.
- Dalenberg, J.W., Jager, G., 1981. Priming effect of small glucose additions to ¹⁴C-labelled soil. *Soil Biology & Biochemistry* 13, 219–223.
- Dalenberg, J.W., Jager, G., 1989. Priming effect of some organic additions to ¹⁴C-labelled soil. *Soil Biology & Biochemistry* 21, 443–448.
- Davidson, E. A., Janssens, I. A., 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440, 165–73.
- De Deyn, G.B., Quirk, H., Oakley, S., Ostle, N., Bardgett, R.D., 2011. Rapid transfer of photosynthetic carbon through the plant-soil system in differently managed species-rich grasslands. *Biogeosciences* 8, 1131–1139.
- De Nobili, M., Contin, M., Mondini, C., Brookes, P., 2001. Soil microbial biomass is triggered into activity by trace amounts of substrate. *Soil Biology & Biochemistry* 33, 1163–1170.
- Denef, K., Roobroeck, D., Manimel Wadu, M.C.W., Lootens, P., Boeckx, P., 2009. Microbial community composition and rhizodeposit-carbon assimilation in differently managed temperate grassland soils. *Soil Biology & Biochemistry* 41, 144–153.
- Dijkstra, F.A., Carrillo, Y., Pendall, E., Morgan, J.A., 2013. Rhizosphere priming: a nutrient perspective. *Frontiers in Microbiology* 4, 1–8.
- Dijkstra, F.A., Cheng, W., 2007. Interactions between soil and tree roots accelerate long-term soil carbon decomposition. *Ecology letters* 10, 1046–53.
- Dijkstra, F.A., Cheng, W., Johnson, D.W., 2006a. Plant biomass influences rhizosphere priming effects on soil organic matter decomposition in two differently managed soils. *Soil Biology & Biochemistry* 38, 2519–2526.
- Dijkstra, F.A., Hobbie, S.E., Reich, P.B., 2006b. Soil processes affected by sixteen grassland species grown under different environmental conditions. *Soil Science Society of America Journal* 70, 770–777.
- Dowling, N.J.E., Widdel, F., White, D.C., 1986. Phospholipid ester-linked fatty acid biomarkers of acetate-oxidizing sulphate-reducers and other sulphide-forming bacteria. *Journal of General Microbiology* 132, 1815–1825.
- Drake, J.E., Gallet-Budynek, A., Hofmockel, K.S., Bernhardt, E.S., Billings, S.A., Jackson, R.B., Johnsen, K.S., Lichter, J., McCarthy, H.R., McCormack, M.L., Moore, D.J.P., Oren, R., Palmroth, S., Phillips, R.P., Phippen, J.S., Pritchard, S.G., Treseder, K.K., Schlesinger, W.H., Delucia, E.H., Finzi, A.C., 2011. Increases in the flux of carbon belowground stimulate

- nitrogen uptake and sustain the long-term enhancement of forest productivity under elevated CO₂. *Ecology letters* 14, 349–357.
- Drake, J.E., Darby, B. a., Giasson, M. -a., Kramer, M. a., Phillips, R.P., Finzi, a. C., 2013. Stoichiometry constrains microbial response to root exudation- insights from a model and a field experiment in a temperate forest. *Biogeosciences* 10, 821–838.
- Elias, K.S., Safir, G.R., 1987. Hyphal elongation of *Glomus fasciculatus* in response to root exudates. *Applied and Environmental Microbiology* 53, 1928–1933.
- Eliasson, P.E., McMurtrie, R.E., Pepper, D.A., Stromgren, M., Linder, S., Agren, G.I., 2005. The response of heterotrophic CO₂ flux to soil warming. *Global Change Biology* 11, 167–181.
- Eschen, R., Müller-Schärer, H., Schaffner, U., 2013. Plant interspecific differences in arbuscular mycorrhizal colonization as a result of soil carbon addition. *Mycorrhiza* 23, 61–70.
- Fontaine, S., Bardoux, G., Abbadie, L., Mariotti, A., 2004a. Carbon input to soil may decrease soil carbon content. *Ecology Letters* 7, 314–320.
- Fontaine, S., Bardoux, G., Benest, D., Verdier, B., Mariotti, A., Abbadie, L., 2004b. Mechanisms of the priming effect in a savannah soil amended with cellulose. *Soil Science Society of America Journal* 68, 125–131.
- Fontaine, S., Henault, C., Aamor, A., Bdioui, N., Bloor, J.M.G., Maire, V., Mary, B., Revaillet, S., Maron, P.A., 2011. Fungi mediate long term sequestration of carbon and nitrogen in soil through their priming effect. *Soil Biology & Biochemistry* 43, 86–96.
- Fontaine, S., Mariotti, A., Abbadie, L., 2003. The priming effect of organic matter: a question of microbial competition? *Soil Biology & Biochemistry* 35, 837–843.
- Frostegård, A., Bååth, E., 1996. The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil. *Biology and Fertility of Soils* 22, 59–65.
- Frostegård, Å., Tunlid, A., Bååth, E., 1991. Microbial biomass measured as total lipid phosphate in soils of different organic content. *Journal of Microbiological Methods* 14, 151–163.
- Frostegård, Å., Tunlid, A., Bååth, E., 2011. Use and misuse of PLFA measurements in soils. *Soil Biology & Biochemistry* 43, 1621–1625.
- Germida, J.J., Siciliano, S.D., Renato de Freitas, J., Seib, A.M., 1998. Diversity of root-associated bacteria associated with field-grown canola (*Brassica napus* L.) and wheat (*Triticum aestivum* L.). *FEMS Microbiology Ecology* 26, 43–50.
- Grayston, S.J., Vaughan, D., Jones, D., 1997. Rhizosphere carbon flow in trees, in comparison with annual plants: the importance of root exudation and its impact on microbial activity and nutrient availability. *Applied Soil Ecology* 5, 29–56.

- Hamer, U., Marschner, B., 2005a. Priming effects in different soil types induced by fructose, alanine, oxalic acid and catechol additions. *Soil Biology & Biochemistry* 37, 445–454.
- Hamer, U., Marschner, B., 2005b. Priming effects in soils after combined and repeated substrate additions. *Geoderma* 128, 38–51.
- Hamilton, E.W., Frank, D.A., 2001. Can plants stimulate soil microbes and their own nutrient supply? Evidence from a grazing tolerant grass. *Ecology* 82, 2397–2402.
- Helal, H.M., Sauerbeck, D.R., 1984. Influence of plant roots on C and P metabolism in soil. *Plant and Soil* 76, 175–182.
- Hodge, A., Campbell, C.D., Fitter, A.H., 2001. An arbuscular mycorrhizal fungus accelerates decomposition and acquires nitrogen directly from organic material. *Nature* 413, 297–299.
- Hungate, B.A., Duval, B.D., Dijkstra, P., Johnson, D.W., Ketterer, M.E., Stiling, P., Cheng, W., Millman, J., Hartley, A., Stover, D.B., 2014. Nitrogen inputs and losses in response to chronic CO₂ exposure in a sub-tropical oak woodland. *Biogeosciences Discussions* 11, 61–106.
- Hütsch, B.W., Augustin, J., Merbach, W., 2002. Plant rhizodeposition — an important source for carbon turnover in soils. *Journal of Plant Nutrition and Soil Science* 165, 397–407.
- Johnson, D., Leake, J.R., Ostle, N., Ineson, P., Read, D.J., 2002. In situ ¹³CO₂ pulse-labelling of upland grassland demonstrates a rapid pathway of carbon flux from arbuscular mycorrhizal mycelia to the soil. *New Phytologist* 153, 327–334.
- Jones, D.L., Hodge, A., Kuzyakov, Y., 2004. Plant and mycorrhizal regulation of rhizodeposition. *New Phytologist* 163, 459–480.
- Kaiser, C., Koranda, M., Kitzler, B., Fuchslueger, L., Schnecker, J., Schweiger, P., Rasche, F., Zechmeister-Boltenstern, S., Sessitsch, A., Richter, A., 2010. Belowground carbon allocation by trees drives seasonal patterns of extracellular enzyme activities by altering microbial community composition in a beech forest soil. *New Phytologist* 187, 843–858.
- Kuzyakov, Y., Cheng, W., 2001. Photosynthesis controls of rhizosphere respiration and organic matter decomposition. *Soil Biology & Biochemistry* 33, 1915–1925.
- Langley, J.A., McKinley, D.C., Wolf, A.A., Hungate, B.A., Drake, B.G., Megonigal, J.P., 2009. Priming depletes soil carbon and releases nitrogen in a scrub-oak ecosystem exposed to elevated CO₂. *Soil Biology & Biochemistry* 41, 54–60.
- Leigh, J., Hodge, A., Fitter, A.H., 2009. Arbuscular mycorrhizal fungi can transfer substantial amounts of nitrogen to their host plant from organic material. *New Phytologist* 181, 199–207.

- Liljeroth, E., Kuikman, P., Veen, J.A., 1994. Carbon translocation to the rhizosphere of maize and wheat and influence on the turnover of native soil organic matter at different soil nitrogen levels. *Plant and Soil* 161, 233–240.
- Lipton, D.S., Blanchar, R.W., Blevins, D.G., 1987. Citrate, Malate, and Succinate concentration in exudates from P-sufficient and P-stressed *Medicago sativa* L. seedlings. *Plant Physiology* 85, 315–317.
- Liu, J.X., Zhang, D.Q., Zhou, G.Y., Faivre-Vuillin, B., Deng, Q., Wang, C.L., 2008. CO₂ enrichment increases nutrient leaching from model forest ecosystems in subtropical China. *Biogeosciences* 5, 1783–1795.
- López-Bucio, J., de La Vega, O.M., Guevara-García, A., Herrera-Estrella, L., 2000. Enhanced phosphorus uptake in transgenic tobacco plants that overproduce citrate. *Nature Biotechnology* 18, 450–453.
- Maire, V., Gross, N., Da Silveira Pontes, L., Picon-Cochard, C., Soussana, J.-F., 2009. Trade-off between root nitrogen acquisition and shoot nitrogen utilization across 13 co-occurring pasture grass species. *Functional Ecology* 23, 668–679.
- Mauna Loa Observatory, 2014. <http://www.esrl.noaa.gov/gmd/ccgg/trends/> [WWW Document]. URL <http://co2now.org/images/stories/data/co2-atmospheric-mlo-monthly-scripps.pdf> (accessed 5.19.14).
- McGonigle, T.P., Miller, M.H., Evans, D.G., Fairchild, G.L., Swan, J.A., 1990. A new method which gives an objective measure of colonization of roots by vesicular—arbuscular mycorrhizal fungi. *New Phytologist* 115, 495–501.
- Medina, A., Vassilev, N., Azcón, R., 2010. The interactive effect of an AM fungus and an organic amendment with regard to improving inoculum potential and the growth and nutrition of *Trifolium repens* in Cd-contaminated soils. *Applied Soil Ecology* 44, 181–189.
- Moghimi, A., Lewis, D.G., Oades, J.M., 1978. Release of phosphate from calcium phosphates by rhizosphere products. *Soil Biology & Biochemistry* 10, 277–281.
- Murashige, T., Skoog, F., 1962. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia Plantarum* 15, 473–497.
- Nguyen, C., 2009. Rhizodeposition of Organic C by Plant: Mechanisms and Controls, in: Lichtfouse, E., Navarrete, M., Debaeke, P., Véronique, S., Alberola, C. (Eds.), *Sustainable Agriculture*. Springer Netherlands, pp. 97–123.
- Nichols, P., Stulp, B.K., Jones, J.G., White, D.C., 1986. Comparison of fatty acid content and DNA homology of the filamentous gliding bacteria *Vitreoscilla*, *Flexibacter*, *Filibacter*. *Archives of Microbiology* 146, 1–6.

- Nottingham, A.T., Griffiths, H., Chamberlain, P.M., Stott, A.W., Tanner, E.V.J., 2009. Soil priming by sugar and leaf-litter substrates: A link to microbial groups. *Applied Soil Ecology* 42, 183–190.
- Ohm, H., Hamer, U., Marschner, B., 2007. Priming effects in soil size fractions of a podzol Bs horizon after addition of fructose and alanine. *Journal of Plant Nutrition and Soil Science* 170, 551–559.
- Olsson, P.A., Bååth, E., Jakobsen, I., Söderström, B., 1995. The use of phospholipid and neutral lipid fatty acids to estimate biomass of arbuscular mycorrhizal fungi in soil. *Mycological Research* 99, 623–629.
- Pascualt, N., Ranjard, L., Kaisermann, A., Bachar, D., Christen, R., Terrat, S., Mathieu, O., Lévêque, J., Mougél, C., Henault, C., Lemanceau, P., Péan, M., Boiry, S., Fontaine, S., Maron, P.-A., 2013. Stimulation of different functional groups of bacteria by various plant residues as a driver of soil priming effect. *Ecosystems* 16, 810–822.
- Paterson, E., 2003. Importance of rhizodeposition in the coupling of plant and microbial productivity. *European Journal of Soil Science* 741–750.
- Personeni, E., Loiseau, P., 2004. How does the nature of living and dead roots affect the residence time of carbon in the root litter continuum? *Plant and Soil* 267, 129–141.
- Perveen, N., Barot, S., Alvarez, G., Klumpp, K., Martin, R., Rapaport, A., Herfurth, D., Louault, F., Fontaine, S., 2014. Priming effect and microbial diversity in ecosystem functioning and response to global change: a modeling approach using the SYMPHONY model. *Global Change Biology* 20, 1174–1190.
- Phillips, R.P., Finzi, A.C., Bernhardt, E.S., 2011. Enhanced root exudation induces microbial feedbacks to N cycling in a pine forest under long-term CO₂ fumigation. *Ecology Letters* 14, 187–194.
- Phillips, R.P., Meier, I.C., Bernhardt, E.S., Grandy, A.S., Wickings, K., Finzi, A.C., 2012. Roots and fungi accelerate carbon and nitrogen cycling in forests exposed to elevated CO₂. *Ecology Letters* 15, 1042–1049.
- Ratnayake, B.M., Leonard, R.T., Menge, J.A., 1978. Root exudation in relation to supply of phosphorus and its possible relevance to mycorrhizal formation. *New Phytologist* 81, 543–552.
- Read, D.J., Perez-Moreno, J., 2003. Mycorrhizas and nutrient cycling in ecosystems - a journey towards relevance? *New Phytologist* 157, 475–492.
- Saleque, M.A., Kirk, G.J.D., 1995. Root-induced solubilization of phosphate in the rhizosphere of lowland rice. *New Phytologist* 129, 325–336.

- Salomé, C., Nunan, N., Pouteau, V., Lerch, T.Z., Chenu, C., 2010. Carbon dynamics in topsoil and in subsoil may be controlled by different regulatory mechanisms. *Global Change Biology* 16, 416–426.
- Sauer, D., Kuzyakov, Y., Stahr, K., 2006. Spatial distribution of root exudates of five plant species as assessed by ^{14}C labeling. *Journal of Plant Nutrition and Soil Science* 169, 360–362.
- Schimel, D.S., 1995. Terrestrial ecosystems and the carbon cycle. *Global Change Biology* 1, 77–91.
- Schimel, J.P., Wetterstedt, J.Å.M., Holden, P.A., Trumbore, S.E., 2011. Drying/rewetting cycles mobilize old C from deep soils from a California annual grassland. *Soil Biology and Biochemistry* 43, 1101–1103.
- Shahzad, T., 2012. Role of plant rhizosphere across multiple species, grassland management and temperature on microbial communities and long term soil organic matter dynamics. AgroParisTech, Paris, France. 180pp PhD Thesis.
- Shahzad, T., Chenu, C., Repinçay, C., Mougin, C., Ollier, J.-L., Fontaine, S., 2012. Plant clipping decelerates the mineralization of recalcitrant soil organic matter under multiple grassland species. *Soil Biology & Biochemistry* 51, 73–80.
- Talbot, J.M., Allison, S.D., Treseder, K.K., 2008. Decomposers in disguise : mycorrhizal fungi as regulators of soil C dynamics in ecosystems under global change. *Functional Ecology* 22, 955–963.
- Tawarayama, K., Watanabe, S., Yoshida, E., Wagatsuma, T., 1996. Effect of onion (*Allium cepa*) root exudates on the hyphal growth of *Gigaspora margarita*. *Mycorrhiza* 6, 57–59.
- Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil microbial biomass C. *Soil Biology & Biochemistry* 19, 703–707.
- Wu, J., Brookes, P.C., Jenkinson, D.S., 1993. Formation and destruction of microbial biomass during the decomposition of glucose and ryegrass in soil. *Soil Biology & Biochemistry* 25, 1435–1441.
- Zelles, L., 1997. Phospholipid fatty acid profiles in selected members of soil microbial communities. *Chemosphere* 35, 275–294.
- Zhu, B., Cheng, W., 2011. Rhizosphere priming effect increases the temperature sensitivity of soil organic matter decomposition. *Global Change Biology* 17, 2172–2183.

Figure Legends

Figure 1. Top: The schematic diagram of sand filled pot that was used to grow plants on, containing soil-filled small PVC boxes. Below: The destructive sampling of plant roots and soil from real pot.

Figure 2. R_s i.e. soil-derived $\text{CO}_2\text{-C}$ ($\text{mg C kg}^{-1} \text{ soil day}^{-1}$) from planted and bare soils under three grassland species. Error bars represent standard error of means.

Figure 3. Relationship between rhizosphere priming effect (RPE) and (a) R_p , the $\text{CO}_2\text{-C}$ coming from plant, mycorrhizae and microbial respiration of rhizodeposits and plant litter ($\text{mg CO}_2\text{-C kg}^{-1} \text{ soil day}^{-1}$) (b) total biomass produced by the plants across all species during final measurement of soil CO_2 efflux. Pt, *Poa trivialis*; Lp, *Lolium perenne*; Tr, *Trifolium perenne*.

Figure 4. Microbial biomass (MB) in control and planted treatments with three meshes. Blank bar represents total MB in control soils where MB_{tot} shows total biomass, MB_{soc} soil-derived microbial biomass and MB_{plant} plant-derived microbial biomass in planted treatments.

Figure 5. Rhizosphere priming effect efficiency i.e. rhizosphere priming effect induced per unit of plant derived carbon assimilated in microbial biomass, in different mesh treatments under three grassland species.

Figure 6. Concentrations of phospholipids fatty acids of AMF, saprophytic fungi, Gram negative and Gram positive bacteria in planted and control soils as affected by C deposition through root exudates only (0.45 μ mesh), root exudates and root-associated mycorrhizae (30 μ mesh) and root exudates plus root-associated mycorrhizae and roots (1000 μ mesh).

Figure S1. Figure S1. (a) Microbial N (mg N kg^{-1} soil) and (b) microbial C/N under three grassland species and mesh treatments as compared to bare soil.

Figure S2: Plant-derived i.e. ^{13}C -labeled C humified in soil organic matter (g C/kg soil) under the three plant species and different mesh treatments.




Figure 1. Diagram of plant-soil microcosm (a) . And soil boxes enclosed by mesh of $0.45\ \mu\text{m}$ (b), $30\ \mu\text{m}$ (c) and $1000\ \mu\text{m}$ (d) being taken out from pots. Circles point to assemblage of roots against mesh.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.




Figure S1.




Figure 2S: Plant-derived i.e. ^{13}C -labeled C humified in soil organic matter (g C/kg soil) under the three species and different mesh treatments.