



HAL
open science

Evaluation and validation of an analytical approach for high-throughput metabolomic fingerprinting using direct introduction–high-resolution mass spectrometry: Applicability to classification of urine of scrapie-infected ewes

Estelle Rathahao-Paris, Sandra Alves, Nawel Boussaid, Nicole Picard-Hagen, Véronique V. Gayrard-Troy, Pierre-Louis Toutain, Jean-Claude Tabet, Douglas N. Rutledge, Alain Paris

► **To cite this version:**

Estelle Rathahao-Paris, Sandra Alves, Nawel Boussaid, Nicole Picard-Hagen, Véronique V. Gayrard-Troy, et al.. Evaluation and validation of an analytical approach for high-throughput metabolomic fingerprinting using direct introduction–high-resolution mass spectrometry: Applicability to classification of urine of scrapie-infected ewes. *European Journal of Mass Spectrometry*, 2019, 25 (2), pp.251-258. 10.1177/1469066718806450 . hal-02317795v1

HAL Id: hal-02317795

<https://agroparistech.hal.science/hal-02317795v1>

Submitted on 18 Oct 2019 (v1), last revised 7 Nov 2019 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Evaluation of direct introduction-high resolution mass spectrometry (DI-HRMS) based metabolomics approach for the discrimination of several exposure groups to pesticides: comparison with liquid chromatography high resolution mass spectrometry (LC/HRMS)

Baninia Habchi^{1,2}, Sandra Alves^{2*}, Delphine Jouan-Rimbaud Bouveresse³, Bilel Moslah¹, Alain Paris⁴, Yannick Lécluse⁵, Pascal Gauduchon⁵, Pierre Lebailly⁵, Douglas N. Rutledge¹, Estelle Rathahao-Paris^{1,2*}

¹ UMR Ingénierie Procédés Aliments, AgroParisTech, Inra, Université Paris-Saclay, 91300 Massy, France

² Sorbonne Université, Faculté des Sciences et de l'Ingénierie, Institut Parisien de Chimie Moléculaire (IPCM), F-75005 Paris, France

³ UMR 914 Physiologie de la Nutrition et du Comportement Alimentaire, INRA, AgroParisTech, Université Paris-Saclay, F-75005 Paris

⁴ Muséum national d'Histoire naturelle, MCAM, UMR7245 CNRS - MNHN, 75005 Paris, France

⁵ ANTICIPE U1086 INSERM & Université Caen-Normandie, Centre François Baclesse, 14076 Caen Cedex 05 France

Abstract

One of the current public health concerns is the evaluation of population exposure to toxicants present in food and environment. To do this, high-throughput methods are required to perform a large number of analyses. The aim of this work is to characterize human exposure to xenobiotics. The classification of urines from farmers occupationally exposed to five pesticides was investigated using two analytical techniques, *i.e.* the

*Corresponding authors :

Dr. Estelle Rathahao-Paris

Present address: UMR CEA-INRA Service de Pharmacologie et d'Immunoanalyse, Laboratoire d'Immuno-Allergie Alimentaire, CEA de Saclay- Bat 136, F-91191, Gif-sur-Yvette cedex, FRANCE. E-mail: Estelle.Paris@inra.fr

or

Dr. Sandra Alves

Address: Sorbonne Université, Faculté des Sciences et de l'Ingénierie, Tour 42-43, 4^{ème} étage, BP 45, 4 place Jussieu F-75005, Paris, FRANCE. E-mail: sandra.alves@sorbonne-universite.fr

1
2
3 classical approach based on liquid chromatography coupled to high resolution mass
4 spectrometry (LC/HRMS) and direct introduction-high resolution mass spectrometry (DI-
5 HRMS) to produce metabolomics data. Discrimination of different exposure groups was
6 obtained from LC/MS data as well as from DI-HRMS data thanks to the Independent
7 Component - Discriminant Analysis (IC-DA). The DIMS approach requiring only a
8 fraction of the LC/MS analysis time, appears to be very promising for high throughput
9 epidemiological studies.
10
11
12
13
14
15
16
17
18
19
20

21 **Keywords:** Metabolomics, Pesticide exposure, Direct introduction mass spectrometry,
22 Liquid chromatography mass spectrometry, High resolution mass spectrometry,
23 Independent component - discriminant analysis.
24
25
26
27
28
29
30

31 **1. Introduction**

32
33 The presence of toxicants (*e.g.*, pesticides, drugs, pollutants or contaminants) in food and
34 environment can impact human health. Due to the great diversity of xenobiotics and the
35 complexity to develop targeted methods specific to a molecule or a compound family,
36 global approaches without a priori such as metabolomics seem to be appropriate in such
37 purpose. Metabolomics ¹ is the study of small molecular mass metabolites found in
38 biological samples, which are involved in growth, metabolic homeostasis or etiology of
39 disease. It aims at the detection, quantification and subsequent identification of different
40 metabolic changes arising between two (or more) well-defined but contrasted situations
41 (*e.g.*, healthy *versus* different unhealthy status). Analytical techniques combined with
42 statistical and/or chemometric processing tools are essential to compare generated
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 metabolomic fingerprints reflecting phenotypic differences and to reveal the metabolic
4
5 biomarkers of a given physiological condition or a pathological state ².

6
7
8 Among the available analytical technologies, mass spectrometry (MS) providing high
9
10 sensitivity, specificity and selectivity, and especially, Fourier transform mass
11
12 spectrometry (FTMS) with its very high mass resolving power, high mass measurement
13
14 accuracy and wide dynamic range should produce high quality of metabolomic profiles in
15
16 term of isobaric species distinction and number of detectable peaks ^{3,4}. MS-based
17
18 metabolomic analysis commonly uses hyphenated methods, typically liquid or gas
19
20 chromatography coupled to mass spectrometry (LC/MS or GC/MS). The hyphenated
21
22 approaches have the main advantages of being able to separate isobaric and some
23
24 isomeric compounds and also to minimize the so-called matrix effects. They should
25
26 enhance the metabolome coverage ^{5,6}. However, they cannot be really considered as high
27
28 throughput approaches because of the analysis time. Although the use of ultra high
29
30 performance liquid chromatography (UHPLC) can reduce the data acquisition time, its
31
32 coupling to the FTMS is done at the expense of the mass resolving power. Furthermore,
33
34 the pre-processing of the anisotropic data generated from several scans acquired along the
35
36 chromatographic separation requires dedicated pretreatment steps such as peak alignment
37
38 and correction for analytical drifts (*e.g.*, retention time and peak intensity). Alternatively,
39
40 the direct introduction mass spectrometry (DIMS) approach enabling simultaneous
41
42 detection of all ionized compounds provides a seemingly isotropic spectral fingerprint in
43
44 a few minutes. Such direct approach can therefore be proposed to increase the throughput
45
46 of analyses ⁷⁻⁹. A comparison between DIMS and LC/MS approaches has been reported
47
48 for the analysis of serum samples from kidney cancer patients ¹⁰. A comparable group
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 classification was obtained from both approaches despite the smaller number of
4
5 metabolites detected from DIMS data (*i.e.* 23 metabolites compared to a total number of
6
7 48 ions from LC/MS data). In addition, only a small fraction of the LC/MS analysis time
8
9 (5%) is required to perform the DIMS experiment. Unfortunately, the DIMS approach
10
11 using electrospray ionization (ESI) has some limitations, especially the occurrence of
12
13 matrix effects. Such phenomena are characterized by ion suppression and/or signal
14
15 enhancement, leading to unpredictable signal changes that may affect the quantitative
16
17 response of the sample constituents and consequently modify the mass spectrum profiles
18
19 ¹¹. These effects should be overcome or at least minimized through the optimization of
20
21 analytical conditions. Sample dilution can be proposed to obtain metabolomics
22
23 fingerprints containing the most informative data with minimized matrix effects without
24
25 loss of peak intensity ¹². The use of high resolution mass spectrometry (HRMS) such as
26
27 FTMS can compensate the absence of chromatographic separation by distinguishing
28
29 compounds based on accurate m/z values and isobar ion resolution. Nevertheless, isomer
30
31 distinction requires additional experiments such as the chromatographic separation and/or
32
33 MS/MS experiments or other alternative technologies, such as the ion mobility
34
35 spectrometry.
36
37
38
39
40
41

42 In our previous work, an in-house computational procedure was developed to process DI-
43
44 HRMS data using a new discriminant analysis tool based on Independent Component -
45
46 Discriminant Analysis (IC-DA). Successfully discrimination of farmers occupationally
47
48 exposed to two types of pesticides was obtained, this constitutes the proof of concept of
49
50 our approach for high-throughput metabolomics ¹³.
51
52
53
54
55
56
57
58
59
60

1
2
3 Most of metabolomic studies concern only few sample groups, typically healthy subjects
4 *versus* one (or two) group(s) of unhealthy subjects ^{9,14,15}. Few investigations involving
5 several classes have been reported, *e.g.*, in the context of dose ¹⁶ and time effects ^{17,18}, but
6 rarely in the context of human exposure during their lifestyle ¹⁹. The main objective of
7 this study is to demonstrate the applicability of our high throughput approach to classify
8 several exposure groups. In addition to DI-HRMS, the hyphenated LC/MS technique was
9 also employed to phenotype urine samples coming from farmers occupationally exposed
10 to five different pesticides. Discrimination of the different exposure groups was obtained
11 from both methods, highlighting the potential of the direct approach to provide
12 satisfactory results comparable to those obtained by the LC/MS approach but with a
13 significant gain in the analysis time.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 **2. Experimental**

32 **2.1. Materials**

33
34 Ultra pure water was produced from a Milli-Q system (Merck Millipore, Guyancourt,
35 France). The highest commercial grade of acetonitrile and formic acid was obtained from
36 Sigma Aldrich Chemicals (L'Isle d'Abeau-Chesnes, France). Urine samples were
37 provided by UMR1086 Cancers and Preventions. They were collected from 41 farmers
38 before and after one day of occupational exposure to pesticides in the cultivated fields.
39 The exposure conditions have been described by Lebailly *et al.* in a previous work ^{20,21}.
40 There are five groups with exposure to: i) isoproturon (with n = 6 individuals), ii) captan
41 (n = 16), iii) cyprodinyl (n = 7), iv) chlorothalonil (n = 8) and v) epoxiconazole (n = 4).
42 Note that self-control samples (n=41) are available and correspond to urine of the same
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 subjects collected one-day before pesticide exposure. Note that all procedures performed
4
5 in this work were in accordance with the ethical standards of the national research
6
7 committee and with the 1964 Helsinki declaration and its later amendments. The
8
9 collected urines were stored at -20°C until MS analysis. Urine sample normalization was
10
11 done based on the creatinine concentration. Finally, urine samples were diluted at the
12
13 chosen dilution factor of 500 after testing various dilution factors. Additionally, a quality
14
15 control sample (QC, a pool of all urine samples) was injected after every set of 10 sample
16
17 analyses to check for eventual analytical drifts and for the quality of the generated
18
19 spectral data.
20
21
22
23

24 **2.2. Mass spectrometry detection**

25
26 A hybrid LTQ-Orbitrap Fourier transform mass spectrometer (LTQ Orbitrap XL, Thermo
27
28 Fisher Scientific, Bremen, Germany) operating in positive ion mode was used to perform
29
30 both DIMS and LC/MS analyses. In the DIMS approach, the following ESI conditions
31
32 were used: electrospray voltage was set at -3 kV, capillary voltage at 13 V and tube lens
33
34 offset at 60 V. Sheath gas flow and auxiliary gas were fixed at 70 and 5 arbitrary units
35
36 (a.u.), respectively. The capillary temperature was maintained at 275 °C.
37
38

39
40 Mass spectra were acquired in the m/z 50-1,000 range with a cycle time of 4 minutes per
41
42 analysis. A theoretical mass resolving power of 100,000 FWHM (Full width at half
43
44 maximum) at m/z 400 was used. Internal mass calibration was performed using mass lock
45
46 option on the $[M + H]^+$ contaminant ions of N-butylbenzene sulfonamide at m/z
47
48 214,089625 for both DIMS and LC/MS experiments. Flow injection analysis (FIA) mode
49
50 was used to perform DIMS analyses. 10 μ L of each 500-fold diluted sample were
51
52 injected into a 100 μ L sample loop using a Surveyor LC chromatographic system
53
54
55
56
57
58
59
60

1
2
3 (Thermo Fisher) at a flow rate of 30 $\mu\text{L}/\text{min}$ of an acetonitrile/water mixture (1:1, v/v)
4
5 plus 0.1 % formic acid as eluent. Tandem mass spectrometry (MS/MS) experiments were
6
7 performed on the monoisotopic peaks of some metabolite species. The collision-induced
8
9 dissociation (CID) conditions, the chromatographic separation and mass spectrometry
10
11 conditions used for LC/MS analyses are described in the Supporting Information.
12
13

14 **2.3. Data analysis**

15 **2.3.1 Data pre-processing**

16
17 An in-house DIMS data processing procedure using Matlab routines (Matlab R2008a,
18
19 The MathWorks, Natick MA, USA) was developed in our previous work ¹³ and was used
20
21 in this study to create the DIMS data matrix. Alternatively, the LC/MS data were pre-
22
23 processed using XCMS tool based on a program written in R (*R version 3.4.0, The R*
24
25 *Foundation for Statistical Computing, Vienna 2017, <https://www.r-project.org>*) as
26
27 described in the Supporting Information. The final number of extracted variables for the
28
29 82 urine samples was 747 and 2106 from DIMS and LC/MS data, respectively.
30
31
32
33
34

35 **2.3.2. Chemometric methods**

36
37 Multivariate analyses were further performed identically on both DIMS and LC/MS
38
39 datasets, including all groups in a single model using two Matlab-based chemometric
40
41 tools, as already described ¹³. Briefly, the first preprocessing tool is the orthogonalization
42
43 part of the OPLS algorithm ²². The second tool is a supervised version of the Independent
44
45 Component Analysis (ICA), named IC-DA, as previously described ¹³. ICA aims to
46
47 extract independent vectors, called Independent Components (ICs), corresponding to
48
49 'source signals' with true physico-chemical meaning ^{23–25}. IC-DA includes group
50
51 membership in its calculation to extract signals that discriminate predefined groups.
52
53
54
55
56
57
58
59
60

1
2
3 Therefore, a second calculation was performed using only the spectral signals that
4 contributed to group discrimination to evaluate if the given group is still separated.
5
6 Nevertheless, because of the unbalanced classes in our case, group discrimination can be
7 oriented by those containing more individuals. To avoid eventual overfitting, a validation
8 step was then performed using a permutation procedure described in the previous work ¹³.
9
10
11
12
13

14 **2.3.3. Selection and annotation of metabolic biomarkers**

15
16 A quantile-quantile (Q–Q) plot was used to visualize variables that contribute most to
17 group discrimination ²⁶. The most discriminant variables are characterized by high or low
18 standard deviation values which depart from that linearity while non-interesting variables
19 that follow a normal distribution are aligned. For each IC, only ions with an absolute
20 intensity greater than 3 times the standard deviation of all the ions were selected.
21
22 Alternatively, an additional procedure was performed to validate the selected
23 discriminant variables, as well as to highlight their contribution to group discrimination.
24
25 In this way, a random selection of 50 % of the individuals from each group, followed by
26 IC-DA calculations was performed and repeated 3,000 times for every IC. Then, a
27 quantile calculation with 95 % confidence limit was done on the generated matrix
28 (variable number \times 5 ICs \times 3,000 repetitions). The discriminant variables characterized
29 by a quantile sign (*i.e.*, positive or negative) were selected based on their high standard
30 deviation values. The quantile sign of some selected variables changed after permutation.
31
32 Those variables were then removed from the discriminant variable list.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 Two computational tools such as the box-whisker plot and the Student's t-test were
50 applied to highlight the contribution of a given variable to the discrimination of one (or
51 several) exposure group(s). Box-whisker plots should show the abundance variations of a
52
53
54
55
56
57
58
59
60

1
2
3 given discriminant variable among all groups, including controls. However, these
4
5 variations between groups were not obvious. Therefore, Student's paired t-tests were also
6
7 done to reveal variable intensity variations before (self-control) and after exposure for
8
9 every individual. The paired data structure was used here only to validate discriminant
10
11 variables but it was not applied to all data because rarely each subject could be its own
12
13 control (*e.g.*, self-control) in general population monitoring.
14

15
16 Afterwards, correlation coefficients were calculated between every selected discriminant
17
18 variable and all other (discriminant or not) variables present in the mass spectra in order
19
20 to find possible ion associations, *i.e.*, isotopic peaks, in-source fragments and/or adduct
21
22 species generated from the same molecule. This calculation improves the compound
23
24 annotation and can also reveal eventual biological relations between metabolites (see
25
26 results and discussion). Finally, the most discriminant and correlated variables were
27
28 putatively annotated using freely available databases (DBs) such as the Human
29
30 Metabolome Database (HMDB) ²⁷, the Kyoto Encyclopedia of Genes and Genomes
31
32 database (KEGG) ²⁸ and the METLIN ²⁹ database. The experimental accurate *m/z* values
33
34 with four digits after the decimal point were used to query DBs with a mass error below
35
36 5 ppm, except for ions with *m/z* values lower than 130 u, where 10 ppm mass error was
37
38 used to find candidate compounds.
39
40
41
42
43
44
45
46

47 **3. Results and discussion**

48 **3.1. Exposure group discrimination**

49
50 Efficient discrimination of control and pesticide-exposed groups was obtained from both
51
52 DIMS and LC/MS data using IC-DA (**Figs. S-1** and **S-2** in the supplementary materials).
53
54
55
56
57
58
59
60

1
2
3 A given group is typically discriminated from the others by a specific IC. However, these
4 initial discriminations being obtained taken into account information on the group
5 membership (*i.e.* dummy variables) are overoptimistic. New calculations only based on
6 the MS signals contributing to each specific group discrimination, still provide distinction
7 of all groups as shown in **Fig. 1** and **Fig. S-3** (supplementary materials) for DIMS and
8 LC/MS, respectively. For the DIMS data, a slight overlap between control subjects and
9 other groups is observed (**Fig. 1a**). Nevertheless, efficient separation is obtained for
10 captan, chlorothalonil, cyprodinil and isoproturon exposure groups (**Figs. 1b-e**,
11 respectively). Additionally, discrimination of the epoxiconazol-exposed group is visible
12 in another projection dimension, *i.e.* IC2, IC3 and IC5 (**Fig. 1b-e**). A comparable group
13 discrimination was obtained from LC/MS data (**Fig. S-3**) but the separation of control
14 subjects from others is less pronounced compared to that for DIMS data (**Fig. S-3a**).
15 Moreover, no clear distinction of the epoxiconazol-exposed group was obtained from the
16 LC/MS data.

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The 3D-plots (IC2 vs IC3 vs IC4) from calculations without the group membership information showed good discrimination of the four exposure groups, *i.e.* those exposed to captan, chlorothalonil, cyprodinyl, and epoxiconazol for both DIMS and LC/MS data (**Fig. 2**). Note that the samples from control group are located in the center of the plot and overlapped with the isoproturon-exposed group into the reported 3D plots. These two groups were separated in another dimension (**Fig. S-4** in the supplementary materials).

Hence, data generated from both the DIMS and the LC/MS approaches give comparable and successful discrimination of all exposure groups. More interesting, less intra-group dispersion is observed for DIMS data, enabling better group separation. Such observation

1
2
3 suggests that less technical drifts may occur in the high throughput DIMS approach
4 compared to LC/MS. Importantly, a component characteristic of the control group despite
5 heterogeneity among individuals could be provided by IC-DA for both data sets. Such
6 characterization may be useful for large-scale epidemiological studies in which
7 assignment of control subjects is required for biological interpretation or clinical
8 assessment.
9

10 **3.2. Detection and characterization of discriminant metabolites**

11
12 The most relevant variables involved in the group discrimination of the five exposure
13 types were extracted to be further identified. However, this step is not so easy when
14 considering several exposure groups. It is less straightforward to highlight the
15 contribution of a given variable to one (or more) specific exposure group(s) in our case,
16 compared to classical studies which tend to find the difference between control and an
17 unhealthy group^{9,14,15}. Therefore, different computational tools were used to improve the
18 selection of the most discriminant ions for every specific exposure group (see
19 experimental section).
20

21
22 First, from the Q–Q plot only discriminant variables characterized by high standard
23 deviations were extracted. 25 variables and 134 variables were selected from the DIMS
24 and LC/MS data, respectively, with only four common variables among them. After
25 concatenating very close m/z variables, the number of highly informative discriminant
26 variables decreased to 17 and 115 for DIMS and LC/MS, respectively, and to 16 and 56
27 after application of a permutation procedure (see experimental section).
28

29
30 In a second step, box-whisker plots were used to show the ion abundance changes among
31 all groups. The example of two discriminant variables at m/z 132.0763 and 138.0546
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 selected from DIMS data analysis is illustrated in **Fig. 3**. The response of m/z 132.0763
4 ions is enhanced for samples coming from farmers exposed to captan (**Fig. 3a**),
5 suggesting a high contribution of these species to reveal captan exposure. For ions at m/z
6 138.0546, intensity change between groups is less clear (**Fig. 3b**). Their response is
7 relatively low for the captan exposure group whereas a slightly higher response is
8 obtained in the other exposure groups, *i.e.* those exposed to isoproturon, chlorothalonil
9 and cyprodinyl. In this latter case, it is difficult to affirm the significant variation of peak
10 intensity between different groups. The use of Student's t-test confirmed that 15 variables
11 among the 16 previously selected were significant for the DIMS data (see experimental
12 section), demonstrating the robustness of the DIMS approach (**Table 1**). In contrast, the
13 use of the t-test decreased the number of significant variables from 56 to 29 for the
14 LC/MS data (**Table S-1**). This indicates that these 27 removed variables are not related to
15 biological perturbations but that their intensity was probably affected by technical drifts
16 occurred over a long period of analysis.

17
18 Finally, correlation coefficients were calculated for each discriminant variable to help in
19 metabolite annotation. The putatively annotated discriminant variables (in bold) and the
20 correlated ions detected in the DIMS and LC/MS experiments are reported in **Table 1**
21 and **Table S-1**, respectively. Among the annotated discriminant metabolites, several di or
22 tri-peptides (*e.g.*, leucyl-proline), as well as amino-acids (*e.g.*, tryptophan) and amino
23 acid metabolism intermediates (*e.g.*, creatine, creatinine) were selected from both DIMS
24 and/or LC/MS data (**Table 1** and **Table S-1**). These di- and tri-peptides have been
25 described as breakdown products of protein digestion or protein catabolism. However,
26 their role in physiological function is not well known. Most of them have been reported

1
2
3 as short-lived intermediates formed during proteolysis (HMDB, <http://www.hmdb.ca>).

4
5 Nicotinuric acid, kynurenic acid and xanthurenic acid ions were determined as
6
7 discriminant variables from LC/MS data (see the **Table S-1**). They display a decreased
8
9 intensity for subjects exposed to captan. Kynurenic acid has already been described as a
10
11 metabolite structurally related to xanthurenic acid ³⁰. Additionally, high level of
12
13 xanthurenic acid has been reported to characterize a metabolic perturbation, for example,
14
15 as a putative inducer of malaria development in mosquitoes ³⁰. In our study, the detection
16
17 of this metabolite may reflect metabolic disruption for subjects exposed to captan.
18
19

20
21 Alternately, acylglycine species (furoylglycine, nicotinuric acid, hippuric acid), and
22
23 carnitine or acylcarnitine derivatives (dehydroxycarnitine, acetyl-carnitine, keto-
24
25 decanoylcarnitine, dodecanedioylcarnitine, decadienylcarnitine) were proposed for the
26
27 annotation of some discriminant variables (**Table 1** and **Table S-1**). These metabolites
28
29 have been described as indicators of disturbed metabolic pathways ³¹⁻³⁴. The unusual
30
31 acylcarnitine excretion levels have been reported to be related to diseases such as obesity,
32
33 type 2 diabetes, cardiovascular disease, encephalopathy and carcinoma ^{32,35}. In our study,
34
35 the discriminant ions at m/z 204.1230 could be annotated as acetylcarnitine based on their
36
37 accurate m/z value (*i.e.* C₉H₁₈NO₄ formulae with 0.5 ppm). However, by comparing with
38
39 the CID mass spectrum of the authentic standard compound, the seemingly differences in
40
41 the fragment ion abundances suggest that the m/z 204.1230 ions could correspond to an
42
43 isomer of acetylcarnitine ³⁶ (**Fig. S-6** of the Supplementary materials).
44
45
46
47

48
49 Another example concerns ions correlated to the discriminant ions at m/z 76.0753. The
50
51 **Fig. 4** displays a reconstituted DIMS mass spectrum of a urine sample collected on a
52
53 subject exposed to chlorothalonil. The m/z 76.0753 ions were putatively annotated as
54
55
56
57
58
59
60

1
2
3 [M + H]⁺ ions of trimethylamine N-oxide (C₃H₁₀NO, 5.3 ppm), an oxidation product of
4
5 the trimethylamine. This compound has been described as a marker of oxidative stress.
6
7 Its high concentration has been reported in transplant patients, reflecting the effects of
8
9 kidney diseases ³⁷. These m/z 76.0753 ions appear to be related to ions at m/z 98.0573,
10
11 136.1077, 151.1438, 152.1471 and 189.1345. The presence of ions at m/z 98.0573,
12
13 151.1438 and 152.1471 reinforced the assignment of elemental composition of the m/z
14
15 76.0753 ions (C₃H₁₀NO, 5.3 ppm) since they could correspond to different species of
16
17 trimethylamine N-oxide such as [M + Na]⁺ (C₃H₉NONa, 3 ppm), [2M + H]⁺ (C₆H₁₉N₂O₂,
18
19 3 ppm) and ¹³C isotope peaks of the [2M + H]⁺ (¹³C₁¹²C₅H₁₉N₂O₂, 3 ppm) ions,
20
21 respectively. Additionally, the correlated ions at m/z 136.1077 and 189.1345 could be
22
23 annotated as [M + NH₄]⁺ ions of urea, N-hydroxy-N-propyl (C₄H₁₀N₂O₂, 3 ppm) and
24
25 [M + H]⁺ (C₇H₁₇N₄O₂, 1 ppm) ions of homo-L-arginine or their isomers, respectively. A
26
27 biological connection between these three compounds has already been reported by He et
28
29 al. ³⁸.
30
31
32
33
34
35
36
37

38 **4. Conclusion**

39
40 Discrimination of individuals exposed to the five pesticides was successfully obtained
41
42 with both DIMS and LC/MS data. This result demonstrates the applicability of DI-HRMS
43
44 approach for metabolomic investigations involving several groups of subjects. More
45
46 importantly, the very short data acquisition time makes the DIMS a valuable approach for
47
48 performing high throughput metabolomic fingerprinting, in particular for large scale
49
50 epidemiological investigations which aim to find associations between exposure to
51
52 toxicants and their effects on human health.
53
54
55
56
57
58
59
60

1
2
3 Besides, our approach enabled the detection of some metabolites already described in the
4 literature (*i.e.* urea, creatinine, carnitine, acylcarnitine, acetylcarnitine,
5 dehydroxycarnitine, keto-decanoylcarnitine, dodecanedioylcarnitine) as indicators of
6 toxic exposure. Pesticide exposure has been reported to cause significant DNA damage
7
8 ²⁰, possible effects on fetuses in pregnant women ¹⁹ and cardiac dysfunction ³⁹. Such an
9 exposure could impact metabolism pathways resulting in changes of metabolite
10 concentrations ^{19,34,40–42}. Further studies are needed to provide more accurate
11 toxicological evaluation and precise identification as well as biological interpretation of
12 biomarkers specifically related to pesticide exposure.
13
14
15
16
17
18
19
20
21
22

23 **Electronic supplementary information**

24
25 Supplementary data related to this article can be found: LC/MS experimental procedure
26 and additional results illustrated by Figures S-1 to S-6 and table S-1.
27
28
29
30

31 **Conflicts of interest**

32
33 The authors declare no conflict of interest.
34
35

36 **Acknowledgments**

37
38 Authors gratefully acknowledge the funding received towards Baninia Habchi's PhD
39 from the Region Ile-de-France and Dim Analytics.
40
41
42

43 **References**

- 44
45
46 1 S. G. Oliver, M. K. Winson, D. B. Kell and F. Baganz, *Trends Biotechnol.*, 1998, **16**, 373–378.
47
48 2 R. Goodacre, S. Vaidyanathan, W. B. Dunn, G. G. Harrigan and D. B. Kell, *Trends Biotechnol.*,
49 2004, **22**, 245–252.
50
51 3 C. Junot, G. Madalinski, J.-C. Tabet and E. Ezan, *Analyst*, 2010, **135**, 2203–2219.
52
53 4 E. Rathahao-Paris, S. Alves, C. Junot and J.-C. Tabet, *Metabolomics*, 2016, **12**, 10.
54
55 5 I. D. Wilson, R. Plumb, J. Granger, H. Major, R. Williams and E. M. Lenz, *J. Chromatogr. B*,
56 2005, **817**, 67–76.
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 6 K. K. Pasikanti, P. C. Ho and E. C. Y. Chan, *J. Chromatogr. B*, 2008, **871**, 202–211.
- 7 B. Habchi, S. Alves, A. Paris, D. N. Rutledge and E. Rathahao-Paris, *TrAC Trends Anal. Chem.*, 2016, **85**, 128–139.
- 8 J. Han, R. M. Danell, J. R. Patel, D. R. Gumerov, C. O. Scarlett, J. P. Speir, C. E. Parker, I. Rusyn, S. Zeisel and C. H. Borchers, *Metabolomics*, 2008, **4**, 128–140.
- 9 G. Madalinski, E. Godat, S. Alves, D. Lesage, E. Genin, P. Levi, J. Labarre, J. C. Tabet, E. Ezan and C. Junot, *Anal. Chem.*, 2008, **80**, 3291–3303.
- 10 L. Lin, Q. Yu, X. Yan, W. Hang, J. Zheng, J. Xing and B. Huang, *R. Soc. Chem.*, 2010, **135**, 2970–2978.
- 11 P. J. Taylor, *Clin. Biochem.*, 2005, **38**, 328–334.
- 12 E. Rathahao-Paris, S. Alves, N. Boussaid, N. Picard-Hagen, V. Gayraud, P. L. Toutain, J. C. Tabet, D. N. Rutledge and A. Paris, *Eur. J. Mass Spectrom.*, 2018, **25**, 251–258.
- 13 B. Habchi, S. Alves, D. Bouveresse Jouan-rimbaud, B. Moslah, A. Paris, Y. Lécluse, P. Gauduchon, P. Lebailly, D. N. Rutledge and E. Rathahao-Paris, *Metabolomics*, DOI:10.1007/s11306-017-1179-x.
- 14 J. Jansson, B. Willing, M. Lucio, A. Fekete, J. Dicksved, J. Halfvarson, C. Tysk and P. Schmitt-Kopplin, *PLoS One*, 2009, **4**, e6386.
- 15 M. Lucio, A. Fekete, C. Weigert, B. Wägele, X. Zhao, J. Chen, A. Fritsche, H.-U. Häring, E. D. Schleicher, G. Xu, P. Schmitt-Kopplin and R. Lehmann, *PLoS One*, 2010, **5**, e13317.
- 16 A. Oikawa, Y. Nakamura, T. Ogura, A. Kimura, H. Suzuki, N. Sakurai, Y. Shinbo, D. Shibata, S. Kanaya and D. Ohta, *Plant Physiol.*, 2006, **142**, 398–413.
- 17 R. J. Raterink, F. M. van der Kloet, J. Li, N. A. Wattel, M. J. M. Schaaf, H. P. Spaink, R. Berger, R. J. Vreeken and T. Hankemeier, *Metabolomics*, 2013, **9**, 864–873.
- 18 C. C. Jacob, G. Dervilly-Pinel, G. Biancotto, F. Monteau and B. Le Bizec, *Metabolomics*, 2015, **11**, 184–197.
- 19 N. Bonvallet, M. Tremblay-Franco, C. Chevrier, C. Canlet, C. Warembourg, J. P. Cravedi and S. Cordier, *PLoS One*, 2013, **8**, e64433.
- 20 P. Lebailly, C. Vigreux, C. Lechevrel, D. Ledemeney, T. Godard, F. Sichel, J. Y. LeTalaër, M. Henry-Amar and P. Gauduchon, *Cancer Epidemiol. Biomarkers Prev.*, 1998, **7**, 929–940.
- 21 P. Lebailly, A. Devaux, D. Pottier, M. Meo De, V. Andre, I. Baldi, F. Severin, J. Bernaud, B. Durand, M. Henry-Amar and P. Gauduchon, *Occup. Environ. Med.*, 2003, **60**, 910–917.
- 22 J. Trygg and S. Wold, *J. Chemom.*, 2002, **16**, 119–128.
- 23 J.-F. Cardoso, *Neural Comput.*, 1999, **11**, 157–192.
- 24 D. N. Rutledge and D. Jouan-Rimbaud Bouveresse, *Trends Anal. Chem.*, 2013, **50**, 22–32.
- 25 D. N. Rutledge and D. Jouan-Rimbaud Bouveresse, *Trends Anal. Chem.*, 2015, **67**, 220.
- 26 M. B. Wilk and R. Gnanadesikan, *Biometrika*, 1968, **55**, 1–17.
- 27 D. S. Wishart, Y. D. Feunang, A. Marcu, A. C. Guo, K. Liang, R. Vázquez-Fresno, T. Sajed, D. Johnson, C. Li, N. Karu, Z. Sayeeda, E. Lo, N. Assempour, M. Berjanskii, S. Singhal, D. Arndt, Y.

- 1
2
3 Liang, H. Badran, J. Grant, A. Serra-Cayuella, Y. Liu, R. Mandal, V. Neveu, A. Pon, C. Knox, M.
4 Wilson, C. Manach and A. Scalbert, *Nucleic Acids Res.*, 2018, **46**, D608–D617.
5
6 28 M. Kanehisa, S. Goto, S. Kawashima and A. Nakaya, *Nucleic Acids Res.*, 2002, **30**, 42–6.
7
8 29 C. A. Smith, G. O’Maille, E. J. Want, C. Qin, S. A. Trauger, T. R. Brandon, D. E. Custodio, R.
9 Abagyan and G. Siuzdak, *Ther. Drug Monit.*, 2005, **27**, 747–51.
10
11 30 O. Billker, V. Lindo, M. Panico, A. E. Etienne, T. Paxton, A. Dell, M. Rogers, R. E. Sinden and H.
12 R. Morris, *Nature*, 1998, **392**, 289–292.
13
14 31 S. Ganti, S. L. Taylor, K. Kim, C. L. Hoppel, L. Guo, J. Yang, C. Evans and R. H. Weiss, *Int. J.*
15 *Cancer.*, 2012, **130**, 2791–2800.
16
17 32 J. Xu, Y. Chen, R. Zhang, J. He, Y. Song, J. Wang, H. Wang, L. Wang, Q. Zhan and Z. Abliz, *Sci.*
18 *Rep.*, 2016, **6**, 35010.
19
20 33 H. Yoon, *Ann. Pediatr. Endocrinol. Metab.*, 2015, **20**, 119–24.
21
22 34 M. G. Murphy, J. F. S. Crocker, P. O’Regan, S. H. S. Lee, L. Geldenhuys, K. Dooley, M. Al-
23 Khalidi and P. D. Acott, *Chemosphere*, 2007, **68**, 1692–1698.
24
25 35 J. Bremer, *Physiol. Rev.*, 1983, **63**, 1420–1480.
26
27 36 M. Peng, X. Fang, Y. Huang, Y. Cai, C. Liang, R. Lin and L. Liu, *J. Chromatogr. A*, 2013, **1319**,
28 97–106.
29
30 37 J. Klepacki, J. Klawitter, J. Klawitter, J. M. Thurman and U. Christians, *Clin. Chim. Acta*, 2015,
31 **446**, 43–53.
32
33 38 Q. He, X. Kong, G. Wu, P. Ren, H. Tang, F. Hao, R. Huang, T. Li, B. Tan, P. Li, Z. Tang, Y. Yin
34 and Y. Wu, *Amino Acids*, 2009, **37**, 199–208.
35
36 39 N. Çetin, E. Çetin, G. Eraslan and A. Bilgili, *Res. Vet. Sci.*, 2007, **82**, 405–408.
37
38 40 L. Du, H. Wang, W. Xu, Y. Zeng, Y. Hou, Y. Zhang, X. Zhao and C. Sun, *Toxicol. Sci.*, 2013, **134**,
39 195–206.
40
41 41 H.-P. Wang, Y.-J. Liang, Q. Zhang, D.-X. Long, W. Li, L. Li, L. Yang, X.-Z. Yan and Y.-J. Wu,
42 *Pestic. Biochem. Physiol.*, 2011, **101**, 232–239.
43
44 42 J. Yang, X. Sun, Z. Feng, D. Hao, M. Wang, X. Zhao and C. Sun, *Toxicol. Lett.*, 2011, **206**, 306–
45 313.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table and Figure captions

Table 1 Annotation of the most discriminant m/z values and their correlated ions from DIMS dataset.

Fig. 1 Discrimination of control subjects (a) and exposure classes: captan (b), chlorothalonil (c), cyprodinyl (d), and isoproturon (e). Results obtained from DIMS data

1
2
3 processing without the contribution of group information, *i.e.*, based only on the extracted
4 MS source signals. Note that discrimination of epoxiconazol-exposed group is visible in
5 another dimension (**b, c and e**).
6
7

8
9
10 **Fig. 2** 3D-plots obtained from DIMS (**a**) and LC/MS (**b**) data processing without the
11 known group memberships.
12

13
14 **Fig. 3** Box-whisker plots showing intensity changes for two discriminant variables, m/z
15 132.0763 and 138.0546 (from DIMS dataset) between control and all exposure groups.
16 The box represents the first and third quartiles, the bold horizontal lines represent the
17 median intensity. The whiskers correspond to the 1.5 times the interquartile range, and
18 the asterisk represent the outlier.
19
20
21
22
23
24
25

26 **Fig. 4** Zoomed mass spectrum (in the m/z 60-200 range) reconstituted from correlation
27 coefficients calculated from DIMS dataset of a urine sample collected on a subject
28 exposed to chlorothalonil. Color coding of the signal points is based on their correlation
29 with the discriminant variables at m/z 76.075. The peaks in red were strongly correlated.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1.

Observed peak (m/z)	Proposed ion elemental composition	Putative ion annotation ^(a)		Isoprotunron **	Captan	Chlorothalonil	Cyprodinil	Epoxiconazol
		Proposed species	Proposed compound name					
76.0753	C ₃ H ₁₀ NO	[M+H] ⁺	Trimethylamine N-oxide	-	-	↗	-	↘
98.0573	C ₃ H ₉ NONa	[M+Na] ⁺		↘	-	-	↘	↘
151.1438	C ₆ H ₁₉ N ₂ O ₂	[2M+H] ⁺		-	-	↗	↘	↘
152.1471 ^(b)	¹² C ₅ ¹³ CH ₁₉ N ₂ O ₂	[2M+H, ¹³ C] ⁺		↘	-	-	↘	↘
136.1077	C ₄ H ₁₀ N ₂ O ₂ NH ₄	[M+NH ₄] ⁺	Urea, N-hydroxy-N-propyl					
189.1345	C ₇ H ₁₇ N ₄ O ₂	[M+H] ⁺	Homo-arginine	-	↘	-	-	-
132.0763	C ₄ H ₁₀ N ₃ O ₂	[M+H] ⁺	Creatine	↘	↗	↘	↘	↘
110.0288				↗	↘	-	↗	-
138.0547	C ₇ H ₈ NO ₂	[M+H] ⁺	Trigonelline	↗	↘	-	↗	-
140.0452	C ₅ H ₆ N ₃ O ₂	[M+H] ⁺	Hydroxypyrazinamide					
144.1016	C ₇ H ₁₄ NO ₂	[M+H] ⁺	Proline betaine	-	-	-	↗	↘
145.1049 ^(b)	¹² C ₆ ¹³ CH ₁₄ NO ₂	[M+H, ¹³ C] ⁺						
229.1545	C ₁₁ H ₂₁ N ₂ O ₃	[M+H] ⁺	Leucyl-Proline	-	-	-	↗	-
230.1578 ^(b)	¹² C ₁₀ ¹³ CH ₂₁ N ₂ O ₃	[M+H, ¹³ C] ⁺						
241.1543	C ₁₂ H ₂₁ N ₂ O ₃	[M+H] ⁺	Pirbuterol					
115.0626 ^(b)	C ₄ H ₈ ¹⁴ N ₂ ¹⁵ N ₁ O	[(M+H), ¹⁵ N] ⁺	Creatinine					
115.0689 ^(b)	¹² C ₃ ¹³ C H ₈ N ₃ O	[(M+H), ¹³ C] ⁺						
227.125	C ₈ H ₁₅ N ₆ O ₂	[2M+H] ⁺						
228.1280 ^(b)	¹² C ₇ ¹³ CH ₁₅ N ₆ O ₂	[2M+H, ¹³ C] ⁺		-	↘	-	-	-
249.1069	C ₈ H ₁₄ N ₆ O ₂ Na	[2M+Na] ⁺						
362.1659	C ₁₂ H ₂₁ N ₉ O ₃ Na	[3M+Na] ⁺						
162.1122	C ₇ H ₁₆ NO ₃	[M+H] ⁺	Carnitine	↘	-	↘	↘	↗

163.1157 ^(b)	¹² C ₆ ¹³ CH ₁₆ NO ₃	[M+H, ¹³ C] ⁺				
146.1171	C ₇ H ₁₆ NO ₂	[M+H] ⁺	Dehydroxycarnitine			
204.1229	C ₉ H ₁₈ NO ₄	[M+H] ⁺	Acetylcarnitine	- ↗	- ↗	-
118.0605	C ₃ H ₈ N ₃ O ₂	[M+H] ⁺	Guanidoacetic acid	↘ -	- ↘	-
257.1604	C ₁₁ H ₂₂ N ₄ O ₄	[M+H-H ₂ O] ⁺	Glutaminy-Lysine	- ↘	↗ -	-

* Changes between each exposed group and their control one: -, no change; ↗, increase; ↘, decrease.

Horizontal lines separate each set of correlated ions; bold m/z values correspond to the selected discriminant variables; the non-bold m/z values are correlated variables with the first one (in bold). ^(a) Note that only one identity was assigned for each variable among the different isomers proposed from DB queries. ^(b) Proposed ¹³C and ¹⁵N isotopic peaks. Correlated ions without DB hit were eliminated from the list.






Fig. 1 Discrimination of control subjects (a) and exposure classes: captan (b), chlorothalonil (c), cyprodinyl (d), and isoprotruron (e). Results obtained from DIMS data processing without the contribution of group information, i.e., based only on the extracted MS source signals. Note that discrimination of epoxiconazole-exposed group is visible in another dimension (b, c and e).

212x150mm (300 x 300 DPI)





(a)




(b)

Ca captan Ch chlorothalonil Cy cyprodinyl E epoxiconazol I isoprotruron T controls



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43



Supplementary Material

Evaluation of direct introduction-high resolution mass spectrometry (DI-HRMS) based metabolomics approach for the discrimination of several exposure groups to pesticides: comparison with liquid chromatography high resolution mass spectrometry (LC/HRMS)

Baninia Habchi^{1,2}, Sandra Alves^{2*}, Delphine Jouan-Rimbaud Bouveresse^{1,3}, Bilel Moslah¹, Alain Paris⁴, Yannick Lécluse⁵, Pascal Gauduchon⁵, Pierre Lebailly⁵, Douglas N. Rutledge¹, Estelle Rathahao-Paris^{1,2*}

¹ UMR Ingénierie Procédés Aliments, AgroParisTech, Inra, Université Paris-Saclay, 91300 Massy, France

² Sorbonne Université, Faculté des Sciences et de l'Ingénierie, Institut Parisien de Chimie Moléculaire (IPCM), F-75005 Paris, France

³ UMR 914 Physiologie de la Nutrition et du Comportement Alimentaire, INRA, AgroParisTech, Université Paris-Saclay, F-75005 Paris

⁴ Muséum national d'Histoire naturelle, MCAM, UMR7245 CNRS - MNHN, 75005 Paris, France

⁵ ANTICIPE U1086 INSERM & Université Caen-Normandie, Centre François Baclesse, 14076 Caen Cedex 05 France

Supplementary material section contains LC/MS and MS/MS experimental conditions. Additional results obtained from both DIMS and LC/MS data are presented in Fig. S-1 to S-5. Table S-1 showed the most discriminant variables and their correlated ions putatively annotated from LC/MS data.

*Corresponding authors :

Dr. Estelle Rathahao-Paris

Present address: UMR CEA-INRA Service de Pharmacologie et d'Immunoanalyse, Laboratoire d'Immuno-Allergie Alimentaire, CEA de Saclay- Bat 136, F-91191, Gif-sur-Yvette cedex, FRANCE. E-mail: Estelle.Paris@inra.fr

or

Dr. Sandra Alves

Address: Sorbonne Université, Faculté des Sciences et de l'Ingénierie, Tour 42-43, 4^{ème} étage, BP 45, 4 place Jussieu F-75005, Paris, FRANCE. E-mail: sandra.alves@sorbonne-universite.fr

LC/MS experimental procedure

1. Mass spectrometry detection

LC/MS experiments were acquired in the m/z 100-1000 range with a cycle time of 14 min (including column re-equilibration) using a mass resolving power of 100,000 FWHM (Full width at half maximum) at m/z 400. The following ESI conditions were applied: electrospray voltage was set at -3 kV, capillary voltage at 3 V, capillary temperature at 275 °C and the tube lens offset at 80 V, sheath gas flow (nitrogen) and auxiliary gas were fixed at 66 and 30 arbitrary units (a.u.), respectively.

The chromatographic separation was performed using a C₁₈ NucleodurHTec column (2 × 100 mm, 3 μm, Macherey-Nagel, Eurl, Hoerd, France) and a high performance liquid chromatography system, Surveyor LC (Thermo Fisher Scientific). 10 μL of sample diluted 10 fold in a mixture of acetonitrile/water (1:1, v/v) were injected at a flow rate of 200 μL/min. The mobile phase consisted of water plus 0.1 % formic acid (A) and acetonitrile plus 0.1% formic acid (B). The gradient started with 2% (B) for 1 min, and increased to 100% (B) in 7 min, then was maintained at 100% (B) during 2 min.

2. Data pre-processing

Raw LC/MS data were converted from profile into centroid mzXML format using MS Convert (<http://proteowizard.sourceforge.net/tools.shtml>). XCMS based on a program written in R (*R version 3.4.0, The R Foundation for Statistical Computing, Vienna 2017, <https://www.r-project.org>*) was used to pre-process generated data. Four basic steps corresponding to the peak picking, the peak grouping, the retention time correction, and a second peak grouping were performed. The *centWave* method was used for peak extraction, followed by the correction of retention time drift using a non-linear LOESS alignment method (*Smith C. A. et al., XCMS: Processing Mass Spectrometry Data for Metabolite Profiling Using Nonlinear Peak Alignment, Matching, and Identification.2006, Anal Chem 78:779–787*). Finally, the fill peaks method was applied to find peak intensity in the raw data to correct missing values. The final number of extracted variables from the 82 samples was 2106.

MS/MS experiments

Tandem mass spectrometry (MS/MS) experiments were performed on the monoisotopic peaks of some selected metabolites using CID (collision-induced dissociation) and/or HCD (higher collision dissociation) conditions. The following parameters were used: isolation width of precursor ions of 1 to 2 u, activation time of 30 ms, and normalized collision energy of about 20 % and 38 % (arbitrary units) for CID and HCD, respectively. The precursor ions were selected, fragmented in the linear ion trap (LTQ) device and final high-resolution detection with a mass resolving power of 7500 (FWHM) for m/z 400 was performed.




Fig.S.1 Results obtained from the DIMS data processing with known group memberships. Discrimination of control subjects (a) and the four exposure classes: captan (b), chlorothalonil (c), cyprodinyl (d) and isoproturon (e). The corresponding extracted source signals show the m/z values which most contribute to group discrimination (f, g, h, i and j). The peak intensity represents the ion contribution to the corresponding IC. The part of the source signals belonging to the group membership matrix show which group(s) contribute(s) the most to build the IC when compared to the influence of the others (k, l, m, n and o). Note that no discrimination is observed for epoxiconazole-exposed subjects.




Fig.S.2 Results obtained from LC/MS data processing with known group memberships. Discrimination of control subjects (a) and all exposure classes: captan (b), chlorothalonil (c), cyprodinyl and epoxiconazol (d) and isoprotruron (e). The corresponding extracted source signals show the m/z values which most contribute to group discrimination (f, g, h, i and j plots). The peak intensity represents the ion contribution to the corresponding IC. The part of the source signals belonging to the group membership matrix show which group(s) contribute(s) the most to build the IC when compared to the influence of the others (k, l, m, n and o).




Fig.S.3 Results from LC/MS data processing without the contribution of group information, i.e., based on the only extracted MS source signals. A separation is still observed for captan, chlorothalonil, cyprodinyl and isoprotruron exposure groups (**b-e** respectively). A discrimination of control subjects (**a**) is less apparent and no discrimination of epoxiconazol exposure group is obtained.




Fig.S.4 3D-plot from DIMS (a) and LC/MS (b) data processing without the contribution of group information. Discrimination of different exposure groups is observed, especially, separation of individuals exposed to isoproturon according to the proportions of IC-5. Cyprodinyl exposure group was discriminated according to another dimension as observed in **Fig.2**.




Fig.S.5 MS/MS spectra of the precursor $[M + H]^+$ ions at m/z 204 (a) from the authentic standard acetylcarnitine and (b) from a urine sample.

Table S.1 The most discriminant m/z values extracted from LC/MS data

Observed peak(m/z)	Proposed ion elemental composition	Putative ion annotation ^(a)		Isoproturon	Captan	Chlorothalonil	Cyprodinil	Epoxiconazol
		Proposed species	Proposed compound name					
102.0524; 736	C ₂ H ₉ NO ₂ Na	[M+Na] ⁺	N-Hydroxy-N-methylmethanamine-water	-	-	-	-	↘
102.0823; 771				-	-	↘	-	↘
102.1069; 549								
271.1515; 776	C ₁₀ H ₁₉ N ₆ O ₃ C ₁₀ H ₁₉ N ₆ O ₃	[M+H-H ₂ O] ⁺ [M+NH ₄] ⁺	GlyGlyArg 3-[2-(6-Amino-9H-purin-9-yl)ethoxy]propane-1,2-diol					
250.3262; 798								
381.2792; 795	C ₂₆ H ₃₇ O ₂	[M+H] ⁺	Benzoic acid 2-[(4-dodecylphenyl)methyl]					
102.1271; 750						↗*		
102.1402; 694				-	↗	-	-	-
102.1406; 805				-	↗	-	-	-
102.1069; 549								
102.1481; 760				-	↗	-	↗	-
102.1054; 518								
102.1483; 723				-	-	-	-	↘
114.0595; 591				-	-	-	↗	-
114.0896; 589	C ₄ H ₁₀ N ₄	[M+NH ₄] ⁺	Methyl-triazin-ium	-	↗	-	↗	-
121.1058; 103				-	↘	-	-	-
136.0478; 63	C ₄ H ₇ N ₃ ONa	[M+Na] ⁺	Creatinine	-	-	↘	-	-
141.1131; 33	C ₆ H ₁₃ N ₄	[M+H] ⁺	1-(2-Methylbutan-2-yl)-1H-tetrazole	-	-	-	-	↘
150.1447; 75				-	↘	-	↘	↗
118.5554; 94								
118.6349; 74								
118.6495; 306								
150.1702; 77								
150.1998; 74								
195.9931; 306								
196.0355; 53	C ₇ H ₆ N ₃ O ₄	[M+H] ⁺	1,2,4-Oxadiazole, 5-methyl-3-(5-nitro-2-furanyl)					
196.0366; 136	C ₁₀ H ₇ NO ₂ Na C ₉ H ₈ O ₅	[M+Na] ⁺ [M+H] ⁺	Quinaldic acid 3-Carboxy-4-(methoxycarbonyl)phenolate					
384.2627; 74	C ₂₃ H ₃₄ N ₃ O ₂ C ₁₈ H ₃₄ N ₅ O ₄	[M+H] ⁺ [M+H-H ₂ O] ⁺	N-Dodecyl-2-(1H-imidazole-carbonyl)benzamide 2AlaAla Lys Leu-	-	↘	-	-	-
170.0410; 50	C ₈ H ₆ F ₂ NO	[M+H] ⁺	3,5-Difluoro-4-(hydroxymethyl)benzotrile	-	-	↗	-	-
103.5474; 53								
129.0141; 50	C ₄ H ₁₀ SK C ₉ H ₂ F	[M+K] ⁺ [M+H-2H ₂ O] ⁺	1-Butanethiol Fluorochromone					
153.0326; 50	C ₃ H ₉ N ₂ O ₃ S	[M+H] ⁺	(Methylamino)(methylimino)met					

				hanesulphonic acid					
189.0468; 50	C ₈ H ₁₀ N ₂ S ₁ Na	[M+Na] ⁺		Ethionamide					
170.0447; 206	C ₇ H ₈ NO ₄	[M+H] ⁺		Furoylglycine				↗*	
180.0675; 484	C ₇ H ₇ FN ₅	[M+H] ⁺		4-Fluoro-3-(1H-tetrazol-1-yl)aniline	↗	-	↗	-	-
180.1005; 480	C ₈ H ₁₅ NO ₂ Na	[M+Na] ⁺		Homostachydrine	-	↘	-	↗	-
181.0608; 447	C ₈ H ₉ N ₂ O ₃	[M+H] ⁺		Nicotinic acid	-	↘	-	-	-
190.0498; 480	C ₁₀ H ₈ NO ₃	[M+H] ⁺		Kynurenic acid					
206.0448; 463	C ₁₀ H ₈ NO ₄	[M+H] ⁺		Xanthurenic acid					
181.0687; 484^(b)	C ₈ ¹³ CH ₁₀ NO ₃	[M+H, ¹³ C] ⁺		Hippuric acid				↘*	
105.0154; 483 ^(b)	C ₇ H ₅ O	[M+H-C ₂ H ₅ NO ₂] ⁺							
179.0188; 480	C ₆ H ₁₀ O ₂ S ₂	[M+H] ⁺		Bissulfine					
181.1212; 480	C ₉ H ₁₅ N ₃ O	[M+H-2H ₂ O] ⁺		Maleic hydrazide choline salt	-	-	↗	-	-
205.0972; 267	C ₁₁ H ₁₃ N ₂ O ₂	[M+H] ⁺		Tryptophan	-	↘	-	-	-
242.9254; 60					-	-	↘	-	-
174.938; 60									
310.9128; 60									
446.8873; 60									
265.1220; 477	C ₁₈ H ₁₇ O ₂	[M+H] ⁺		Cinnamylcinnamate	-	↘	-	-	↘
274.122; 390	C ₁₉ H ₁₆ NO	[M+H] ⁺		Methoxyphenylcarbazole	-	-	-	-	↘
	C ₁₁ H ₂₀ N ₃ O ₃ S	[M+H-H ₂ O] ⁺		CysLeuGly					
114.0497; 463									
114.0569; 459	C ₄ H ₁₂ PNa	[M+Na] ⁺		Phosphonium, tetramethyl					
299.1828; 750	C ₁₂ H ₂₃ N ₆ O ₃	[M+H-H ₂ O] ⁺		Ala AlaArg	-	↘	-	-	-
310.2011; 497	C ₁₇ H ₂₈ NO ₄	[M+H] ⁺		Decadienylcarnitine				↘*	
311.2044; 497 ^(b)	¹² C ₁₆ ¹³ CH ₂₈ NO ₄	[M+H, ¹³ C] ⁺							
304.4917; 497									
286.1716; 528	C ₁₈ H ₂₅ NP	[M+H] ⁺		Ethanamine, 2-(diphenylphosphino)-N,N-diethyl					
326.0869; 433	C ₁₄ H ₁₆ NO ₈	[M+H] ⁺		Dihydroxy-1H-indole glucuronide I	-	↘	-	-	-
348.0687; 433	C ₁₄ H ₁₅ NO ₈ Na	[M+Na] ⁺							
464.1909; 528	C ₂₃ H ₃₀ NO ₉	[M+H] ⁺		Dihydroisomorphine-glucuronide					
330.2273; 487	C ₁₇ H ₃₂ NO ₅	[M+H] ⁺		Keto-decanoylcarnitine	-	-	-	↘	-
331.2306; 487 ^(b)	¹² C ₁₆ ¹³ CH ₃₂ NO ₅	[M+H, ¹³ C] ⁺							
374.2533; 490	C ₁₉ H ₃₆ NO ₆	[M+H] ⁺		Dodecanedioylcarnitine					
229.1433; 514	C ₁₂ H ₂₁ O ₄	[M+H] ⁺		Traumatic acid					
269.1494; 504	C ₁₁ H ₂₅ N ₃ O ₂	[M+K] ⁺		Diacetylspermidine					
	C ₁₃ H ₂₁ N ₂ O ₄	[M+H] ⁺		Dihydroxymelphalan					
356.2429; 507	C ₁₉ H ₃₃ NO ₅	[M+H] ⁺		Derivative of carnitine					

* Changes between each exposed group and their control one: -, no change; ↗, increase; ↘, decrease; change between all exposed groups and all control: ↗*, increase; ↘*, decrease. Horizontal lines separate each set of correlated ions; bold m/z value correspond to the most discriminant variables; the non-bold m/z values are correlated variables with the first one (in bold). ^(a) Note that only one identity was assigned for each variable among the different isomers proposed from DB queries. ^(b) Proposed ion fragment composition or ¹³C isotopic peak. Correlated ions without DB hit were eliminated. Some candidates considered unreliable were rejected in the absence of isotopic peaks on the mass spectrum (e.g., absence of the ³⁷Cl isotopic peak in the mass spectrum since its expected abundance should correspond to 30% and 60% for one and two chlorine atoms, respectively, compared to the monoisotopic peak).