



**HAL**  
open science

## Improved analysis of arabinoxylan-bound hydroxycinnamate conjugates in grass cell walls

Alexis Eugene, Catherine Lapierre, John Ralph

### ► To cite this version:

Alexis Eugene, Catherine Lapierre, John Ralph. Improved analysis of arabinoxylan-bound hydroxycinnamate conjugates in grass cell walls. *Biotechnology for Biofuels*, 2020, 13, 10.1186/s13068-020-01841-6 . hal-03083492

**HAL Id: hal-03083492**

**<https://agroparistech.hal.science/hal-03083492>**

Submitted on 19 Dec 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

METHODOLOGY

Open Access



# Improved analysis of arabinoxylan-bound hydroxycinnamate conjugates in grass cell walls

Alexis Eugene<sup>1</sup> , Catherine Lapierre<sup>2</sup> and John Ralph<sup>1,3\*</sup>

## Abstract

**Background:** Arabinoxylan in grass cell walls is acylated to varying extents by ferulate and *p*-coumarate at the 5-hydroxy position of arabinosyl residues branching off the xylan backbone. Some of these hydroxycinnamate units may then become involved in cell wall radical coupling reactions, resulting in ether and other linkages amongst themselves or to monolignols or oligolignols, thereby crosslinking arabinoxylan chains with each other and/or with lignin polymers. This crosslinking is assumed to increase the strength of the cell wall, and impedes the utilization of grass biomass in natural and industrial processes. A method for quantifying the degree of acylation in various grass tissues is, therefore, essential. We sought to reduce the incidence of hydroxycinnamate ester hydrolysis in our recently introduced method by utilizing more anhydrous conditions.

**Results:** The improved methanolysis method minimizes the undesirable ester-cleavage of arabinose from ferulate and *p*-coumarate esters, and from diferulate dehydrodimers, and produces more methanolysis vs. hydrolysis of xylan-arabinosides, improving the yields of the desired feruloylated and *p*-coumaroylated methyl arabinosides and their diferulate analogs. Free ferulate and *p*-coumarate produced by ester-cleavage were reduced by 78% and 68%, respectively, and 21% and 39% more feruloyl and *p*-coumaroyl methyl arabinosides were detected in the more anhydrous method. The new protocol resulted in an estimated 56% less combined diferulate isomers in which only one acylated arabinosyl unit remained, and 170% more combined diferulate isomers conjugated to two arabinosyl units.

**Conclusions:** Overall, the new protocol for mild acidolysis of grass cell walls is both recovering more ferulate- and *p*-coumarate-arabinose conjugates from the arabinoxylan and cleaving less of them down to free ferulic acid, *p*-coumaric acid, and dehydrodiferulates with just one arabinosyl ester. This cleaner method, especially when coupled with the orthogonal method for measuring monolignol hydroxycinnamate conjugates that have been incorporated into lignin, provides an enhanced tool to measure the extent of crosslinking in grass arabinoxylan chains, assisting in identification of useful grasses for biomass applications.

**Keywords:** Ferulate, *p*-Coumarate, Acylation, Crosslinking, Dehydrodiferulates, Methanolysis

## Background

Hydroxycinnamate conjugates in the cell wall have been the subject of increasing interest. *p*-Coumarate (*p*CA), especially, has been long known to acylate lignins and has been shown to arise on lignin polymers via

polymerization of pre-formed monolignol *p*CA conjugates, along with the canonical monolignols [1–3]. Transferase genes have now been identified for this monolignol acylation via the activated intermediate *p*-coumaroyl-CoA [4–6]; the equivalent alternative viewpoint is that *p*CA (via *p*-coumaroyl-CoA) is esterified by monolignols. More recently, plants have been engineered to produce analogous monolignol ferulate (FA) conjugates [3, 7, 8], that also participate in lignification and, because ferulate is more compatible with the monolignols in its radical

\*Correspondence: jralph@wisc.edu

<sup>1</sup> Great Lakes Bioenergy Research Center, Wisconsin Energy Institute,

University of Wisconsin, Madison, WI, USA

Full list of author information is available at the end of the article




coupling reactions [2, 9], biosynthesize lignins with readily cleavable ester bonds in the lignin backbone, allowing more facile delignification of such materials [8, 10–12]. The DFRC (derivatization by reductive cleavage) method, which cleaves lignin  $\beta$ -ethers while leaving  $\gamma$ -esters intact, was extended to provide unambiguous evidence for the existence of such conjugates in lignin, and to assess their (relative) levels in the polymer [7, 13–17]; DFRC releases both *p*CA and FA conjugates (and in fact also benzoate, vanillate, and other conjugates) that are involved in  $\beta$ -ether units in lignin that can be cleaved by the method [13, 18–20]. The availability of the method led to the discovery that various plants naturally use monolignol FA (ML-FA) conjugates in their lignification [15].

FA and *p*CA esters have also long been known on arabinoxylans in grasses, and commelinid monocots in general [2, 21–29], Fig. 1. Both, but particularly FA-polysaccharide esters, have been demonstrated to be involved

in radical coupling reactions to cross-link cell wall polysaccharides, both with other hydroxycinnamate-bearing polysaccharide chains and with lignin. The genes/enzymes involved in arabinoxylan acylation have been more difficult to elucidate and the actual nature of the acceptor (poly)saccharide moiety remains elusive; a gene for *p*-coumaroylation is reasonably certain [30, 31], but the ferulate analog remains putative [32].

Having orthogonal analytical methods to delineate between hydroxycinnamates that acylate monolignols vs. polysaccharides is particularly beneficial for research in this area. We recently introduced a method to clip arabinosyl units from arabinoxylan polysaccharides releasing hydroxycinnamoylated methyl arabinosides for analysis [33], Fig. 1. The 1-*O*-methyl arabinosides vs. the parent (1-*O*-H) arabinosides themselves, allow for improved HPLC producing sharper chromatograms than the original method using




aqueous HCl [34], presumably because the  $\alpha$ - and  $\beta$ -anomers of the parent arabinose could interconvert during chromatography, whereas the methyl arabinosides are stable. The major species of interest identified with this method are methyl 5-*O*-feruloyl arabinofuranoside (FA-MeAra **4**) and methyl 5-*O*-*p*-coumaroyl arabinofuranoside (*p*CA-MeAra **3**), as well as various dehydrodiferulates (DFA) **6** and ferulate-monolignol cross-products **7** in which the ferulate moieties are esterified as 1-*O*-methyl arabinosides [33, 34], Fig. 1.

The mild acidolysis procedure provides a simple, straightforward way of determining the relative acylation of grass arabinoxylan, but we came to realize that the presence of  $\sim 10\%$  water in the acidolysis reagent leads to some undesirable hydrolysis that cleaves the esters and results in small amounts of arabinose rather than 1-*O*-methyl-arabinofuranoside units. As a result, some monomeric free hydroxycinnamic acids are produced (where as free acids are essentially not present in the cell wall) along with a selection of mono-methyl-arabinosylated diferulates that complicate the chromatogram. Here, we describe an improved ‘anhydrous’ method that limits such hydrolytic reactions, while still being able to cleave the glycoside linking arabinose units to the xylan backbone, Fig. 1.

## Results and discussion

As described in “Methods” section, the new procedure utilizes anhydrous methanolic HCl (with dioxane as a co-solvent) to limit aqueous hydrolysis of both the hydroxycinnamate ester and the arabinosyl unit from the xylan backbone. The methanolic-HCl reagent is conveniently prepared from commercially available anhydrous methanol (and dioxane) by carefully adding acetyl chloride in the simple method described by Fieser and Fieser [35]. We admit that the conditions are not strictly anhydrous as the biomass sample comes with its own moisture and exchangeable hydroxyls. We were originally not certain that such conditions would continue to efficiently clip off the arabinose units, fearing that some water might be necessary; that trepidation appeared to be groundless, however, as methanolysis was found to be essentially as facile as hydrolysis. We tested whether yields could be improved by adding the acetyl chloride directly to samples already suspended in anhydrous methanol/dioxane; however, as the FA-MeAra **4** yield only increased by about 2%, we contend that the simplicity of using the pre-prepared hydrolysis solution outweighs any slight yield benefit that might be realized.


Figure 2 shows the UV chromatograms for a sample of maize insoluble dietary fiber (IDF) that underwent both the normal (black) and anhydrous (red) acidolysis processes. The main products of mild acidolysis are



**Fig. 2** UV-vis chromatograms of maize IDF after mild acidolysis. Black trace was processed using the original acidolysis method; red trace was the new anhydrous method described herein. Internal standard,  $I_{std}$ ; ferulate **1**; FA-Ara **2** (unmethylated,  $m/z$  325); *p*CA-MeAra **3**, FA-MeAra **4**; MeFA, methyl ferulate, **5**; DFAs **6** (one isomer, the 8-*O*-4-DFA, is shown) with two or one (**6'**) MeAra attached, also shown vertically expanded  $\times 3$ ; monolignol-FA-MeAra coupling products **7** (Fig. 1) are also in this region

marked 1–7. Peak 1 is free FA, formed from the hydrolysis of arabinose-bound ferulate. This peak is significantly decreased when mild acidolysis is performed under the more anhydrous conditions, indicating that less hydrolysis of FA-MeAra is occurring. *p*-Coumaric acid (*p*CA, eluting at 4.8 min) is also reduced, although the peak is too small to see in this chromatogram. Peak 2 appears only in the original method and is due to unmethylated FA-Ara (*m/z* 325); similar small peaks occur throughout the chromatogram for the (partially) unmethylated analogs of all the possible products when aqueous HCl is used. Performing the acidolysis in the near-absence of water clearly aids in the methanolysis (rather than the competing hydrolysis) of the arabinosyl units, simplifying the array of products formed. Peaks 3 and 4 are *p*CA-MeAra and FA-MeAra, typically the most abundant desired products from mild acidolysis of grasses. The peak at 14.5 min (with an apparent mass of 472, that is more abundant in the older method) remains unidentified at this time. Peak 5 is due to methyl ferulate. The elevation of peak 5 (and also methyl *p*-coumarate, not shown) suggests that more ferulate is clipped from its arabinose unit but peak 4 is also enhanced under the more anhydrous acidolysis conditions, so the method remains an improvement for the methyl-arabinosylated compounds of most interest. Finally, the grouped region in Fig. 2 is where a series of different DFAs 6 conjugated to either two or one (denoted as 6') methyl arabinoside units elutes. Monolignol ferulate (ML-FA) cross-products 7, Fig. 1, are also found in this region. The presence of these species is direct evidence of the crosslinking nature of arabinose-bound ferulate. It is clear from the UV chromatograms that there are again more of these products recovered with the dry method (red chromatogram).


Figure 3 shows the key differences between wet and anhydrous acidolysis in more detail. Panel a displays the extracted-ion chromatogram (EIC) for *m/z* 193 from the samples in Fig. 2, in which *Z*- and *E*-ferulic acid 1 elute at 7.0 and 7.5 min; both are reduced (as noted above) indicating that less ester hydrolysis has taken place. Panel b is the EIC for *m/z* 339, corresponding to FA-MeAra 4, which elutes at 13.1 min; the level of this crucial marker compound (for ferulates on arabinoxylan) is elevated. Similarly, panels c and d represent related compounds at various stages of hydrolysis. Figure 3c presents the EIC for DFA-MeAra 6' at *m/z* 531; all of the mono-MeAra DFAs are reduced, again indicating that less arabinose hydrolysis has occurred. Finally, panel d shows the EIC for *m/z* 677, which is DFA-MeAra<sub>2</sub> 6; these crucial peaks are all substantially elevated, again because less ester hydrolysis, and more methanolysis and less hydrolysis of the arabinosyl glycosidic position, has occurred. There are many different isomers of DFA arabinosides



**Fig. 3** Extracted-ion chromatograms. Extracted-ion chromatograms (EIC) are shown for *m/z* 193, 339, 531, and 677 from the samples in Fig. 2. Black trace was processed using the original acidolysis method; red trace was the new anhydrous method. **a, c** Show that levels of the hydrolyzed products FA and DFA-MeAra are reduced under the drier acidolysis conditions. **b, d** Show that the unhydrolyzed esters with fully methanolized Ara units, FA-MeAra 4 and DFA-MeAra<sub>2</sub> 6, are enhanced in the anhydrous method

6 possible as identified and discussed previously [33], accounting for the plethora of peaks in this chromatogram. An estimate of the yield changes from using anhydrous conditions for acidolysis is presented in Fig. 4, clearly showing how the peaks of interest are enhanced over those resulting from hydrolysis.

The aim of this paper was to present the improved method for the analysis of hydroxycinnamates, monomeric and dimeric, on grass arabinoxylans. The method minimizes the undesirable ester-cleavage of arabinose from ferulate and *p*-coumarate esters, and from diferulate dehydrodimers, and produces more methanolysis vs. hydrolysis of xylan-arabinosides, improving the yields of the desired feruloylated and *p*-coumaroylated methyl arabinosides and their diferulate analogs. Free ferulate and *p*-coumarate produced by ester-cleavage were reduced by 78% and 68%, respectively, and 21% and 39% more feruloyl and *p*-coumaroyl methyl arabinosides (4



**Table 1** Amounts of methyl hydroxycinnamoyl arabinosides released from selected monocot samples

Sample	pCA-MeAra 3 (mg/g)	FA-MeAra 4 (mg/g)
Sorghum	2.51 ± 0.57	11.22 ± 0.73
Switchgrass	7.83 ± 0.39	8.96 ± 0.41
Maize IDF	3.62 ± 0.26	33.10 ± 1.71
Rye IDF	0.81 ± 0.05	19.08 ± 1.06

Analyzed using the improved method described in “Methods” section. Values are means of duplicate runs ± the standard deviation

and 3) were detected in the more anhydrous method. The new protocol resulted in an estimated 56% less combined diferulate isomers 6' in which only one acylated arabinosyl unit remained, and 170% more combined diferulate isomers 6 conjugated to two arabinosyl units. The results of quantifying the monomeric methyl hydroxycinnamoyl arabinosides 3 and 4 from a few grass samples, using the new method, are given in Table 1. This protocol will be used in our laboratories from this point on and is recommended to other research groups pursuing such analyses.

## Conclusions

Overall, mild acidolysis is a powerful tool for analyzing the relative acylation of grass arabinoxylans by pCA and FA. Although not all peaks resulting from either hydrolysis of the ester or of the arabinosyl unit from the xylan were eliminated, their levels were significantly reduced. For the analysis of polysaccharide-bound pCA and FA, the levels of the methyl arabinosides 3 and 4 were elevated. In addition, for the DFAs that are most important for evaluation of the extent of polysaccharide crosslinking in grasses, the mono-methyl-arabinosylated DFAs (DFA-MeAra 6') were reduced, and the fully di-methyl-arabinosylated DFAs (DFA-MeAra<sub>2</sub> 6) were significantly

elevated. Because of the higher yields of desirable products and the simplification of the chromatograms from the significant elimination of hydrolysis (vs. methanolysis) products, we highly recommend this method for determining the (relative) levels of hydroxycinnamates and their dimers (and monolignol cross-products) on arabinoxylans, and for its use with the complimentary DFRC-based method to independently analyze lignin-bound hydroxycinnamates.

## Methods

### Chemicals

ACS grade dioxane (Sigma-Aldrich) was distilled immediately prior to use. Anhydrous methanol (Alfa Aesar, 99.9%, packed under argon with septum) was used as is. Aqueous 2 M HCl was made from concentrated HCl (Sigma-Aldrich, 37%) in ultrapure water (18.2 MΩ cm<sup>-1</sup>, Thermochemical E-pure). Acetyl chloride (98%) was from Sigma-Aldrich. FA (Aldrich, 99%), pCA (Sigma, 98%), and *m*-coumaric acid (Acros, 99%) were used as is. FA-MeAra, and pCA-MeAra were synthesized as previously described [33, 36, 37]. Chromatography was performed with LC-MS grade 0.1% formic acid and acetonitrile obtained from Fisher Scientific.

### Plant materials

Insoluble dietary fiber (IDF) from maize and rye was obtained as described previously [38]. Briefly, maize or rye bran was ground to <0.5 mm, defatted with *n*-hexane, and treated with α-amylase. The residue was extracted several times with water and ethanol, and freeze dried for 48 h. Energy sorghum and switchgrass were supplied by the Great Lakes Bioenergy Research Center's material production chain from the 2019 harvest year. Whole-cell-wall samples were pre-ground to <0.5 mm and then extracted three times with water and ethanol, and once with acetone before being freeze dried for 48 h; the product is generally termed the 'cell wall residue' or CWR, or in the food arena as IDF. The dried extract-free IDF material was then ball milled.

### Mild acidolysis

Aqueous mild acidolysis ('normal acidolysis') was performed as described previously [33]. In a glass vial (Chemglass Life Sciences, 2 dram), 10 mg of extract-free plant material was mixed into 1 mL of aqueous acidolysis reagent consisting of dioxane, methanol, and 2 M HCl<sub>(aq)</sub> in a 60/30/10 (v/v) ratio. The vial was capped and heated at 80 °C for 3 h with stirring. Anhydrous mild acidolysis was performed as described above except the samples were incubated with 1 mL of a dry acidolysis reagent. This reagent was prepared by carefully adding 0.7 mL of acetyl chloride to 20 mL of 70/30 (v/v) freshly distilled

dioxane/absolute methanol, following the Fieser and Fieser method for making anhydrous methanolic HCl [35]. As acetyl chloride is moisture sensitive, the dioxane and methanol were kept as dry as possible (using only syringe extraction through the septum), to prevent hydrolysis. After reaction, 100  $\mu$ L of 30 mM *m*-coumaric acid was added as internal standard. Then,  $\sim$ 2 mL of water was added and the mixture was extracted with three portions of ethyl acetate. The organic layers were combined and dried over anhydrous magnesium sulfate, gravity-filtered (VWR 415 filter paper), and the solvent evaporated to dryness on a rotavap. The residue was dissolved in 1.0 mL of acetonitrile and syringe-filtered (0.2  $\mu$ m) before liquid chromatography–mass spectrometric (LC–MS) analysis.

### LC–MS analysis

The samples were analyzed on a Shimadzu Nexera X-2 UHPLC equipped with a Phenomenex XB-C18 column (1.7  $\mu$ m, 100  $\times$  2.1 mm) and a 2.0  $\mu$ L injection volume. Gradient elution was performed using solvents *A* = 0.1% formic acid and *B* = acetonitrile. The gradient program was as follows: 0–2 min, 5% *B*; ramp to 18% *B* at 17.5 min; ramp to 50% *B* at 32 min; hold for 1 min. Detection was performed with a Shimadzu SPD M30A photodiode array and a Bruker Impact II Ultra-high Resolution QqTOF mass spectrometer operating in negative-ion mode electrospray ionization. UV spectra were recorded in the 250–400 nm range and the MS was operated with capillary voltage, 3.5 kV; nebulizing gas pressure, 4.0 bar; drying gas temperature, 210  $^{\circ}$ C; drying gas flow, 8 L min $^{-1}$ , *m/z* range, 50–1000. FA (Aldrich, 99%), *p*CA (Sigma, 98%), FA-MeAra **4**, and *p*CA-MeAra **3** were quantified in the MS using authentic standards with *m*-coumaric acid (Acros, 99%) as the internal standard; the arabinose-containing standards were the same as those used in the original method [33], and their synthesis has been described [36, 37]. Standards were not available for DFA-MeAra and DFA-MeAra<sub>2</sub>, so changes in these were estimated from a simple comparison of MS peak areas.

### Abbreviations

FA: Ferulic acid **1**; *p*CA: *p*-Coumaric acid; FA-Ara: 5-*O*-Feruloyl arabinofuranoside **2**; FA-MeAra: Methyl 5-*O*-feruloyl arabinofuranoside **4**; *p*CA-MeAra: Methyl 5-*O*-*p*-coumaroyl arabinofuranoside **3**; DFA: Dehydrodiferulate or (dehydro) diferulic acid; DFA-MeAra: Methyl dehydrodiferuloyl monoarabinofuranoside **6**; DFA-MeAra<sub>2</sub>: Dimethyl dehydrodiferuloyl diarabinofuranoside **6**; EIC: Extracted-ion chromatogram; IDF: Insoluble dietary fiber ( $\approx$  CWR, cell wall residue); LC–MS: Liquid chromatography–mass spectrometry (or spectrometer).

### Acknowledgements

The authors would like to thank Dr. Steven Karlen and Prof. Richard F. Helm for providing the quantification standards, and Dr. Mirko Bunzel for providing the maize and rye IDF.

### Authors' contributions

JR originated the research idea based on CL's original method; AE designed and performed the experiments. AE, JR, CL wrote the manuscript and produced the figures. All authors read and approved the final manuscript.

### Funding

This research was funded by the DOE Great Lakes Bioenergy Research Center (DOE Office of Science BER DE-FC02-07ER64494).

### Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### Ethics approval and consent to participate

Not applicable.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup> Great Lakes Bioenergy Research Center, Wisconsin Energy Institute, University of Wisconsin, Madison, WI, USA. <sup>2</sup> Institute Jean-Pierre Bourgin, INRAE, AgroParisTech, Université Paris-Saclay, Versailles, France. <sup>3</sup> Department of Biochemistry, University of Wisconsin, Madison, WI, USA.

Received: 30 September 2020 Accepted: 24 November 2020

Published online: 10 December 2020

### References

- Nakamura Y, Higuchi T. Ester linkage of *p*-coumaric acid in bamboo lignin. III. Dehydrogenative polymerization of coniferyl *p*-hydroxybenzoate and coniferyl *p*-coumarate. *Cell Chem Technol*. 1978;12:209–21.
- Ralph J. Hydroxycinnamates in lignification. *Phytochem Rev*. 2010;9:65–83.
- Mottiar Y, Vanholme R, Boerjan W, Ralph J, Mansfield SD. Designer lignins: harnessing the plasticity of lignification. *Curr Opin Biotechnol*. 2016;37:190–200.
- Withers S, Lu F, Kim H, Zhu Y, Ralph J, Wilkerson CG. Identification of a grass-specific enzyme that acylates monolignols with *p*-coumarate. *J Biol Chem*. 2012;287:8347–55.
- Petrik DL, Karlen SD, Cass CL, Padmakshan D, Lu F, Liu S, Le Bris P, Antelme S, Santoro N, Wilkerson CG, et al. *p*-Coumaroyl-CoA: monolignol transferase (PMT) acts specifically in the lignin biosynthetic pathway in *Brachypodium distachyon*. *Plant J*. 2014;77:713–26.
- Marita JM, Hatfield RD, Rancour DM, Frost KE. Identification and suppression of the *p*-coumaroyl CoA: hydroxycinnamyl alcohol transferase in *Zea mays* L. *Plant J*. 2014;78:850–64.
- Wilkerson CG, Mansfield SD, Lu F, Withers S, Park J-Y, Karlen SD, Gonzales-Vigil E, Padmakshan D, Unda F, Rencoret J, Ralph J. Monolignol ferulate transferase introduces chemically labile linkages into the lignin backbone. *Science*. 2014;344:90–3.
- Ralph J, Lapierre C, Boerjan W. Lignin structure and its engineering. *Curr Opin Biotechnol*. 2019;56:240–9.
- Grabber JH, Hatfield RD, Lu F, Ralph J. Coniferyl ferulate incorporation into lignin enhances the alkaline delignification and enzymatic degradation of maize cell walls. *Biomacromol*. 2008;9:2510–6.
- Zhou S, Runge T, Karlen SD, Ralph J, Gonzales-Vigil E, Mansfield SD. Chemical pulping advantages of Zip-lignin hybrid poplar. *ChemSusChem*. 2017;10:3565–73.
- Kim KH, Dutta T, Ralph J, Mansfield SD, Simmons BA, Singh S. Impact of lignin polymer backbone esters on ionic liquid pretreatment of poplar. *Biotechnol Biofuels*. 2017;10(101):1–10.
- Bhalla A, Bansal N, Pattathil S, Li M, Shen W, Particka CA, Semaan R, Gonzales-Vigil E, Karlen SD, Ralph J, et al. Engineered lignin in poplar biomass facilitates Cu-AHP pretreatment. *ACS Sustain Chem Eng*. 2018;6:2932–41.

13. Regner M, Bartuce A, Padmakshan D, Ralph J, Karlen SD. Reductive cleavage method for quantitation of monolignols and low-abundance monolignol conjugates. *ChemSusChem*. 2018;11:1600–5.
14. Karlen SD, Free HCA, Padmakshan D, Smith BG, Ralph J, Harris PJ. Commelinid monocotyledon lignins are acylated by *p*-coumarate. *Plant Physiol*. 2018;177:513–21.
15. Karlen SD, Zhang C, Peck ML, Smith RA, Padmakshan D, Helmich KE, Free HCA, Lee S, Smith BG, Lu F, et al. Monolignol ferulate conjugates are naturally incorporated into plant lignins. *Sci Adv*. 2016;2:e1600393.
16. Smith RA, Gonzales-Vigil E, Karlen SD, Park J-Y, Lu F, Wilkerson CG, Samuels L, Mansfield SD, Ralph J. Engineering monolignol *p*-coumarate conjugates into Poplar and Arabidopsis lignins. *Plant Physiol*. 2015;169:2992–3001.
17. Lu F, Karlen SD, Regner M, Kim H, Ralph SA, Sun R-C, Kuroda K-I, Augustin MA, Mawson R, Sabarez H, et al. Naturally *p*-hydroxybenzoylated lignins in palms. *BioEnergy Res*. 2015;8:934–52.
18. Lu F, Ralph J. Detection and determination of *p*-coumaroylated units in lignins. *J Agric Food Chem*. 1999;47:1988–92.
19. Karlen SD, Smith RA, Kim H, Padmakshan D, Bartuce A, Mobley JK, Free HCA, Smith BG, Harris PJ, Ralph J. Highly decorated lignins occur in leaf base cell walls of the Canary Island date palm *Phoenix canariensis*. *Plant Physiol*. 2017;175:1058–67.
20. Lu F, Ralph J. The DFRC (derivatization followed by reductive cleavage) method and its applications for lignin characterization. In: Lu F, editor. *Lignin: structural analysis, applications in biomaterials, and ecological significance*. Hauppauge: Nova Science Publishers, Inc; 2014. p. 27–65.
21. Jacquet G, Pollet B, Lapiere C, Mhamdi F, Rolando C. New ether-linked ferulic acid-coniferyl alcohol dimers identified in grass straws. *J Agric Food Chem*. 1995;43:2746–51.
22. Ralph J, Grabber JH, Hatfield RD. Lignin-ferulate crosslinks in grasses: active incorporation of ferulate polysaccharide esters into ryegrass lignins. *Carbohydr Res*. 1995;275:167–78.
23. Buanaafina MMD. Feruloylation in grasses: current and future perspectives. *Mol Plant*. 2009;2:861–72.
24. Bento-Silva A, Patto MCV, Bronze MD. Relevance, structure and analysis of ferulic acid in maize cell walls. *Food Chem*. 2018;246:360–78.
25. Harris PJ, Hartley RD. Phenolic constituents of the cell walls of monocotyledons. *Biochem Syst Ecol*. 1980;8:153–60.
26. Hartley RD, Jones EC, Wood TM. Lignin-carbohydrate linkages in plant cell walls. Part 3. Carbohydrates and carbohydrate esters of ferulic acid released from cell walls of *Lolium multiflorum* by treatment with cellulolytic enzymes. *Phytochemistry*. 1976;15:305–7.
27. Mueller-Harvey I, Hartley RD, Harris PJ, Curzon EH. Linkage of *p*-coumaroyl and feruloyl groups to cell wall polysaccharides of barley straw. *Carbohydr Res*. 1986;148:71–85.
28. Hartley RD, Morrison WH III, Himmelsbach DS, Borneman WS. Cross-linking of cell wall phenolic arabinoxylans in graminaceous plants. *Phytochemistry*. 1990;29:3705–9.
29. Ralph J, Hatfield RD, Grabber JH, Jung HG, Quideau S, Helm RF. Cell wall cross-linking in grasses by ferulates and diferulates. In: Lewis NG, Sarkanen S, editors. *Lignin and lignan biosynthesis*, vol. 697. ACS symposium series. Washington, DC: American Chemical Society; 1998. p. 209–36.
30. Li G, Jones KC, Eudes A, Pidatala VR, Sun J, Xu F, Zhang C, Wei T, Jain R, Birdseye D, et al. Overexpression of a rice BAHD acyltransferase gene in switchgrass (*Panicum virgatum* L.) enhances saccharification. *BMC Biotechnol*. 2018;18:54.
31. Bartley LE, Peck ML, Kim SR, Ebert B, Manisseri C, Chiniquy DM, Sykes R, Gao L, Rautengarten C, Vega-Sanchez ME, et al. Overexpression of a BAHD acyltransferase, OsAt10, alters rice cell wall hydroxycinnamic acid content and saccharification. *Plant Physiol*. 2013;161:1615–33.
32. de Souza WR, Martins PK, Freeman J, Pellny TK, Michaelson LV, Sampaio BL, Vinecky F, Ribeiro AP, da Cunha BADB, Kobayashi AK, et al. Suppression of a single BAHD gene in *Setaria viridis* causes large, stable decreases in cell wall feruloylation and increases biomass digestibility. *New Phytol*. 2018;218:81–93.
33. Lapiere C, Voxeur A, Boutet S, Ralph J. Arabinose conjugates diagnostic of ferulate-ferulate and ferulate-monomolignol cross-coupling are released by mild acidolysis of grass cell walls. *J Agric Food Chem*. 2019;67:12962–71.
34. Lapiere C, Voxeur A, Karlen SD, Helm RF, Ralph J. Evaluation of feruloylated and *p*-coumaroylated arabinosyl units in grass arabinoxylans by acidolysis in dioxane/methanol. *J Agric Food Chem*. 2018;66:5418–24.
35. Fieser LF, Fieser M. *Reagents for organic synthesis*, vol. 1. New York: Wiley; 1967. p. 192.
36. Hatfield RD, Helm RF, Ralph J. Synthesis of methyl 5-*O*-*trans*-feruloyl- $\alpha$ -L-arabinofuranoside and its use as a substrate to assess feruloyl esterase activity. *Anal Biochem*. 1991;194:25–33.
37. Helm RF, Ralph J, Hatfield RD. Synthesis of feruloylated and *p*-coumaroylated methyl glycosides. *Carbohydr Res*. 1992;229:183–94.
38. Allerdings E, Ralph J, Schatz P, Gniechwitz D, Steinhart H, Bunzel M. Isolation and structural identification of di-arabinosyl 8-*O*-4-dehydridiferulate from maize bran insoluble fibre. *Phytochemistry*. 2005;66:113–24.
39. Busse-Wicher M, Li A, Silveira RL, Pereira CS, Tryfona T, Gomes TC, Skaf MS, Dupree P. Evolution of xylan substitution patterns in gymnosperms and angiosperms: implications for xylan interaction with cellulose. *Plant Physiol*. 2016;171:2418–31.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

