

Digestive and metabolic bioavailability in healthy humans of 15N-labeled rapeseed and flaxseed protein incorporated in biscuits

Romain Tessier, Juliane Calvez, Gheorghe Airinei, Nadezda Khodorova, Sylvie Dauguet, Olivier Galet, Martin Chapelais, Daniel Tomé, Robert Benamouzig, Claire Gaudichon

▶ To cite this version:

Romain Tessier, Juliane Calvez, Gheorghe Airinei, Nadezda Khodorova, Sylvie Dauguet, et al.. Digestive and metabolic bioavailability in healthy humans of 15N-labeled rapeseed and flaxseed protein incorporated in biscuits. The American Journal of Clinical Nutrition, 2023, 117, pp.896 - 902. 10.1016/j.ajcnut.2023.02.020 . hal-04204013

HAL Id: hal-04204013 https://agroparistech.hal.science/hal-04204013

Submitted on 11 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The American Journal of CLINICAL NUTRITION



journal homepage: www.journals.elsevier.com/the-american-journal-of-clinical-nutrition

Original Research Article

Digestive and metabolic bioavailability in healthy humans of ¹⁵N-labeled rapeseed and flaxseed protein incorporated in biscuits

Romain Tessier¹, Juliane Calvez¹, Gheorghe Airinei¹, Nadezda Khodorova¹, Sylvie Dauguet², Olivier Galet³, Martin Chapelais¹, Daniel Tomé¹, Robert Benamouzig¹, Claire Gaudichon^{1,*}

¹ Université Paris-Saclay, AgroParisTech, INRAE, UMR PNCA, Palaiseau, France; ² Terres Inovia, Pessac, France; ³ Avril, Paris, France

ABSTRACT

Background: In the search to diversify protein sources for humans, oilseeds are good candidates due to the high protein content of their coproducts after oil extraction. Among them, rapeseed presents a well-balanced amino acid (AA) profile. Flaxseed is an emerging source but the nutritional value of its protein is not yet documented.

Objectives: This study aimed to determine the nitrogen (N) and AA bioavailability of these protein sources.

Methods: Nineteen healthy volunteers were intubated with a naso-ileal tube. They ingested 156 g biscuits containing intrinsically labeled ¹⁵N rapeseed (n = 10) or flaxseed (n = 9) protein over a 4-h period. Ileal digesta, blood, and urine were sampled over 8 h after the first meal ingestion. N and ¹⁵N enrichment and AAs were measured to determine digestive and deamination losses. Ileal digestibility, the digestible indispensable AA score (DIAAS) and net postprandial protein utilization (NPPU) were calculated.

Results: Real ileal digestibility was $80.7 \pm 6.5\%$ for rapeseed protein and $92.2 \pm 2.0\%$ for flaxseed protein (P = 0.0002). Mean indispensable AA (IAA) digestibility reached $84.1 \pm 6.9\%$ and $93.3 \pm 6.7\%$ for rapeseed and flaxseed, respectively, lysine being the lowest digestible IAA for both sources. Despite moderate digestibility, the DIAAS was 1.1 for rapeseed but only 0.6 for flaxseed due to lysine insufficiency. Deamination losses accounted for $20.0 \pm 6.5\%$ of dietary N for flaxseed and $11.0 \pm 2.8\%$ for rapeseed (P = 0.002). The NPPU did not differ between the protein sources, with $71.3 \pm 6.5\%$ for flaxseed and $69.7 \pm 7.6\%$ for rapeseed.

Conclusions: Despite good digestibility, flaxseed protein cooked in biscuits was penalized by both lysine insufficiency and poor lysine digestibility that decreased its DIAAS and increased deamination. By contrast, rapeseed was moderately digestible but presented no limiting IAA, resulting in an excellent DIAAS and low deamination.

This study was registered at clinicaltrials.gov as NCT04024605.

Keywords: stable isotopes, ileal digestibility, intestinal tubes, oilseed protein, digestible indispensable amino acid score, humans

Introduction

In view of the diversification of plant-based protein foods, the protein quality of these sources needs to be clarified. Oilseeds are good candidates as their oil coproduct, oilcake, is rich in protein. Sunflower and rapeseed represent a significant reservoir of protein as they are among the 10 major crops in France and Europe [1]; in 2021, 1911 Mt of sunflower and 3286 Mt of rapeseed were produced in France [2]. Although a minor crop, flaxseed has been the subject of increased

interest regarding the nutritional quality of its oil, but its protein quality has not been assessed.

We recently determined the ileal digestibility of amino acid (AA) from a sunflower isolate incorporated in a biscuit and found a mean digestibility of indispensable amino acid (IAA) that reached 89% [3]. In a previous study, we had also determined that the ileal nitrogen (N) digestibility of rapeseed protein isolate was only 84% [4], whereas it was around 90%–91% for soy [5] and pea isolates [6] as well as gluten [7]. During these studies, the digestibility of individual AAs was not

https://doi.org/10.1016/j.ajcnut.2023.02.020

Received 21 October 2022; Received in revised form 10 February 2023; Accepted 21 February 2023

Available online 24 February 2023

Abbreviations used: AA, amino acid; Asx, aspartate and asparagine; AP, atom percent; CS, chemical score; DAA, dispensable amino acid; DIAAS, digestible indispensable AA score; DM, dry matter; Glx, glutamate and glutamine; IAA, indispensable amino acid; IRMS, isotope ratio mass spectrometer; N, nitrogen; NPPU, net postprandial utilization; PEG, polyethylene glycol.

^{*} Corresponding author.

E-mail address: claire.gaudichon@agroparistech.fr (C. Gaudichon).

^{0002-9165/© 2023} The Author(s). Published by Elsevier Inc. on behalf of American Society for Nutrition. This is an open access article under the CC BY-NC-ND license (http:// creativecommons.org/licenses/by-nc-nd/4.0/).

determined. Since the FAO recommended in 2013 that the digestible indispensable AA score (DIAAS) should be the gold standard to assess the capacity of protein to satisfy AA requirements [8], the measurement of AA digestibility has become necessary. However, its determination faces certain methodological barriers. The ileal digestibility of several protein isolates from milk, soy, corn, and sunflower has been measured in humans using a naso-ileal tube to collect digesta [3,9–11].

The aim of our study was to generate data on the ileal digestibility of AAs, the DIAAS, and the metabolic bioavailability of dietary N in 2 oilseed sources: rapeseed, which is known to have moderate protein digestibility but a good biological value [4], and flaxseed, where the quality of its protein has not yet been documented. For this purpose, the proteins were labeled intrinsically with ¹⁵N and extracted from the seeds. They were then incorporated in biscuits that were given as a ¹⁵N test meal to healthy volunteers equipped with a naso-ileal tube.

Methods

Subjects

This study formed part of the PRODIGE (Optimized PROcess for sustainable production of highly DIGEstible oilseed proteins for humans) project and was conducted at the Human Nutrition Research Center in Avicenne Hospital (AP-HP, Bobigny). Recruitment started in July 2020 and was completed in May 2021. The study was approved by the Sud Mediterranée IV Ethics Committee and the French Agency for Medicinal and Health Products Safety. It was registered in the www .clinicaltrials.gov database (NCT04024605).

The number of volunteers was determined in accordance with previous studies on the AA digestibility of protein sources [9,10,12]. These studies were performed on 6 to 10 volunteers per group. Based on the data on rapeseed protein digestibility [4] and that of other plant protein isolates studied in humans [6,7,9], a difference of $6 \pm 4\%$ of N digestibility could be expected. In view of the between subject design (2-tailed unpaired Student's *t* test), risks at $\alpha = 5\%$, and $(1 - \beta) = 80\%$, the sample size thus determined was 9 per group (G*Power 3.1).

The eligibility criteria were: BMI between 18 and 30 kg/m², age between 18 and 65 y, negative serology for human immunodeficiency virus, hepatitis C virus antibody, hepatitis B virus surface antigen, and an absence of dietary allergies and digestive disease. All volunteers signed an informed consent form. Thirty-four volunteers were recruited, 13 of whom did not complete the protocol. The causes of these failures included an impossibility to introduce the tube into the nose (n = 1), poor tolerance of the tube (n = 5), nonmigration of the tube through the pylorus (n = 4), and insufficient migration in the small intestine (n = 3). Twenty-one volunteers were thus allocated to the rapeseed and flaxseed groups. Randomization was not possible because the flaxseed and rapeseed biscuits could not be produced at the same time. Two batches of each biscuit were produced within a short interval of around a month, thus permitting overlapping of the enrollment of volunteers in each group. One volunteer who completed the protocol had to be excluded because the tube was not positioned in the terminal ileum and another because the tube had migrated in the colon (Supplemental Figure 1).

The final sample size was therefore n = 10 for rapeseed and n = 9 for flaxseed. The characteristics of the subjects were similar in the 2 groups (Table 1).

Test meals

Rapeseed and flaxseed were labeled by Terres Inovia using ¹⁵N ammonium nitrate enriched at 10% during growth, as detailed in

TABLE 1

munopointerie entracteristics of the subjects	ŀ	Anthro	pometric	charac	teristics	of	the	subje	cts
---	---	--------	----------	--------	-----------	----	-----	-------	-----

Characteristic	Rapeseed	Flaxseed
Sex (F/M)	4/6	3/6
Weight (kg)	70 ± 12	68 ± 10
Height (cm)	172 ± 12	169 ± 5
BMI (kg/m ²)	23.5 ± 3	23.7 ± 3
Age (y)	34 ± 11	36 ± 12

¹ Values are means \pm SDs, n = 19.

Supplemental Table 1. The seeds were hulled and de-oiled (Olead). 15 N protein was extracted and purified according to the processes described in Supplemental Figure 2. A rapeseed isolate could not be produced because of the small quantity of harvested grains. The protein content (N × 6.25) in the 15 N labeled protein fractions was 50% for the rapeseed concentrate and 80% for the flaxseed isolate. The enrichment of rapeseed and flaxseed protein was 0.73 atom percent (AP) and 0.51 AP, respectively. The protein fractions were incorporated in the biscuits that were cooked at 180°C for 21 min (Foodinnov), with a target protein concentration of 15%–16% dry matter (DM). The wheat flour contained a small quantity of gluten (2.8% of the biscuit) (Supplemental Table 2).

The volunteers ingested 156 g of biscuits, divided into 9 portions. The first portion contained 4 biscuits and the other 8 contained 1 biscuit. The protein intake was 18 g and 20 g for rapeseed and flaxseed biscuits, respectively. Alongside the biscuits, the subjects ingested a total of 100 g apple puree, in accordance with our study on sunflower [3]. Overall, energy supply was 670 kcal (rapeseed) to 720 kcal (flaxseed).

Rapeseed protein contained 39% IAA and flaxseed 34%. Rapeseed contained twice as much lysine than flaxseed (Supplemental Table 3).

Clinical design

The day before the experiment, the subjects were equipped with an intestinal tube as previously described [3,10,13]. The protocol design is presented in Figure 1. After fasting overnight, a solution of polyethylene glycol-4000 (PEG, 20 g/L), a nonabsorbable marker of liquid flow rate, was infused via the intestinal tube at 1 mL/min. Ileal effluent, blood, and urine were collected at baseline. The first meal portion was given to the volunteers, followed by a portion every half hour for 4 h. Volunteers were not informed on the type of protein they had to ingest (single-blind). The digestive contents were collected continuously and pooled by half hour for 8 h after the first meal. Blood was sampled every 30 min for 4 h and then every hour. All urine was collected and pooled by 2-h periods.

Analytical methods

The concentration of PEG in the digesta was measured using the turbidimetric method [14] to determine the ileal flow rate. Urea concentrations in plasma and urine were measured by means of the Urease-Berthelot method (Urea assay, Randox).

Rapeseed and flaxseed proteins, together with the digestive contents, were hydrolyzed for 24 h with HCl 6N at 110°C with norvaline as an internal standard. For methionine and cysteine, a performic acid oxidation was done before acid hydrolysis. For tryptophan, proteins were hydrolyzed for 20 h at 110°C in the presence of barium hydroxide 2N, with 5-methyl-tryptophan as internal standards.

For AA quantification, calibration standards were composed of a mixture of AA (Waters) to which norvaline was added. AA and

standards were derivatized using the AccQ-Tag Ultra Derivatization kit (Waters) according to the manufacturer's protocol. The AA analysis was performed on an Acquity H-class Ultra-HPLC system with a photo diode array detector (Waters), as described previously [11].

Before measuring the ¹⁵N enrichments, AA from digesta and meals were isolated using a hydrogen form of ion exchange resin and derivatized with ethyl chloroformate, as previously described [15]. ¹⁵N enrichments were analyzed by gas chromatography (GC 6890N, Agilent Technologies) in constant flow mode, coupled to an isotope ratio mass spectrometer (IRMS) (Isoprime GV Instrument) via the GC5 Isoprime interface [11].

Urinary urea was isolated on a sodium form of cation exchange resin after the isolation of urinary ammonia and treatment with urease. Plasma was split into 3 fractions: proteins, free AAs, and urea, as described previously [7]. Resin-containing urea was eluted with KHSO₄ (2.5M).

¹⁵N recovery in the digestive contents, urea pools, plasma free AA, biscuits, and protein was measured using an elemental analyzer (Vario Micro Cube, Elementar) coupled with an IRMS (Isoprime, GV Instrument). Atropine (Thermo Electron) and L-glutamic acid (USGS41, Sigma Aldrich) were used as elemental and isotopic standards, respectively.

N, AA digestibility, and DIAAS were the primary outcomes, and the net postprandial protein utilization (NPPU) was the secondary outcome.

Calculations

The ileal flow rate was evaluated for each period of 30 min (F, mL/ 30 min):

 $F = \frac{[PEG]_{solution}}{[PEG]_{digesta}} \times perfusion flow rate \times 30$

where [PEG] is the concentration of PEG in the infused solution (1 mL/min) and in the digestive contents.

The total N flow rate ($N_{tot \ ileum}$, mmol/30 min) in the ileum was assessed for each period:

$$N_{\text{tot ileum}} = \frac{N_{\text{s}} \times DM_{\text{s}} \times F}{14 \times 10}$$

where N_s is the percentage of N, DM_s is the DM of the ileal sample (g/100 g), F is the ileal flow rate (mL/30 min), and 14 is the molar mass of N (g/mol).

The dietary N flow rate ($N_{diet \ ileum}$, mmol/30 min) was calculated as follows:

$$N_{diet \ ileum} = N_{tot \ ileum} imes rac{APE_{ileum}}{APE_{meal}}$$

with APE the enrichment excess compared to ${}^{15}N$ basal enrichment that was measured at t = 0 in each volunteer.

Endogenous N (N_{endo ileum} in mmol/30 min) was the difference between total and dietary N recovered in the ileum.

The real ileal digestibility (RID, % of ingested N) was:

$$RID = \frac{N_{ingested} - \sum N_{diet \ ileum}}{N_{ingested}} \times 100$$

With $N_{ingested}$ the amount of N ingested by the subjects (mmol) and $\Sigma N_{diet \ ileum}$ the cumulative dietary N recovered in the ileum over 8 h (mmol).

The quantity of dietary AA in digesta (AA_{diet i}, mmol) at each period was:

$$AA_{diet \; i}(t) = \left[AA\right]_{ileum \; i}(t) \times DM \; (t) \times F(t) \times \frac{APE_{ileum \; i}(t)}{APE_{protein}}$$

where [AA]_{ileum} i(t) is the quantity of each AA "i" in the digestive contents at each period "t" (mmol/g), DM the dry matter (g/100 mL), F the ileal flow rate (mL/30 min), APE the ¹⁵N excess enrichment of each AA "i" at each period "t," and APE_{protein} that of rapeseed or flaxseed protein. Basal abundance was measured at t = 0 in each volunteer.

The chemical score (CS) was defined for each IAA:

 $CS = \frac{mass \text{ of dietary IAA}_i \text{ in 1g of dietary protein}}{mass \text{ of the same dietary IAA}_i \text{ in 1g of reference protein}}$

The reference protein profile was the recommended AA scoring pattern for older children, adolescents, and adults [8].

The digestible IAA ratio (DIAA ratio), was calculated for IAA:

DIAA ratio = $CS_i \times AA$ ileal digestibility_i

The DIAAS was the smallest DIAA ratio calculated. The protein mass was calculated using N \times 6.25 factor, or a more specific factor recommended for plants, which is 5.6 [16].

The dietary N incorporation in plasma proteins, body, and urinary urea pools was calculated using the same dilution principle as for digesta. Plasma protein and body urea pools were calculated as described previously [7].

The NPPU (% of ingested N) was the amount of N retained in the body after 8 h. It corresponded to the N ingested minus metabolic and digestive losses:

$$NPPU = \frac{N_{ingested} - \left(\sum N_{diet \ ileum} + \sum N_{diet \ urinary \ urea} + N_{diet \ body \ urea}\right)}{N_{ingested}} \times 100$$

with Ndiet body urea the remaining dietary N in body urea at 8 h.

Statistical analysis

Data are expressed as means \pm SD. Differences between the groups were estimated using an unpaired Student's *t* test after verification that normal distribution of data was not rejected. Kinetic data were analyzed in a mixed model with the group as a fixed factor and time as a repeated factor (SAS 9.4 Software, SAS Institutes). Differences between groups at each time point were determined by contrast analyses when the time \times group interaction was significant. Differences were considered to be statistically significant for a *P* value lower than 0.05.

Results

Digestive flow rates and ileal digestibility

The liquid flow rate (Figure 2A) was similar in the 2 groups during the first 2 h. It increased in the rapeseed group at 3 h but remained stable in the flaxseed group. This flow rate averaged 139 ± 61 mL/h with flaxseed ingestion and 166 ± 51 mL/h in the rapeseed group.

Dietary N (Figure 2B) appeared in the ileum 1 h after ingestion of the first meal, the peak being observed at 5 h in the rapeseed group and with a plateau after 2 h for flaxseed. The amount of dietary N was significantly higher in the rapeseed group than in the flaxseed group from 2.5 h to 7.5 h after the meal. The real ileal protein digestibility of flaxseed protein was therefore $92.2 \pm 2.0\%$, but it was only $80.7 \pm 6.5\%$ for rapeseed (P = 0.0002).

The endogenous N ileal flow (Figure 2C) was ~6.5 mmol/h in both groups and did not display any marked variations during the post-prandial window nor did it differ between the groups.

Ileal AA digestibility and the DIAAS

Ileal AA digestibility values of rapeseed and flaxseed protein were determined (Table 2). In the flaxseed group, values ranged from 78.2 \pm 6.2% for lysine to 96.5 \pm 3.3% for methionine. In the rapeseed group, lysine was also the lowest (75.3 \pm 10.1%) and methionine the highest digestible AA (92.6 \pm 3.6%). For all AAs except leucine and lysine, digestibility was higher with flaxseed than with rapeseed.

Chemical and DIAA ratios were calculated using either the default N-to-protein conversion factor of 6.25 or a specific factor for plant protein of 5.6 [16] (Table 3). Flaxseed was limited in lysine, with a CS ranging from 0.7 to 0.8 depending on the conversion factor. For rapeseed, the CS was higher than 1 whatever the conversion factor. The DIAAS of rapeseed ranged from 1.1 at its lowest (N \times 6.25) to 1.2 (N \times 5.6), whereas it was only 0.6 to 0.7 for flaxseed, due to lysine insufficiency.

Deamination losses of dietary N and NPPU

Digestive and deamination losses of dietary N are shown in Figure 3. The transfer of dietary N to urinary urea during the 8 h after the rapeseed meal was $3.4 \pm 1.0\%$ of ingested N, whereas it reached $6.4 \pm 2.9\%$ in the flaxseed group (P = 0.01). Similarly, the amount of dietary N present in the body urea pool at 8 h was almost doubled in the flaxseed group when compared to rapeseed (P = 0.02). Consequently, total deamination losses were $11.0 \pm 2.8\%$ in the rapeseed group but reached $20.0 \pm 6.5\%$ in the flaxseed group (P = 0.002). However, this difference was balanced by higher digestive losses reported in the rapeseed group; as a result, NPPU did not differ between the groups ($69.8 \pm 7.6\%$ for rapeseed compared with $71.5 \pm 6.5\%$ for flaxseed).

Discussion

This study aimed to evaluate the nutritional quality of protein fractions from rapeseed and flaxseed. Plant proteins were labeled intrinsically with ¹⁵N before being purified and incorporated into biscuits. We determined precisely the ileal digestibility of protein and AAs and subsequently the DIAAS, as well as N retention in healthy humans.

Ileal losses

To our knowledge, our work is the first to have measured the protein digestibility of flaxseed protein isolate, which was good (higher than 92%). Few data are available in the literature regarding the protein quality of flaxseed. When given as a linseed meal to pigs, the standardized ileal digestibility was 76%, whereas it was 72% for a canola meal [17]. Similar values were obtained in rats when flaxseed was given as cereal bars or in an extruded form [18]. The higher

digestibility we obtained could be ascribed to the fact that flaxseed was given as an isolate. This value of 92% was therefore similar to other protein isolates we have assessed using the same methodology, namely soy, pea, and gluten [5–7,11]. By contrast, we had previously found lower protein digestibility values for sunflower (87%) and rapeseed (84%) protein isolates [3,4]. The moderate digestibility of rapeseed is known to be due to its albumin fractions, and particularly napin, which is rich in cysteine and presents some disulfide bonds [19]. The value we determined during the present study was 3% lower than that previously obtained [4]. This result was not surprising because the rapeseed protein fraction we tested was a concentrate rather than an isolate. Further, it was cooked in a biscuit whereas during the study by Bos et al. [4], it had been incorporated in a liquid meal. The degree of purification of plant protein has been identified as a favorable factor in terms of its digestibility because of the removal of antinutritional factors such as the phenolic compounds and phytates that are present in rapeseed [20].

The digestibility of individual AAs was found to be similar to that determined for global protein in both oilseeds, with good homogeneity of AA digestibility values, except for lysine. Indeed, the digestibility of lysine was markedly lower than that of other AAs, as well as leucine in flaxseed only. A lower digestibility of lysine has already been reported in pigs with values 8% to 11% lower than that of mean AA digestibility for flaxseed and canola meal, respectively [17]. However, such a low lysine digestibility of oilseed meals in pigs has not been observed by others [21]. In our study, low lysine digestibility could partly be explained by the fact that the protein was cooked in biscuits, together with sugars, thus altering the lysine through Maillard reactions. Moreover, cooking with gluten-containing products can create aggregates. In particular, methylglyxocal, a product of the Maillard reaction, may react with glutenin, leading to a 30% reduction in in vitro digestibility [22]. The loss of lysine digestibility was especially high for flaxseed, the digestibility of which was 20% lower than that of other AAs.

DIAAS

The determination of AA digestibility enabled us to calculate the DIAAS. Using the default factor for N-to-protein conversion, 6.25, the DIAAS of flaxseed was only 0.6 due to lysine limitation. Although this factor has been acknowledged to be overestimated whatever the protein source, especially from plants [23,24], it is employed universally, and particularly for regulatory and commercial purposes. When using a more specific factor for plant proteins (5.6) [16], the DIAAS of flaxseed rose to 0.7. Rapeseed had no limiting IAA and a DIAAS that revealed a 10% to 23% excess of lysine, depending on the conversion factor.

The AA profile of rapeseed was sufficiently balanced, with an excess of IAAs of \geq 42% regarding the CS, so was not penalized by its



FIGURE 1. Experimental design of the study.



FIGURE 2. (A) Liquid ileal flow rate. (B) Dietary N flow in the ileum. (C) Endogenous N flow in the ileum. Values are means \pm SDs. n = 9 in flaxseed group (black circle) and n = 10 in rapeseed group (gray square). The data were analyzed using a mixed model with repeated measurements (unstructured matrix for the liquid low rate and Toeplitz matrix for dietary and endogenous N). * indicates a significant difference between groups at a particular time point (P < 0.05).

moderate digestibility. By contrast, the low lysine digestibility enhanced lysine deficiency in flaxseed. It would have been interesting to assess the digestibility of the raw flaxseed isolate in order to determine whether the heating process was responsible for this particular effect on lysine and thus lowered the score. As total digestible lysine can be higher than digestible reactive lysine [25], the analysis of reactive lysine would have helped to refine the results, but this would require developing a specific method for gas chromatography combustion IRMS analysis.

Deamination losses

We found a marked effect for the protein, insofar as ¹⁵N transfer was almost doubled in the urine and residual body urea pool at 8 h in the flaxseed group compared to rapeseed. This resulted in a total deamination rate of around 15% for rapeseed compared with 26% for flaxseed. The low deamination rate observed for rapeseed was similar to what we had observed previously [4], where we found a postprandial transfer of 6% in urine and a similar amount in residual body urea. This low deamination rate could be ascribed to the adequate AA profile of rapeseed, with no limiting AA. By contrast, the high deamination rate of flaxseed could be explained by the low lysine content. Consistent with this, we had reported a deamination rate of 27% for wheat [7] in which lysine was limiting to a concentration of 50% below the requirement. In this latter study, gluten had been incorporated in French biscuits that had been toasted, as during the present study.

Finally, the postprandial retention of both proteins was similar, the high deamination losses of flaxseed being compensated for by low digestive losses, whereas the picture was the reverse for rapeseed.

Strengths and limitations of the study

Our study was the first to assess the postprandial quality of flaxseed protein and determine the DIAAS of rapeseed and flaxseed in human volunteers. The labeling of plants with ¹⁵N offered a particularly interesting tool because in addition to digestibility and the DIAAS score, it also provides data on N retention. It was thus possible to obtain a complete overview of protein AA and N bioavailability. However, plant labeling in the field involves a degree of risk and in our case, the rapeseed crop yield was not sufficiently good to enable a complete purification of protein, unlike flaxseed. However, this work did reveal that even with a lower degree of purification, rapeseed concentrate was sufficient to meet human AA requirements and even to compensate for lysine deficiency from other sources. In addition, ¹⁵N recycling in the intestine can result in a slight underestimation of digestibility of around 1.5% [26]. In contrast, the repeated meal protocol of this study could have led to an overestimation of digestibility and NPPU. Nevertheless, this is neither supported by our former study, where rapeseed was given as a bolus [4], nor by results obtained for casein and pea given as bolus [6,27] or repeated meals [11]. Lastly, the lack of true randomization could also be considered as a bias [28], but this was limited by the fact that both groups displayed homogenous basal variables and were studied at almost the same time and during successive periods.

In conclusion, we found that flaxseed and rapeseed displayed similar postprandial protein retention, with high levels of digestive losses for rapeseed and of the deamination rate for flaxseed, the latter being associated with the quantity of digestible lysine. Interestingly, despite its moderate AA digestibility, and particularly for lysine, no

TABLE 2

Ileal digestibility of AA from flaxseed and rapeseed protein cooked in biscuits¹

Amino Acid	Flaxseed	Rapeseed	P^2
IAA digestibility (%)			
Histidine	93.8 ± 3.4	89.5 ± 5.3	0.03
Isoleucine	96.2 ± 1.9	83.6 ± 5.3	0.0007
Leucine	87.2 ± 1.4	89.0 ± 6.1	0.002
Lysine	78.2 ± 6.2	75.3 ± 10.1	0.3
Methionine	96.5 ± 3.3	92.6 ± 3.6	0.002
Phenylalanine	96.2 ± 2.4	85.2 ± 8.7	0.002
Threonine	93.4 ± 3.4	84.2 ± 7.2	0.002
Valine	95.2 ± 2.5	83.9 ± 8.1	0.001
Mean IAA	93.3 ± 6.7	84.1 ± 6.9	0.0009
DAA digestibility (%)			
Alanine	95.1 ± 2.3	88.5 ± 6.2	0.005
Asx	92.3 ± 4.1	82.4 ± 8.7	0.002
Glycine	91.0 ± 4.0	77.4 ± 10.4	0.003
Glx	95.0 ± 3.4	87.7 ± 7.3	0.007
Proline	92.6 ± 4.0	77.8 ± 9.1	0.0003
Serine	94.7 ± 2.7	82.4 ± 8.9	0.001
Tyrosine	94.8 ± 3.3	87.2 ± 6.3	0.002
Mean DAA	93.6 ± 2.5	82.7 ± 8.1	0.002
Mean of all AA ³	93.8 ± 3.1	83.9 ± 7.3	0.0008

AA, amino acid; Asx, aspartate and asparagine; DAA, dispensable amino acid; Glx, glutamate and glutamine; IAA, indispensable amino acid.

¹ Values are means \pm SDs. n = 9 in the flaxseed group and n = 10 in the rapeseed group. The digestibility of tryptophan, arginine, and cysteine was not determined.

² Differences between groups were estimated using an unpaired Student's *t* test. Considering the number of tests for AA digestibility (n = 15), the overall concentration to conclude to a significant difference for each AA was 0.05/15 = 0.0033 (Bonferroni correction). *P* values above this threshold are indicated in italics.

³ Means of AA digestibility are weighted by the proportion of each AA in the protein.

limiting AAs were revealed in rapeseed by a DIAAS higher than 1, in contrast to flaxseed.

We thank Francis Cacérès from Terres Inovia for their participation in the labeling of flaxseed and rapeseed and Alain Quinsac for its participation to the project management. We thank Mbalo Ndiaye from



FIGURE 3. Digestive (black bars) and deamination losses (urinary urea: light gray; body urea: medium gray) of dietary nitrogen 8 h after the ingestion of flaxseed or rapeseed protein cooked in biscuits. Values are means \pm SDs. n = 9 in flaxseed group and n = 10 in rapeseed group. The data were analyzed using Student's *t* test with unequal variances. * indicates a significant difference between the groups.

Avril for his involvement in the production of protein ingredients and biscuits. We also thank Marie Claude Amard and Bruna Neves from Avicenne Hospital for recruiting the subjects and collecting their blood samples.

Author contribution

The authors' responsibilities were as follows—CG, JC, OG, and DT: designed the research; RT, JC, NK, MC, GA, CG, and RB: conducted the research; SD and OG: provided essential materials; RT and CG: analyzed data; RT, CG, and JC: wrote the article; RT and CG: had primary responsibility for final content; and all authors: read, edited and approved the final manuscript.

Conflict of interest

The authors report no conflicts of interest.

Data availability

Data described in the manuscript will be made available uponrequest, pending application and approval.

 TABLE 3

 Chemical and DIAAS for flaxseed and rapeseed protein cooked in biscuits and calculated using 2 different nitrogen-to-protein conversion factors¹

Amino Acid	Chemical ratio	Chemical ratio				DIAAS ratio			
	N × 5.6		N × 6.25		N × 5.6		N × 6.25		
	Flaxseed	Rapeseed	Flaxseed	Rapeseed	Flaxseed	Rapeseed	Flaxseed	Rapeseed	
Histidine	1.8	2.3	1.6	2.1	1.7	2.1	1.5	1.9	
Isoleucine	2.0	1.8	1.8	1.6	2.0	1.5	1.8	1.4	
Leucine	1.3	1.6	1.1	1.2	1.2	1.4	1.1	1.3	
Lysine	0.8	1.6	0.7	1.5	0.7	1.2	0.6	1.1	
SAA	1.7	3.0	1.6	2.7	1.7	2.8	1.5	2.5	
AAA	2.5	2.3	2.3	2.0	2.4	1.9	2.2	1.7	
Threonine	1.8	2.3	1.6	2.0	1.7	1.9	1.5	1.7	
Tryptophan	3.5	3.1	3.1	2.8	_	_	_	_	
Valine	1.8	1.7	1.6	1.5	1.7	1.4	1.5	1.3	
SCORE ²	0.8 (Lys)	1.6 (Leu)	0.7 (Lys)	1.2 (Leu)	0.7 (Lys)	1.26 (Lys)	0.6 (Lys)	1.1 (Lys)	

AAA, aromatic amino acids (phenylalanine and tyrosine); DIAAS, digestible indispensable amino acid score; Leu, leucine; Lys, lysine; SAA, sulfur amino acids (methionine and cysteine).

¹ Values are calculated based on the mean amino acid composition of protein (Table 2) and mean ileal digestibility for the IAA (Table 3).

² The score is the chemical score for columns 1–4 and DIAAS for column 5 to 8. DIAAS corresponds to the lowest DIAA score, and the IAA concerned is shown in brackets.

R. Tessier et al.

Funding

The French Research Agency and FASO (the Oilseed Strategic Action Fund) co-funded our experimental costs. The PhD grant for Romain Tessier was co-funded by FASO and INRAE.

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.ajcnut.2023.02.020.

References

- B. Schauberger, H. Kato, T. Kato, D. Watanabe, P. Ciais, French crop yield, area and production data for ten staple crops from 1900 to 2018 at county resolution, Sci. Data 9 (2022) 38, https://doi.org/10.1038/s41597-022-01145-4.
- [2] FranceAgriMer database [Internet], VisioNet Données économiques agricoles et alimentaires. [cited, 2022 Jul 20]. Available from: https://visionet.franceagri mer.fr/Pages/Statistiques.aspx?menuurl=Statistiques/productions%20v%C3% A9g%C3%A9tales/grandes%20cultures/revue%20March%C3%A9%20des% 2001%C3%A9o-prot%C3%A9agineux/France.
- [3] R. Tessier, J. Calvez, G. Airinei, N. Khodorova, R. Kapel, A. Quinsac, et al., The true amino acid digestibility of 15N-labelled sunflower biscuits determined with ileal balance and dual isotope methods in healthy humans, J. Nutr. 152 (2022) 698–706, https://doi.org/10.1093/jn/nxab423.
- [4] C. Bos, G. Airinei, F. Mariotti, R. Benamouzig, S. Bérot, J. Evrard, et al., The poor digestibility of rapeseed protein is balanced by its very high metabolic utilization in humans, J. Nutr. 137 (2007) 594–600, https://doi.org/10.1093/jn/ 137.3.594.
- [5] F. Mariotti, S. Mahé, R. Benamouzig, C. Luengo, S. Daré, C. Gaudichon, et al., Nutritional value of [¹⁵N]-soy protein isolate assessed from ileal digestibility and postprandial protein utilization in humans, J. Nutr. 129 (1999) 1992–1997, https://doi.org/10.1093/jn/129.11.1992.
- [6] F. Mariotti, M.E. Pueyo, D. Tomé, S. Bérot, R. Benamouzig, S. Mahé, The influence of the albumin fraction on the bioavailability and postprandial utilization of pea protein given selectively to humans, J. Nutr. 131 (2001) 1706–1713.
- [7] C. Bos, B. Juillet, H. Fouillet, L. Turlan, S. Daré, C. Luengo, et al., Postprandial metabolic utilization of wheat protein in humans, Am. J. Clin. Nutr. 81 (2005) 87–94, https://doi.org/10.1093/ajcn/81.1.87.
- [8] FAO. Dietary protein quality evaluation in human nutrition. Report of an FAO Expert Consultation, Food and Nutrition Paper n°92, Rome, Italy, 2013, pp. 1–66.
- [9] C. Gaudichon, C. Bos, C. Morens, K.J. Petzke, F. Mariotti, J. Everwand, et al., Ileal losses of nitrogen and amino acids in humans and their importance to the assessment of amino acid requirements, Gastroenterology 123 (2002) 50–59, https://doi.org/10.1053/gast.2002.34233.
- [10] J. Calvez, S. Benoit, J. Piedcoq, N. Khodorova, D. Azzout-Marniche, D. Tomé, et al., Very low ileal nitrogen and amino acid digestibility of zein compared to whey protein isolate in healthy volunteers, Am. J. Clin. Nutr. 113 (2021) 70–82, https://doi.org/10.1093/ajcn/nqaa274.
- [11] F.M. Guillin, C. Gaudichon, L. Guérin-Deremaux, C. Lefranc-Millot, G. Airinei, N. Khodorova, et al., Real ileal amino acid digestibility of pea protein compared to casein in healthy humans: a randomized trial, Am. J. Clin. Nutr. 115 (2022) 353–363, https://doi.org/10.1093/ajcn/nqab354.
- [12] S. Devi, A. Varkey, M. Dharmar, R.R. Holt, L.H. Allen, M.S. Sheshshayee, et al., Amino acid digestibility of extruded chickpea and yellow pea protein is high and comparable in moderately stunted south Indian children with use of a dual stable isotope tracer method, J. Nutr. 150 (2020) 1178–1185, https:// doi.org/10.1093/jn/nxaa004.

- [13] M. Oberli, A. Marsset-Baglieri, G. Airinei, V. Santé-Lhoutellier, N. Khodorova, D. Rémond, et al., High true ileal digestibility but not postprandial utilization of nitrogen from bovine meat protein in humans is moderately decreased by high-temperature, long-duration cooking, J. Nutr. 145 (2015) 2221–2228.
- [14] R.W. Russell, A.D. Mc Gilliard, P.J. Berger, J.W. Young, Evaluation of turbidimetric determination of polyethylene glycol, J. Dairy Sci. 65 (1982) 1798–1803.
- [15] C. Fromentin, P. Sanders, F. Nau, M. Anton, G. Fromentin, D. Tomé, et al., A pilot study for the intrinsic labeling of egg proteins with ¹⁵N and ¹³C, Rapid Commun. Mass Spectrom. 26 (2012) 43–48, https://doi.org/10.1002/rcm.5291.
- [16] Afssa. Apport en protéines: consommation, qualité, besoins et recommandations, Report from Agence française de sécurité sanitaire des aliments, n° NUT-Ra-Proteines. Maison Alfort, France, 2007, pp. 1–461.
- [17] B. Cotten, D. Ragland, J.E. Thomson, O. Adeola, Amino acid digestibility of plant protein feed ingredients for growing pigs, J. Anim. Sci. 94 (2016) 1073–1082.
- [18] S. Giacomino, E. Peñas, V. Ferreyra, N. Pellegrino, M. Fournier, N. Apro, et al., Extruded flaxseed meal enhances the nutritional quality of cereal-based products, Plant Foods Hum. Nutr. 68 (2013) 131–136, https://doi.org/10.1007/ s11130-013-0359-8.
- [19] I. Schmidt, D. Renard, D. Rondeau, P. Richomme, Y. Popineau, M.A. Axelos, Detailed physicochemical characterization of the 2S storage protein from rape (*Brassica napus* L.), J. Agric. Food Chem. 52 (2004) 5995–6001, https:// doi.org/10.1021/jf0307954.
- [20] M. Das Purkayastha, J. Gogoi, D. Kalita, P. Chattopadhyay, K.S. Nakhuru, D. Goyary, et al., Physicochemical and functional properties of rapeseed protein isolate: influence of antinutrient removal with acidified organic solvents from rapeseed meal, J. Agric. Food Chem. 62 (2014) 7903–7914, https:// doi.org/10.1021/jf5023803.
- [21] C.M.E. Heyer, L.F. Wang, E. Beltranena, R.T. Zijlstra, Nutrient digestibility of extruded canola meal in ileal-cannulated growing pigs and effects of its feeding on diet nutrient digestibility and growth performance in weaned pigs, J. Anim. Sci. 99 (2021) 11, https://doi.org/10.1093/jas/skab135.
- [22] Y. Wang, J. Wang, S. Wang, J. Guo, S. Wang, Modification of glutenin and associated changes in digestibility due to methylglyoxal during heat processing, J. Agric. Food Chem. 67 (2019) 10734–10743, https://doi.org/10.1021/ acs.jafc.9b04337.
- [23] F. Mariotti, D. Tomé, P.P. Mirand, Converting nitrogen into protein—beyond 6.25 and Jones' factors, Crit. Rev. Food Sci. Nutr. 48 (2008) 177–184, https:// doi.org/10.1080/10408390701279749.
- [24] N. Sriperm, G.M. Pesti, P.B. Tillman, Evaluation of the fixed nitrogen-toprotein (N: P) conversion factor (6.25) versus ingredient specific N:P conversion factors in feedstuffs, J. Sci. Food Agric. 91 (2011) 1182–1186, https://doi.org/10.1002/jsfa.4292.
- [25] S.M. Rutherfurd, K. Bains, P.J. Moughan, Available lysine and digestible amino acid contents of proteinaceous foods of India, Br. J. Nutr. 108 (suppl 2) (2012) S59–S68, https://doi.org/10.1017/S0007114512002280.
- [26] A. Deglaire, P.J. Moughan, G. Airinei, R. Benamouzig, D. Tomé, Intact and hydrolyzed casein lead to similar ileal endogenous protein and amino acid flows in adult humans, Am. J. Clin. Nutr. 111 (2020) 90–97, https://doi.org/ 10.1093/ajcn/nqz268.
- [27] A. Deglaire, C. Fromentin, H. Fouillet, G. Airinei, C. Gaudichon, C. Boutry, et al., Hydrolyzed dietary casein as compared with the intact protein reduces postprandial peripheral, but not whole-body, uptake of nitrogen in humans, Am. J. Clin. Nutr. 90 (2009) 1011–1022, https://doi.org/10.3945/ ajcn.2009.27548.
- [28] H. Naci, What to do (or not to do) when randomization is not possible, J. Heart Lung Transplant. 36 (2017) 1174–1177, https://doi.org/10.1016/ j.healun.2017.07.008.